

3 different columns in one!

It's the new Whatman Multi-system - 3 different columns in one. How's it done? By using Whatman's unique interchangeable sealing pistons in one very durable glass column. Polypropylene pistons are fitted standard for ion exchange celluloses and most organic solvents, polyamide mesh is available for gel filtration and PTFE for difficult organic and acid solvents. So you don't have to buy lots of separate columns—just select the appropriate sealing piston. And as end units are identical, chromatography can be ascending or descending. Multi-system is made to the high standards of precision you'd expect from Whatman and incorporates their sample input/delivery system to ensure homogenous loading. Multi-system is designed for easy use, versatility and durability. It offers high accuracy in a full range of column chromatography systems without triplication—all at a reasonable cost.

New Whatman Multi-system— the cleverest column in chromatography!

For full details, contact—



Whatman®

WHATMAN BIOCHEMICALS LIMITED,
Springfield Mill, Maidstone, Kent, England.
Telephone: Maidstone 61688/9 Telex: 96113



Hoppe-Seyler's Zeitschrift für Physiologische Chemie

Begründet 1877 · Fortgeführt von A. Kossel, F. Knoop und K. Thomas

Editors in Chief:

A. BUTENANDT · F. LYNEN · G. WEITZEL

Subscription Rates:

For one volume (12 parts) DM 480,—

Vol. 355 No. 9

Contents

September 1974

Modification of histidine residue in yeast pyruvate kinase by diethylpyrocarbonate
L. Bornmann and B. Hess

Isolation of elastase-like and chymotrypsin-like neutral proteases from human granulocytes
W. Schmidt and K. Havemann

Chemistry and mechanism of new angiotensin II antagonists: 8-C-phenylglycine analogues of [5-isoleucine]angiotensin
H. Wissmann, B. Schoelkens, E. Lindner and R. Geiger

Fluorometric histamine determination in canine plasma under normal conditions, following application of exogenous histamine, and during histamine release by Haemaccel
W. Lorenz, H. Barth, M. Thermann, A. Schmal, P. Dormann and I. Niemeyer

An investigation of the microheterogeneity and the subunit structure of carboxylesterases by the quantitative determination of the terminal amino acids
E. Heymann and J. Iglesia

Membranes of glyoxysomes from castor bean endosperm; Further investigations on the membrane bound enzymes of fatty acid degradation and of the glyoxylate cycle
Ch. Biegelmayer, G. Nahler and H. Ruis

Enzymatic cleavage of the N^ε-acetyl protecting group from lysine in proteins (Kunitz trypsin-kallikrein inhibitor) in vitro and in vivo
H. Jering, G. Schorp and H. Tschesche

Binding of calcium ions by amino acids and peptides, especially by poly- α -L-glutamate
W. Dirscherl and H. Dohr

Activity of 8-C-bromo-derivatives of ATP and ADP in the mitochondrial adenine nucleotide translocation system
E. Schlimme and K.-W. Stahl

Excretion of isotopically labeled metabolites of testosterone in the bile of male rats
W.-D. Hetzel, R. Kiehscher and W. Staib

Influence of L-ascorbate on the concentrations microsomal cytochrome P-450 and cytochrome b₅ in adrenals, kidneys and spleen of guinea pigs
E. Degkwitz, S. Walsch and M. Dubberstein

Short Communications

Dependence of increased iron absorption by iron-deficient rats on an elutable component of jejunal mucosa
H. Huebers, E. Huebers and W. Rummel

Antamamide protects hepatocytes from phalloidin destruction
H. Faulstich, Th. Wieland, A. Walli and K. Birkmann

The primary structure of a monoclonal immunoglobulin L-chain of κ -type, subgroup IV (Bence-Jones protein Len.). A new subgroup of the κ -type L chain
M. Schneider and N. Hilschmann

Indexed in Current Contents



Walter de Gruyter · Berlin · New York



Walter de Gruyter

Berlin · New York

Neuausgabe

**Eckhart
Buddecke**

Grundriß der Biochemie

**Für Studierende der Medizin,
Zahnmedizin und Naturwissenschaften**

4., neubearbeitete Auflage. 1974.

Mit mehr als 400 Formeln, Tabellen und Diagrammen

Groß-Oktav. XXXII, 516 Seiten. Plastik flexibel DM 36,—
(de Gruyter Lehrbuch)

Die progressive Zunahme des biochemischen Fachwissens erfordert eine überschaubare und zusammenfassende Darstellung der Biochemie als Hilfsmittel für den Unterricht. Der Grundriß der Biochemie von E. Buddecke gliedert den Wissenschaft in die Kapitel „Stoffe und Stoffwechsel“, „Stoffwechselregulation“ und „Funktionelle Biochemie der Organe und Gewebe“ mit dem Ziel, durch knappe Darstellung gesicherte Fakten und gezielte Stoffauswahl dem Leser einerseits eine rasche Information zu bieten, andererseits jedoch auf die vielfältigen Beziehungen und Anwendungsmöglichkeiten der Biochemie zur klinischen Chemie und Molekularpathologie hinzuweisen, um damit dem Studierenden der Medizin und Zahnmedizin, sowie der anderen Gebiete der Naturwissenschaften das Verständnis klinischen Fachwissens zu erleichtern.

Neuerscheinung

**Curtius—Roth
(Editors)**

Clinical Biochemistry

Principles and Methods

2 Volumes

1974. Large-octavo. XXXVI, 1677 pages. With 177 tables, 362 figures, 3 coloured plates and 4463 references.
Cloth DM 460,— ISBN 3 11 001622 2

Special price until December, 31, 1974: DM 390,—

NEW 4 WAYS TO IMPROVE A GREAT SAMPLER

(WITH SAFETY IN MIND)


1 The design of the 'Model Q' is based on that of the established Oxford Sampler but features both forward and reverse modes of operation.

This gives the same degree of precision ($\pm 1\%$) with viscous and aqueous liquids.

2 Incorporation of a calibration stop halts the plunger action automatically to ensure precisely measured samples and even further elimination of human error.

3 A simple twist of the new release mechanism ejects the tip for disposal so you've no fear of personal contamination.

4 Available now for the 'Model Q' and other Oxford Samplers are individually wrapped sterile tips.



THE 'MODEL Q' SAMPLER



For further details please contact
The Boehringer Corporation (London) Ltd.,
Bilton House, Uxbridge Road, London W5 2TZ Telephone: 01-579 6944 (7 lines) Telex: 934289
If in the Republic of Ireland contact our Dublin Office at 5 Lower Hatch Street, Dublin, Tel: Dublin 760512

CLINICAL SCIENCE AND MOLECULAR MEDICINE

Formerly *Clinical Science*

EDITORIAL BOARD

For the Biochemical Society

J. A. OWEN, *Deputy Chairman*

B. H. BILLING, A. M. BRECKENRIDGE, C. N. HALES, T. D. R. HOCKADAY,
R. G. HUNTSMAN, G. H. LATHE, J. LIDDELL, I. MACINTYRE,
K. L. MANCHESTER, R. W. E. WATTS

For the Medical Research Society

R. D. COHEN, *Chairman*

G. CUMMING, D. C. FLENLEY, N. F. JONES, D. M. MATTHEWS,
M. E. M. NOBLE, D. K. PETERS, A. POLAK, J. I. S. ROBERTSON,
B. ROBINSON, J. D. M. SLATER

VOL. 47, No. 5

NOVEMBER 1974

CONTENTS

Neutral protease from the polymorphonuclear leucocytes of human rheumatoid synovial fluid.
By R. H. PRYCE-JONES, J. SAKLATVALA and G. C. WOOD

Unesterified fatty acids and the binding of tryptophan in human plasma. By G. CURZON,
JUNE FRIEDEL, BHARATI D. KATAMANENI, H. GREENWOOD and M. H. LADER

Influence of deoxycholic acid feeding on the elimination of cholesterol in normolipaemic subjects.
By K. EINARSSON, K. HELISTRÖM and M. KALLNER

Blood pressure changes produced by kidney cross-transplantation between spontaneously hypertensive rats and normotensive rats. By G. BIANCHI, U. FOX, G. F. DI FRANCESCO, A. M. GIOVANETTI and D. PAGETTI.

Vitamin E protection of the liver from paracetamol in the rat. By B. E. WALKER, J. KELLEHER
M. F. DIXON and M. S. LOSOWSKY

Studies of the peripheral vasodilator response to acute insulin-induced hypoglycaemia in man.
By W. G. MIDDLETON and E. B. FRENCH

The inhibition of renin secretion by alpha-adrenergic stimulation in the isolated rat kidney. By
R. VANDONGEN and W. S. PEART

The effects of exogenous parathyroid hormone on plasma and urinary adenosine 3',5'-cyclic monophosphate in man. By S. TOMLINSON, P. M. BARLING, J. D. M. ALBANO, B. L. BROWN
and J. L. H. O'RIORDAN

Muscle metabolism during exercise in patients with Parkinson's disease. By S. LANDIN,
L. HAGENFELDT, B. SALTIN and J. WAHREN

Platelet survival in acute proliferative glomerulonephritis. By J. A. CARRUTHERS, I. RALFS,
T. N. D. GIMLETTE and R. FINN

SHORT COMMUNICATION

Human erythrocyte glutathione reductase and glucose 6-phosphate dehydrogenase activities in
normal subjects and in persons exposed to lead. By J. K. HOWARD

MEDICAL RESEARCH SOCIETY (Meeting on 27 and 28 September 1974)

Demonstrations
Communications


*Subscription: £2.00 (\$7.50) per part; £20.00 (\$70.00) per year.
Orders may be placed with your bookseller, or sent direct to the publishers.*

BLACKWELL SCIENTIFIC PUBLICATIONS LTD

5 ALFRED STREET, OXFORD OX1 4HB, ENGLAND

BORIC ACID GELS

for the column chromatographic separation
of sugars and nucleic acids



18.445-4
Boric acid gel, particle size 0.1 - 0.4mm.
30ml \$75.00

18.518-3
Boric acid gel, particle size <0.1mm.
30ml \$150.00

Boric Acid Gels are cross-linked polymers insoluble in water and organic solvents. They are prepared by the cross-linking copolymerization of dihydroxyborylanilino-substituted methacrylic acid with 1,4-butanediol dimethacrylate.¹

These gels serve as useful packing material for the column chromatographic separation of mixtures whose components form complexes of varying stability with boric acid. Complex formation is dependent upon pH and molarity of the buffer used.

The separation of mono- and oligosaccharides,² ribonucleosides-deoxyribonucleosides,¹ oligonucleotides³ and t-RNA has been reported.

Boric Acid Gels may be used for preparative as well as analytical procedures, and are available in two porosities.

References

- 1) H. Schott, *Angew. Chem., Intern. Ed. Engl.*, **11**, 824 (1972); [*Angew. Chem.*, **84**, 819 (1972).]
- 2) K. Reske and H. Schott, *Angew. Chem., Intern. Ed. Engl.*, **12**, 417 (1973); [*Angew. Chem.*, **85**, 412 (1973).]
- 3) H. Schott, E. Rudloff, P. Schmidt, R. Roychoudhury and H. Kössel, *Biochemistry*, **12**, 932 (1973).

Do you have our free 1975 Catalog/Handbook?

It's THE SOURCE for more than 18,000 organic and biochemicals, polymers, and deuterated solvents, their disposal methods, handling precautions, physical constants, and reference data. And over 95% of the items listed are in stock, ready for immediate delivery. For your copy, send us your name, school or firm, and address.



Aldrich Chemical Company, Inc.

Craftsmen in Chemistry



Home Office:

Aldrich Chemical Co., Inc.
940 W. St. Paul Ave.
Milwaukee, Wisconsin 53233

In Great Britain:

Ralph N. Emanuel Ltd.
264 Water Rd., Wembley, Middx.
HAO 1PY, England

In Continental Europe:

Aldrich-Europe
B-2340 Beerse
Belgium

In Germany:

EGA-Chemie KG
7924 Steinheim am Albuch
Germany