



The hottest available

25-Hydroxy [23,24-³H] cholecalciferol
60-100Ci/ mmol TRK.558

L-[³⁵S]Methionine SJ.204

The specific activity of the last ten batches
has been 550-820Ci/mmol, with an
average of 627Ci/mmol.

Please enquire for further details



The Radiochemical Centre Amersham

The Radiochemical Centre Limited, Amersham, England.
Tel: Little Chalfont (024 04) 4444.

In the Americas: Amersham/Searle Corp, Illinois 60005.
Tel: 312-593-6300.

In W. Germany: Amersham Buchler GmbH & Co, KG,
Braunschweig. Tel: 05307-4693-97.

THE BIOCHEMICAL SOCIETY
SPECIAL PUBLICATIONS SERIES

No. 1 BIOCHEMISTRY AND MENTAL ILLNESS

Edited by L. L. IVERSEN *and* S. P. R. ROSE

1973 251 pp. £4.50 (\$12.00)

No. 2 BIOCHEMISTS IN INDUSTRY

Edited by K. W. FULLER

1973 58 pp. £1.25 (U.S. \$3.50)

No. 3 THE METABOLISM AND FUNCTION OF VITAMIN D

Edited by D. R. FRASER

1974 114 pp. £4.00 (\$10.50)

**No. 4 MEMBRANE ADENOSINE TRIPHOSPHATASES AND
TRANSPORT PROCESSES**

Edited by J. R. BRONK

1974 197 pp. £5.00 (\$12.50)

Available from your bookseller or agent, or direct from

THE BIOCHEMICAL SOCIETY (PUBLICATIONS)
P.O. Box 32, Commerce Way,
Colchester CO2 8HP, Essex

Leaflets giving details of contents of the above publications are available on request

Now Available...

VOLUME 2

ESSAYS IN MEDICAL BIOCHEMISTRY

Edited by V. MARKS and C. N. HALES

187 pp. £4.00 (US \$8.00) by surface mail to overseas addresses

Volume 2 of *Essays in Medical Biochemistry* continues the practice established in the first of providing, in readable form, highly authoritative but personal reviews of topics that are of general interest to medical biochemists. The subjects chosen for review are those that have undergone growth as a result of major conceptual and technological advances, and include essays by acknowledged world experts on complement, trace metals, acquired disorders of haem synthesis and chronobiochemistry.

List of Contents and Authors:

The Human Complement System: Biochemistry, Biology and Pathobiology

By D. T. Fearon and K. F. Austen

The Clinical Value of Trace-Metal Measurements

By H. T. Delves

Acquired Disorders of Haem Synthesis

By G. H. Elder

A New Perspective: Chronobiochemistry

By H. W. Simpson

Still Available:

ESSAYS IN MEDICAL BIOCHEMISTRY VOLUME 1

131 pp. £2.00 (US \$5.00)

Order From:




**THE BIOCHEMICAL SOCIETY BOOK DEPOT,
P.O. BOX 32,
COMMERCE WAY, COLCHESTER CO2 8HP, ENGLAND**
(or through your bookseller)



SYMPOSIA No. 41

BIOCHEMICAL ADAPTATION TO ENVIRONMENTAL CHANGE

Edited by **R. M. S. Smellie** and **J. F. Pennock**



240 pp. (ISBN 0 904498 01 8) £10.00 (U.S. \$20.00)

As biochemists have become more interested in the pathways of metabolism and mechanisms of control in a wider range of living systems, it has become apparent that, although there are many features in common to organisms inhabiting different environments, a variety of adaptations or refinements has been necessary so as to enable particular species to perform efficiently under the widely differing and often rapidly changing conditions to which they are exposed. Some of the more fully documented examples of these adaptations, including modifications in enzyme structure, function and activity, alteration in membrane lipids and performance and the substitution of one class of carrier molecule by another, are discussed in this volume.

List of contents and authors:

Chairman's Introduction by C. J. Duncan.

Design of Metabolic and Enzymic Machinery to Fit Lifestyle and Environment by P. W. Hochachka.

Temperature: A 'Shaping Force' in Protein Evolution by G. N. Somero and P. S. Low.

Temperature Adaptation in Fish by M. W. Smith.

Substrate Cycles in Metabolic Regulation and in Heat Generation by E. A. Newsholme and B. Crabtree.

Biochemical Adaptations for Flight in the Insect by B. Sacktor.

Facultative Anaerobiosis in Molluscs by A. de Zwaan, J. H. F. M. Kluytmans and D. I. Zandee.

Metabolic Consequences of Submersion Asphyxia in Mammals and Birds by A. S. Blix.

Adaptations with Respect to Salinity by E. Schoffeniels.

Adaptations of Enzymes for Regulation of Catalytic Function by D. E. Atkinson.

THE BIOCHEMICAL SOCIETY BOOK DEPOT

P.O. Box 32, Commerce Way, Colchester CO2 8HP, Essex.

CLINICAL SCIENCE AND MOLECULAR MEDICINE

Formerly *Clinical Science*

EDITORIAL BOARD

For the Biochemical Society

D. J. GALTON, *Deputy Chairman*

H. S. BACHELARD, B. H. BILLING, R. FRASER, D. GOMPERTZ,
D. A. HEMS, W. P. T. JAMES, D. E. M. LAWSON, B. LEWIS,
B. E. C. NORDIN, J. L. REID, M. SANDLER, P. G. WALKER

For the Medical Research Society

D. C. FLENLEY, *Chairman*

A. J. BELLINGHAM, I. R. CAMERON, F. J. CONWAY, P. T. FLUTE, P. J. HILTON,
D. B. MORGAN, M. I. M. NOBLE, P. RICHARDS, G. E. SLADEN,
J. D. SWALES, N. VEALL, M. M. WEBB-PEPLOE

Vol. 52, No. 5

May 1977

CONTENTS

- Editorial Review: Claude Bernard's (extended) *milieu interieur* revisited: autoregulation of cell and subcell integrity. By E. R. ROBIN
- Factors affecting angiotensin II concentrations in the human infant at birth. By FIONA BROUGHTON PIPKIN and E. M. SYMONDS
- Catabolic rate of α_1 -antitrypsin of Pi type M and Z in man. By C. B. LAURELL, B. NOSSLIN and J.-O. JEPSSON
- Porphyrins in erythrocytes and blood plasma: preliminary studies with qualitative thin-layer chromatography. By T. K. WITH
- Intrarenal distribution of plasma flow in cirrhosis as measured by transit renography: relationship with plasma renin activity, and sodium and water excretion. By S. P. WILKINSON, I. K. SMITH, M. CLARKE, V. ARROYO, J. RICHARDSON, H. MOODIE and R. WILLIAMS
- Plasma noradrenaline concentrations in experimental renovascular hypertension in the rat. By H. J. DARGIE, S. S. FRANKLIN and J. L. REID
- Protein metabolism in human neonates: nitrogen-balance studies, estimated obligatory losses of nitrogen and whole-body turnover of nitrogen. By P. B. PENCHARZ, W. P. STEFFEE, W. COCHRAN, N. S. SCRIMSHAW, W. M. RAND and V. R. YOUNG
- The effect of 25-hydroxy vitamin D₃ in the osteomalacia of chronic renal failure. By J. B. EASTWOOD, T. C. B. STAMP, H. E. DE WARDENER, P. J. BORDIER and C. D. ARNAUD
- Human hepatic 5-aminolaevulinate synthase: requirement of an exogenous system for succinyl-coenzyme A generation to demonstrate increased activity in cirrhotic and anticonvulsant-treated subjects. By H. L. BONKOWSKY and JOANNE S. POMEROY
- Bone necrosis and urinary hydroxyproline excretion in rabbits. By C. R. WEATHERLEY, G. DALE, J. MCGURK and D. N. WALDER
- Muscle-protein catabolism after injury in man, as measured by urinary excretion of 3-methyl-histidine. By D. H. WILLIAMSON, ROSE FARRELL, ALISON KERR and R. SMITH
- Proline and hydroxyproline excretion and vitamin C status in elderly human subjects. By C. J. BATES

SHORT COMMUNICATION

- Intrarenal role of angiotensin II in controlling sodium excretion during dehydration in dogs. By N. C. TRIPPEDO, J. E. HALL, T. E. LOHMEIER and A. C. GUYTON

The subscription price is £35.00 (U.K.), £35.00 (Overseas) and \$120.00 (North America) per annum, post free.


BLACKWELL SCIENTIFIC PUBLICATIONS LTD
OSNEY MEAD, OXFORD OX2 0EL, ENGLAND

Enzacryl[®] Gel



Packings for column and thin-layer permeation chromatography

Enzacryl Gel, a bead-polymerized chromatographic support for column and thin-layer gel permeation chromatography (GPC), is prepared by free-radical polymerization of *N*-acryloylmorpholine and *N,N'*-



methylenediacylamide in aqueous solution. The reaction mixture is dispersed in liquid paraffin containing suitable surfactants, and the resulting beads are washed and graded according to size.^{1,2,3}

Enzacryl Gel is available in five grades or porosities, designated K0, K1, K2, K4 and K10. The swelling properties of a given grade vary with solvent. Thus, although pore size is predetermined largely by the total monomer concentration and ratio of monomer to cross-linker in the aqueous polymerization mixture, useful secondary variations in mean pore size may be obtained as a result of swelling variation in different chromatographic solvents.³ **Enzacryl Gel K0** has the lowest mean pore size and is ideal for separation of small molecules in water, DMF and pyridine.

The most important advantage of **Enzacryl Gel** over other GPC supports is its utility in water, a wide range of organic solvents and an almost infinite variety of mixed-solvent systems. **Enzacryl Gel** easily withstands sterilization by heat. The matrix is relatively stable to

extremes of pH,⁴ and is not susceptible to microbial attack.

For details on the chemical and physical characteristics, applicability, molecular weight exclusion limits and other properties of **Enzacryl Gel**, please request free copies of *Aldrichimica Acta*, vol. 9 (ref. 3 below) and a technical data sheet.

References:

- 1) R. Epton, C. Holloway, and J.V. McLaren, *J. Appl. Polymer Sci.*, **18**, 179 (1974).
- 2) R. Epton, C. Holloway, and J.V. McLaren, *J. Chromatogr.*, **90**, 249 (1974).
- 3) R. Epton, *Aldrichimica Acta*, **9**, 23 (1976).
- 4) R. Epton, C. Holloway, and J.V. McLaren, *J. Chromatogr.*, **117**, 245 (1976).

Availability

Each of **Enzacryl Gel** grades K0, K1 and K2 is available in four bead sizes: extra fine (< 45 μ for thin-layer GPC), fine (40-75 μ), medium (75-150 μ) and coarse (150-300 μ). Please order by grade and bead size.

25g \$37.50 100g \$100.00

- 86,080-8 Enzacryl[®] Gel K4**, in pre-swollen form in water, for thin-layer GPC **25g \$32.00**
- 86,081-6 Enzacryl[®] Gel K10**, in pre-swollen form in water, for thin-layer GPC **25g \$33.70**
- 86,009-3 Enzacryl Gel[®] CO₂H Na** form, an acidic ion-exchange derivative of **Enzacryl Gel K2** in bead form, diameter 75-150 μ , exchange capacity 1.0-1.1 meq/g **10g \$15.00**
50g \$50.00

Please request information on our **Enzacryl[®] Carriers** and **Enzacryl[®] C Immobilized Enzymes**.

[®] Registered trademark of Koch-Light Laboratories, Colnbrook, Buckinghamshire, England. Aldrich is the exclusive distributor of **Enzacryl Products** in the U.S.

Aldrich Chemical Company, Inc.

Craftsmen in Chemistry

Corporate Offices:
Aldrich Chemical Co., Inc.
940 W. Saint Paul Ave.
Milwaukee, Wisconsin 53233
U. S. A.

Great Britain:
Aldrich Chemical Co., Ltd.
The Old Brickyard, New Road
Gillingham, Dorset SP8 4JL
England

Belgium/
Continental Europe:
Aldrich-Europe
B-2340 Beerse
Belgium

West Germany/
Continental Europe:
EGA-Chemie KG
7924 Steinheim am Albuch
West Germany