

Cellular Aspects

Contents of Volume 172, No. 2, 1978

FULL PAPERS

	PAGE		PAGE
Biosynthesis and Degradation			
An energy requirement for the degradation of intravenously injected ¹²⁵ I-labelled albumin in mouse liver and kidney slices. <i>By J. L. Mego & R. M. Farb</i>	233	Effect of nutrition on subcellular localization of rat fat-cell lipoprotein lipase. <i>By A. Vanhove, C. Wolf, M. Breton & M.-C. Glangeaud</i>	239
The simultaneous release by bone explants in culture and the parallel activation of pro-collagenase and of a latent neutral proteinase that degrades cartilage proteoglycans and denatured collagen <i>By G. Vaes, Y. Eeckhout, G. Lenaers-Claeys, C. Francois-Gillet & J.-E. Druetz</i>	261	Rapid stimulation by vasopressin, oxytocin and angiotensin II of glycogen degradation in hepatocyte suspensions. <i>By D. A. Hems, L. M. Rodrigues & P. D. Whitton</i>	311
Degradation of cartilage proteoglycans by a neutral proteinase secreted by rabbit bone-marrow macrophages in culture. <i>By P. Hauser & G. Vaes</i>	275	Effects of chronic hyperinsulinaemia on hepatic enzymes involved in lipogenesis and carbohydrate metabolism in the young rat. <i>By K. L. McCormick, J. A. Widness, J. B. Susa & R. Schwartz</i>	327
Calcium-dependent Golgi-vesicle fusion and cathepsin B in the conversion of proalbumin into albumin in rat liver. <i>By P. S. Quinn & J. D. Judah</i>	301	Is the adenine nucleotide translocator rate-limiting for oxidative phosphorylation? <i>By M. Stubbs, P. V. Vignais & H. A. Krebs</i>	333
Bioenergetics			
The properties of adenosine triphosphatase from exponential and synchronous cultures of <i>Alcaligenes eutrophus</i> H16. <i>By C. Edwards, J. A. Spode & C. W. Jones</i>	253	Development	
Cellular Interactions and Control Processes			
Rate-limiting factors in urate synthesis and gluconeogenesis in avian liver. <i>By J. P. Mapes & H. A. Krebs</i>	193	A radiochemical assay for glycine <i>N</i> -acyl-transferase activity. Some properties of the enzyme in rat and rabbit. <i>By M. O. James & J. R. Bend</i>	285
The contribution of serum triacylglycerol to hepatic triacylglycerol turnover in the starved rat. <i>By E. W. Lipkin, C. Cooper & R. A. Shipley</i>	205	Perinatal development of, and effect of chemical pretreatment on, glycine <i>N</i> -acyltransferase activities in liver and kidney of rabbit and rat. <i>By M. O. James & J. R. Bend</i>	293
Rate of release of hepatic triacylglycerol into serum in the starved rat. <i>By J. F. Palmer, C. Cooper & R. A. Shipley</i>	219	Changes in the lipoprotein lipase (clearing-factor lipase) activity of white adipose tissue during development of the rat. <i>By A. Cryer & H. M. Jones</i>	319
Thyrotropin-stimulated recruitment of free monoribosomes on to membrane-bound thyroglobulin-synthesizing polyribosomes. <i>By E. Davies, J. E. Dumont & G. Vassart</i>	227	Metabolism in Whole Organisms	
		The availability of inorganic sulphate in blood for sulphate conjugation of drugs in rat liver <i>in vivo</i> . [³⁵ S]Sulphate incorporation into harmol sulphate. <i>By G. J. Mulder & E. Scholtens</i>	247

RAPID PAPERS

	PAGE		PAGE
Subcellular Structures		Cellular Interactions and Control Processes	
Evidence that the coproporphyrinogen oxidase activity of rat liver is situated in the inter-membrane space of mitochondria. <i>By G. H. Elder & J. O. Evans</i>	345	Phospholipase A ₂ of rat liver mitochondria in vitamin E deficiency. <i>By A. S. Pappu, P. Fatterpaker & A. Sreenivasan</i>	349