

New labelled compounds from Amersham

We are continually extending our range of labelled compounds and radionuclides to maintain our position as the leading supplier of radiochemicals. A supplement is now available listing the 55 new products added to our range since the publication of our current radiochemicals catalogue 9 months ago.

Improvements in specifications and changes in availability are also detailed.

Write today for your copy of our supplement.



The Radiochemical Centre Amersham

The Radiochemical Centre Limited, Amersham, England.
Tel: Little Chalfont (024 04) 4444.

In the Americas: Amersham Corporation, Illinois 60005.
Tel: 312-593-6300.

In W. Germany: Amersham Buchler GmbH & Co, KG,
Braunschweig. Tel: 05307-4693-97.



Wiley

NEW YORK · CHICHESTER
BRISBANE · TORONTO

New Journal — Call For Papers

BIOPHARMACEUTICS & DRUG DISPOSITION

EDITORS

G.L. Mattok

Drug Research Laboratories,
Health Protection Branch,
Health and Welfare Canada
Tunney's Pasture,
Ottawa K1A 0L2,
Canada.

S.H. Curry

Department of Pharmacology and Therapeutics,
The London Hospital Medical College,
Turner Street,
London E1 2AD.

AIMS AND SCOPE

The Journal will publish original reports of studies in biopharmaceutics and pharmacokinetics, especially those which have a direct relation to the therapeutic use of drugs. This includes bioavailability and other human pharmacological studies, and therapeutic response and toxicity related to plasma and other tissue concentrations of drugs. Research on the design of drug dosage regimens, on factors affecting the disposition of and clinical response to drugs, and on the treatment of overdose, based on pharmacokinetic principles, will be accepted. Papers on analytical methodology, *in vitro* drug metabolism, and on animal models, will also be published, provided that they either facilitate the preceding types of investigation, or may otherwise be related to the use of drugs in man. The Journal will also carry review articles on specific subjects within the scope as defined above.

Starting Early 1979 — Initially Quarterly

FULL DETAILS AVAILABLE ON REQUEST FROM THE ADDRESS BELOW



John Wiley & Sons Limited

Baffins Lane · Chichester · Sussex PO19 1UD · England

now you can measure testosterone and dihydrotestosterone separately or together



Our new RIA kit is the simplest and fastest available for the assay of testosterone and dihydrotestosterone, separately or together in research applications in a wide range of biological samples.

To assay both androgens together a high specific activity tritium label gives the assay a very sensitive detection limit of 3 pg dihydrotestosterone per incubation tube. Also, a wide range of concentration levels can be assayed, up to 400 pg testosterone and 800 pg dihydrotestosterone per incubation tube.

Separation of the two androgens is achieved without the need for column chromatography by the use of a novel oxidation step which destroys the testosterone present. The assay of the unaffected dihydrotestosterone is achieved with a simple protocol incorporating a short incubation time of 2½ hours. Standards for both androgens are provided. Accurate results are obtained easily and quickly.



**The Radiochemical Centre
Amersham**


**new
testosterone/dihydrotestosterone RIA kit**

For further information please write or phone
The Radiochemical Centre Limited, Amersham, England. Telephone: 024-04-4444
In W. Germany: Amersham Buchler GmbH & Co KG, Braunschweig. Telephone: 05307-4693-97

If your isolated cells look like this?



or your nucleic acid like this?



You should use one of the range of enzymes from BCL prepared specifically to help you prepare **intact cells** and **high molecular weight nucleic acid**.

Item	Cat. No.	Use
Cellulase	238104	Preparation of protoplast, from plant cells.
Collagenase	103578	Disaggregation of liver cells, islets adrenal cells-etc.
Dispase (Protease neutral)	165859	Specifically for preparation of cells in culture.
Proteinase K	161519	When you need high molecular weight nucleic acid free from nuclease damage.
Pronase	165921	Broad spectrum protease for preparation of nucleic acid and isolated animal cells.



For full details of the above enzymes, please contact our Technical Department. To place your order contact:—
The Boehringer Corporation (London) Ltd.,
Bell Lane, Lewes, East Sussex BN7 1LG.
Telephone: Lewes (07916) 71611

22/78