

Molecular Aspects

Contents of Volume 185, No. 3, 1980

Proteins

| | | |
|--|---|---------|
| The chemical composition of bovine vitreous-humour collagen fibres | D. A. SWANN & S. S. SOTMAN | 545-554 |
| Messenger ribonucleic acids from pig intestinal mucosa direct synthesis of calcium-binding protein in a cell-free translation system | H. MELLERSH, S. TOMLINSON & A. POLLOCK | 601-607 |
| Amino acid sequences C-terminal to the cross-links in bovine elastin | K. M. BAIG, M. VLAOVIC & R. A. ANWAR | 611-616 |
| A biochemical comparison of normal human liver and hepatocellular carcinoma ferritins | S. BULLOCK, A. BOMFORD & R. WILLIAMS | 639-645 |
| Purification of <i>Torpedo californica</i> post-synaptic membranes and fractionation of their constituent proteins | J. ELLIOTT, S. G. BLANCHARD, W. WU, J. MILLER, C. D. STRADER, P. HARTIG, H.-P. MOORE, J. RACS & M. A. RAFTERY | 667-677 |
| Unoccupied nuclear oestradiol-receptor sites in normal human endometrium | C. LÉVY, R. MORTEL, B. EYCHENNE, P. ROBEL & E.-E. BAULIEU | 733-738 |
| The preparation of calmodulins from barley (<i>Hordeum</i> sp.) and Basidiomycete fungi | R. J. A. GRAND, A. C. NAIRN & S. V. PERRY | 755-760 |
| Poly(ADP-ribose) polymerase from Ehrlich ascites-tumour cells. Amino acid composition, N-terminal analysis and chemical cleavage of the purified protein | J. HOLTLUND, T. KRISTENSEN & K. SLETTEN | 779-782 |

Nucleic Acids

| | | |
|--|--------------------------|---------|
| Alkylation by 1,2:5,6-dianhydrogalactitol of deoxyribonucleic acid and guanosine | E. INSTITÓRIS & J. TAMÁS | 659-666 |
|--|--------------------------|---------|

Lipids

| | | |
|---|------------------------------|---------|
| Ubiquinones have surface-active properties suited to transport electrons and protons across membranes | P. J. QUINN & M. A. ESFAHANI | 715-722 |
| Enzymic hydrolysis of sphingomyelin in the presence of bile salts | S. YEDGAR & S. GATT | 749-754 |

Carbohydrates

| | | |
|---|-----------------------------|---------|
| Extraction and characterization of proteoglycan from human meniscus | D. MCNICOL & P. J. ROUGHLEY | 705-713 |
|---|-----------------------------|---------|

Small Molecules

| | | |
|---|----------------------------------|---------|
| Isolation, characterization and seasonal variations in the concentration of <i>N</i> ^ε (γ -glutamyl)-lysine isodipeptide in the blood plasma of the winter flounder (<i>Pseudopleuronectes americanus</i>) | E. J. SQUIRES & L. A. W. FELTHAM | 761-766 |
|---|----------------------------------|---------|

Enzymology

| | | |
|--|---|---------|
| An adenosine triphosphate-dependent carbamoylphosphate-3-hydroxymethylcephem- <i>O</i> -carbamoyltransferase from <i>Streptomyces clavuligerus</i> | S. J. BREWER, P. M. TAYLOR & M. K. TURNER | 555-564 |
| The isolation and preliminary characterization of <i>N</i> -acetyl-D-glucosamine kinase from rat kidney and liver | M. B. ALLEN & D. G. WALKER | 565-575 |
| Kinetic characterization of <i>N</i> -acetyl-D-glucosamine kinase from rat liver and kidney | M. B. ALLEN & D. G. WALKER | 577-582 |
| Interaction of activating protein and surfactants with human liver hexosaminidase A and G _{M2} ganglioside | P. HECHTMAN & K. KACHRA | 583-591 |

| | | |
|--|--|----------------|
| The use of arginine analogues for investigating the functional organization of the arginine-binding site in lobster muscle arginine kinase. Role of the 'essential' thiol group | D. C. WATTS, E. O. ANOSIKE, B. MORELAND, R. J. POLLITT & C. R. LEE | 593-599 |
| Graphical determination of pK values of the active-site groups of enzymes. An analysis of the bell-shaped curves | O. PARKASH & I. S. BHATIA | 609-610 |
| Comparison of bone and osteosarcoma adenylate cyclase. Partial purification of membranes and kinetic properties of enzyme | S. B. RODAN, J. J. EGAN, E. E. GOLUB & G. A. RODAN | 617-627 |
| Comparison of bone and osteosarcoma adenylate cyclase. Effects of Mg^{2+} , Ca^{2+} , ATP^{4-} and $HATP^{3-}$ in the assay mixture | S. B. RODAN, E. E. GOLUB, J. J. EGAN & G. A. RODAN | 629-637 |
| Pathways in the activation of human coagulation factor X | K. MERTENS & R. M. BERTINA | 647-658 |
| Properties of freshly purified and thiol-treated spinach chloroplast fructose biphosphatase | S. A. CHARLES & B. HALLIWELL | 689-693 |
| Venom from the snake <i>Bothrops asper</i> Garman. Purification and characterization of three phospholipases A_2 | A. C. ALAGÓN, R. R. MOLINAR, L. D. POSSANI, P. L. FLETCHER, JR., J. E. CRONAN, JR. & J. Z. JULIA | 695-704 |
| Influence of the position of the double bond in steroid substrates on the efficiency of the proton-transfer reaction by <i>Pseudomonas testosteroni</i> 3-oxo steroid Δ^4 - Δ^5 -isomerase | H. WEINTRAUB, E.-E. BAULIEU & A. ALFSEN | 723-732 |
| Reactivation and aging of neurotoxic esterase inhibited by a variety of organophosphorus esters | B. CLOTHIER & M. K. JOHNSON | 739-747 |
| Rapid type 2 molybdenum(V) electron-paramagnetic resonance signals from xanthine oxidase and the structure of the active centre of the enzyme | J. P. G. MALTHOUSE, S. GUTTERIDGE & R. C. BRAY | 767-770 |
| Kinetics of suicide substrates | S. G. WALEY | 771-773 |
| Novel inhibitors of poly(ADP-ribose) synthetase | M. R. PURNELL & W. J. D. WHISH | 775-777 |
| Maleylacetate reductase from <i>Trichosporon cutaneum</i> | A. B. GAAL & H. Y. NEUJAHN | 783-786 |
| Organization of thiol groups of electric-eel electric-organ sodium-plus-potassium ion-stimulated adenosine triphosphatase studied with bifunctional reagents | W. E. HARRIS & W. L. STAHL | 787-790 |
| General Methods | | |
| The development of a radioimmunoassay for Tamm-Horsfall glycoprotein in serum | A. DAWNAY, C. MCLEAN & W. R. CATTELL | 679-687 |

INDEXES

| | |
|----------------------|----------------|
| Author Index | 791-792 |
| Subject Index | 793-798 |

Contents of Volume 185, No. 1

Proteins

| | | |
|--|---|---------|
| Fibrinogen–fibrin conversion. The mechanism of fibrin-polymer formation in solution | G. F. SMITH | 1–11 |
| Chemical modification of the haem propionate of cytochrome <i>c</i> | R. TIMKOVICH | 47–57 |
| Cholic acid binding by glutathione <i>S</i> -transferases from rat liver cytosol | J. D. HAYES, R. C. STRANGE & I. W. PERCY-ROBB | 83–87 |
| Comparison of phosphorylation of ribosomal proteins from HeLa and Krebs II ascites-tumour cells by cyclic AMP-dependent and cyclic GMP-dependent protein kinases | O.-G. ISSINGER, H. BEIER, N. SPEICHERMANN, V. FLOKERZI & F. HOFMANN | 89–99 |
| Equilibrium-binding studies of pig laryngeal cartilage proteoglycans with hyaluronate oligosaccharide fractions | I. A. NIEDUSZYNSKI, J. K. SHEEHAN, C. F. PHELPS, T. E. HARDINGHAM & H. MUIR | 107–114 |
| Immuno-biochemical studies of a non-histone chromosomal protein in embryonic and mature chick oviduct | C. T. TENG & C. S. TENG | 169–175 |
| Identification of cyanogen bromide peptides involved in intermolecular cross-linking of bovine type III collagen | A. C. NICHOLLS & A. J. BAILEY | 195–201 |
| Purification and properties of different forms of modeccin, the toxin of <i>Adenia digitata</i> . Separation of subunits with inhibitory and lectin activity | L. BARBIERI, M. ZAMBONI, L. MONTANARO, S. SPERTI & F. STIRPE | 203–210 |
| Sensitivity and other factors affecting biospecific desorption in chromatography of proteins. A study by computer simulation | R. J. YON | 211–216 |
| Reaction of 1-fluoro-2,4-dinitrobenzene and 2,4,6-trinitrobenzenesulphonic acid with membranes of sarcoplasmic reticulum | D. A. THORLEY-LAWSON & N. M. GREEN | 223–226 |
| Purification and characterization of mouse α -lactalbumin and preparation of its antibody | Y. NAGAMATSU & T. OKA | 227–237 |
| The amino acid sequence of ferredoxin from <i>Brassica napus</i> (rape) | I. TAKRURI & D. BOULTER | 239–243 |
| Labelling of the cyclitol permease in <i>Klebsiella aerogenes</i> | G. REBER, M. ROPARS & J. DESHUSSES | 253–257 |
| Properties of the non-specific calcium-binding sites of rabbit skeletal-muscle myosin | J. WIKMAN-COFFELT | 265–268 |
| Purification and some properties of a lectin from the fruit juice of the tomato (<i>Lycopersicon esculentum</i>) | D. C. KILPATRICK | 269–272 |
| Haemoglobin binding with haptoglobin. Unequivocal demonstration that the β -chains of human haemoglobin bind to haptoglobin | A. L. KAZIM & M. Z. ATASSI | 285–287 |

Nucleic Acids

| | | |
|---|------------------------------------|-------|
| The isolation and characterization of a sequence-specific endonuclease from <i>Anabaena subcylindrica</i> | S. G. HUGHES, T. BRUCE & K. MURRAY | 59–63 |
| The nucleotide sequences recognized by endonucleases <i>Ava</i> I and <i>Ava</i> II from <i>Anabaena variabilis</i> | S. G. HUGHES & K. MURRAY | 65–75 |

Lipids

| | | |
|---|-------------------------|-------|
| Lipopolysaccharides of R mutants isolated from <i>Vibria cholerae</i> | K. HISATSUNE & S. KONDO | 77–81 |
|---|-------------------------|-------|

Carbohydrates

| | | |
|--|--------------------------|-------|
| Factors influencing proteoglycan size in rachitic-chick growth cartilage | P. ROUGHLEY & I. DICKSON | 33–39 |
|--|--------------------------|-------|

| | | |
|--|---|---------|
| Quantitative analysis of hyaluronate in nanogram amounts | A. TENGBLAD | 101–105 |
| The glycoprotein of measles virus. External radioactive labelling of its carbohydrate and partial characterization of the glycopeptide | O. ANTONEN, M. JOKINEN, A. SALMI, R. VAINIONPÄ & C. G. GAHMBERG | 189–194 |
| Small Molecules | | |
| Metabolism of [¹⁴ C]diethylstilboestrol epoxide by rat liver <i>in vitro</i> | P. H. JELLINCK & J. H. BOWEN | 129–137 |
| An accurate and sensitive analysis by high-pressure liquid chromatography of conjugated and unconjugated bilirubin IX- α in various biological fluids | S. ONISHI, S. ITOH, N. KAWADE, K. ISOBE & S. SUGIYAMA | 281–284 |
| Enzymology | | |
| Specificity of P2 primary alkylsulphohydrolase induction in the detergent-degrading bacterium <i>Pseudomonas</i> C12B. Effects of alkanesulphonates, alkyl sulphates and other related compounds | J. M. CLOVES, K. S. DODGSON, G. F. WHITE & J. W. FITZGERALD | 13–21 |
| Purification and properties of the P2 primary alkylsulphohydrolase of the detergent-degrading bacterium <i>Pseudomonas</i> C12B | J. M. CLOVES, K. S. DODGSON, G. F. WHITE & J. W. FITZGERALD | 23–31 |
| The binding of the coenzyme pyridoxal 5'-phosphate and analogues of the substrate-coenzyme complex to tyrosine decarboxylase | A. ORLACCHIO, C. BORRI-VOLTATTORNI & C. TURANO | 41–46 |
| Analysis of bilirubin and bilirubin mono- and di-conjugates. Determination of their relative amounts in biological samples | N. BLANCKAERT | 115–128 |
| Characterization of the low-temperature intermediates of the reaction of fully reduced soluble cytochrome oxidase with oxygen by electron-paramagnetic-resonance and optical spectroscopy | G. M. CLORE, L.-E. ANDRÉASSON, B. KARLSSON, R. AASA & B. G. MALMSTRÖM | 139–154 |
| Characterization of the intermediates in the reaction of mixed-valence-state soluble cytochrome oxidase with oxygen at low temperatures by optical and electron-paramagnetic-resonance spectroscopy | G. M. CLORE, L.-E. ANDRÉASSON, B. KARLSSON, R. AASA & B. G. MALMSTRÖM | 155–167 |
| Inhibition of β -lactamase of <i>Bacillus licheniformis</i> 749/C by compound PS-5, a new β -lactam antibiotic | Y. FUKAGAWA, T. TAKEI & T. ISHIKURA | 177–185 |
| <i>Appendix:</i> A computer-assisted spectrophotometric enzyme assay. A method for the calculation of K_m and V from a single reaction progress curve by the integrated form of the Lineweaver–Burk plot with particular reference to β -lactamase | Y. FUKAGAWA | 186–188 |
| A kinetic method for the study of solvent environments of thiol groups in proteins involving the use of a pair of isomeric reactivity probes and a differential solvent effect. Investigation of the active centre of ficin by using 2,2'- and 4,4'-dipyridyl disulphides as reactivity probes | J. P. G. MALTHOUSE & K. BROCKLEHURST | 217–222 |
| Investigation of human erythrocyte superoxide dismutase by ¹ H nuclear-magnetic-resonance spectroscopy | H. A. O. HILL, W.-K. LEE, J. V. BANNISTER & W. H. BANNISTER | 245–252 |
| Demonstration and cytosolic location of an endo- <i>N</i> -acetyl- β -D-glucosaminidase activity towards an asialo- <i>N</i> -acetyl-lactosaminic-type substrate in rat liver | R. J. PIERCE, G. SPIK & J. MONTREUIL | 261–264 |
| Electrophoretically distinct forms of uridine kinase in the rat. Tissue distribution and age-dependence | J. ABSIL, M. TUILIE & J.-M. ROUX | 273–276 |
| Intermolecular Organization | | |
| DNA-binding specificity of a chromatin non-histone protein fraction of HeLa cells | J. H. J. DUNN, R. M. LYALL, R. C. BRIGGS, A. M. CAMPBELL & S. HNILICA | 277–279 |

Contents of Volume 185, No. 2

Proteins

- | | | |
|--|---|---------|
| Granulocyte/macrophage-, megakaryocyte-, eosinophil- and erythroid-colony-stimulating factors produced by mouse spleen cells | A. W. BURGESS, D. METCALF, S. H. M. RUSSELL & N. A. NICOLA | 301-314 |
| The organization of NADH dehydrogenase polypeptides in the inner mitochondrial membrane | S. SMITH & C. I. RAGAN | 315-326 |
| Isolation of a serum component that stimulates the spreading of cells in culture | J. G. WHATELEY & P. KNOX | 349-354 |
| The chemistry of the collagen cross-links. Purification and characterization of cross-linked polymeric peptide material from mature collagen containing unknown amino acids | N. D. LIGHT & A. J. BAILEY | 373-381 |
| The isolation, characterization and partial sequences of the chicken erythrocyte non-histone chromosomal proteins HMG14 and HMG17. Comparison with the homologous calf thymus proteins | J. M. WALKER & E. W. JOHNS | 383-386 |
| Sequence of the N-terminal half of <i>Bacillus amyloliquefaciens</i> α -amylase | H. CHUNG & F. FRIEDBERG | 387-395 |
| A low-molecular-weight inhibitor of the neutral proteinase from rat intestinal smooth muscle | I. T. CARNEY, C. G. CURTIS, J. KAY & N. BIRKET | 423-433 |
| Precolicin E1, the major gene product of plasmid-ColE1 deoxyribonucleic acid <i>in vitro</i> | D. H. WATSON | 463-471 |
| The distribution of iron between the metal-binding sites of transferrin in human serum | J. WILLIAMS & K. MORETON | 483-488 |
| Isoelectric-focusing properties and carbohydrate content of pea (<i>Pisum sativum</i>) legumin | J. A. GATEHOUSE, R. R. D. CROY & D. BOULTER | 497-503 |
| An investigation of pyridinoline, a putative collagen cross-link | D. F. ELSDEN, N. D. LIGHT & A. J. BAILEY | 531-534 |

Nucleic Acids

- | | | |
|---|--|---------|
| Isolation and preliminary characterization of epithelial-specific messenger ribonucleic acids and their products during embryonic tooth development | M. ZEICHNER-DAVID, B. G. WELIKY & H. C. SLAVKIN | 489-496 |
|---|--|---------|

Carbohydrates

- | | | |
|---|---|---------|
| Partial deglycosylation of blood-group-specific glycoproteins | A. S. R. DONALD | 327-337 |
| Isolation and characterization of sulphated mucopolysaccharides from rat leukaemic (RBL-1) basophils | D. D. METCALFE, S. I. WASSERMAN & K. F. AUSTEN | 367-372 |
| The relationship between acetyl content and sugar composition in expanding leaves of beech (<i>Fagus sylvatica</i> L.) | J. S. D. BACON | 405-409 |

Small Molecules

- | | | |
|--|---|---------|
| Identification of <i>O</i> -acetyl-5-methoxytryptophol in the pineal gland by gas-chromatography-mass spectrometry | I. SMITH, P. FRANCIS, R. M. LEONE & P. E. MULLEN | 537-540 |
|--|---|---------|

Enzymology

- | | | |
|---|--|---------|
| A steady-state kinetic analysis of the fructose 1,6-bisphosphate-activated pyruvate kinase from <i>Carcinus maenas</i> hepatopancreas | I. G. GILES & P. C. POAT | 289-299 |
| New relations for activity analysis of associating enzymes | J. C. THOMES, J. ARCHAMBAULT DE VENCAY & R. JULIEN | 339-347 |

| | | |
|--|---|---------|
| Calcium ion-regulated thin filaments from vascular smooth muscle | S. B. MARSTON, R. M. TREVETT & M. WALTERS | 355-365 |
| Electron-paramagnetic-resonance parameters of molybdenum(V) in sulphite oxidase from chicken liver | M. T. LAMY, S. GUTTERIDGE & R. C. BRAY | 397-403 |
| Mechanistic and stereochemical studies on 3-oxo steroid Δ^4 - Δ^5 -isomerase from human placenta | M. AKHTAR, M. CALDER, T. SMITH & J. N. WRIGHT | 411-421 |
| 3-Hydroxy-3-methylglutaryl-coenzyme A reductase. A comparison of the modulation <i>in vitro</i> by phosphorylation and dephosphorylation to modulation of enzyme activity by feeding cholesterol- or cholestyramine-supplemented diets | K. A. MITROPOULOS, B. L. KNIGHT & B. E. A. REEVES | 435-441 |
| Partial purification and characterization of a novel neutral proteinase from human uterine cervix | A. ITO, H. IHARA & Y. MORI | 443-450 |
| Activation and inactivation of phosphoenolpyruvate carboxykinase by ferrous ions | C. H. REYNOLDS | 451-454 |
| Purification and properties of α -D-mannosidase from the germinated seeds of <i>Medicago sativa</i> (alfalfa) | A. CURDEL & F. PETEK | 455-462 |
| Active-site amino acid residues in γ -glutamyltransferase and the nature of the γ -glutamyl-enzyme bond | J. S. ELCE | 473-481 |
| The site of phosphorylation of troponin I in the perfused rabbit heart. The effect of adrenaline | A. J. G. MOIR, R. J. SOLARO & S. V. PERRY | 505-513 |
| Partial purification and characterization of a new intracellular β -glucosidase of <i>Trichoderma reesei</i> | M. INGLIN, B. A. FEINBERG & J. R. LOEWENBERG | 515-519 |
| The phospholipid-dependence of uridine diphosphate glucuronyltransferase. Temperature-dependence of microsomal enzyme activity (reverse reaction) | J. CUMMINGS, A. B. GRAHAM & G. C. WOOD | 521-526 |
| Oxygen kinetics of frozen cytochrome oxidase. The capacity of the oxygen pocket | K. DE FONSEKA & B. CHANCE | 527-530 |
| Validity of the jack-knife technique for analysing enzyme-kinetic data | A. CORNISH-BOWDEN & J. T.-F. WONG | 535-536 |
| Isoenzymes of membrane-bound β -glucosidase of human spleen | M. YAQOUB & M. CARROLL | 541-543 |

Contents of Volume 185, No. 3

Proteins

| | | |
|--|---|---------|
| The chemical composition of bovine vitreous-humour collagen fibres | D. A. SWANN & S. S. SOTMAN | 545-554 |
| Messenger ribonucleic acids from pig intestinal mucosa direct synthesis of calcium-binding protein in a cell-free translation system | H. MELLERSH, S. TOMLINSON & A. POLLOCK | 601-607 |
| Amino acid sequences C-terminal to the cross-links in bovine elastin | K. M. BAIG, M. VLAOVIC & R. A. ANWAR | 611-616 |
| A biochemical comparison of normal human liver and hepatocellular carcinoma ferritins | S. BULLOCK, A. BOMFORD & R. WILLIAMS | 639-645 |
| Purification of <i>Torpedo californica</i> post-synaptic membranes and fractionation of their constituent proteins | J. ELLIOTT, S. G. BLANCHARD, W. WU, J. MILLER, C. D. STRADER, P. HARTIG, H.-P. MOORE, J. RACS & M. A. RAFTERY | 667-677 |
| Unoccupied nuclear oestradiol-receptor sites in normal human endometrium | C. LÉVY, R. MORTEL, B. EYCHENNE, P. ROBEL & E.-E. BAULIEU | 733-738 |
| The preparation of calmodulins from barley (<i>Hordeum</i> sp.) and Basidiomycete fungi | R. J. A. GRAND, A. C. NAIRN & S. V. PERRY | 755-760 |

| | | |
|--|--|---------|
| Poly(ADP-ribose) polymerase from Ehrlich ascites-tumour cells. Amino acid composition, <i>N</i> -terminal analysis and chemical cleavage of the purified protein | J. HOLTUN, T. KRISTENSEN & K. SLETTEN | 779-782 |
| Nucleic Acids | | |
| Alkylation by 1,2:5,6-dianhydrogalactitol of deoxyribo-nucleic acid and guanosine | E. INSTITÓRIS & J. TAMÁS | 659-666 |
| Lipids | | |
| Ubiquonones have surface-active properties suited to transport electrons and protons across membranes | P. J. QUINN & M. A. ESFAHANI | 715-722 |
| Enzymic hydrolysis of sphingomyelin in the presence of bile salts | S. YEDGAR & S. GATT | 749-754 |
| Carbohydrates | | |
| Extraction and characterization of proteoglycan from human meniscus | D. MCNICOL & P. J. ROUGHLEY | 705-713 |
| Small Molecules | | |
| Isolation, characterization and seasonal variations in the concentration of <i>N</i> ^ε -(γ -glutamyl)-lysine isodipeptide in the blood plasma of the winter flounder (<i>Pseudopleuronectes americanus</i>) | E. J. SQUIRES & L. A. W. FELTHAM | 761-766 |
| Enzymology | | |
| An adenosine triphosphate-dependent carbamoylphosphate-3-hydroxymethylcephem <i>O</i> -carbamoyltransferase from <i>Streptomyces clavuligerus</i> | S. J. BREWER, P. M. TAYLOR & M. K. TURNER | 555-564 |
| The isolation and preliminary characterization of <i>N</i> -acetyl-D-glucosamine kinase from rat kidney and liver | M. B. ALLEN & D. G. WALKER | 565-575 |
| Kinetic characterization of <i>N</i> -acetyl-D-glucosamine kinase from rat liver and kidney | M. B. ALLEN & D. G. WALKER | 577-582 |
| Interaction of activating protein and surfactants with human liver hexosaminidase A and G _{M2} ganglioside | P. HECHTMAN & K. KACHRA | 583-591 |
| The use of arginine analogues for investigating the functional organization of the arginine-binding site in lobster muscle arginine kinase. Role of the 'essential' thiol group | D. C. WATTS, E. O. ANOSIKE, B. MORELAND, R. J. POLLITT & C. R. LEE | 593-599 |
| Graphical determination of <i>pK</i> values of the active-site groups of enzymes. An analysis of the bell-shaped curves | O. PARKASH & I. S. BHATIA | 609-610 |
| Comparison of bone and osteosarcoma adenylate cyclase. Partial purification of membranes and kinetic properties of enzyme | S. B. RODAN, J. J. EGAN, E. E. GOLUB & G. A. RODAN | 617-627 |
| Comparison of bone and osteosarcoma adenylate cyclase. Effects of Mg ²⁺ , Ca ²⁺ , ATP ⁴⁻ and HATP ³⁻ in the assay mixture | S. B. RODAN, E. E. GOLUB, J. J. EGAN & G. A. RODAN | 629-637 |
| Pathways in the activation of human coagulation factor X | K. MERTENS & R. M. BERTINA | 647-658 |
| Properties of freshly purified and thiol-treated spinach chloroplast fructose bisphosphatase | S. A. CHARLES & B. HALLIWELL | 689-693 |
| Venom from the snake <i>Bothrops asper</i> Garman. Purification and characterization of three phospholipases A ₂ | A. C. ALAGÓN, R. R. MOLINAR, L. D. POSSANI, P. L. FLETCHER, JR., J. E. CRONAN, JR. & J. Z. JULIA | 695-704 |
| Influence of the position of the double bond in steroid substrates on the efficiency of the proton-transfer reaction by <i>Pseudomonas testosteroni</i> 3-oxo steroid Δ^4 - Δ^5 -isomerase | H. WEINTRAUB, E.-E. BAULIEU & A. ALFSEN | 723-732 |
| Reactivation and aging of neurotoxic esterase inhibited by a variety of organophosphorus esters | B. CLOTHIER & M. K. JOHNSON | 739-747 |

| | | |
|--|--|---------|
| Rapid type 2 molybdenum(V) electron-paramagnetic resonance signals from xanthine oxidase and the structure of the active centre of the enzyme | J. P. G. MALTHOUSE, S. GUTTERIDGE & R. C. BRAY | 767-770 |
| Kinetics of suicide substrates | S. G. WALEY | 771-773 |
| Novel inhibitors of poly(ADP-ribose) synthetase | M. R. PURNELL & W. J. D. WHISH | 775-777 |
| Maleylacetate reductase from <i>Trichosporon cutaneum</i> | A. B. GAAL & H. Y. NEUJAHN | 783-786 |
| Organization of thiol groups of electric-eel electric-organ sodium-plus-potassium ion-stimulated adenosine triphosphatase studied with bifunctional reagents | W. E. HARRIS & W. L. STAHL | 787-790 |
| General Methods | | |
| The development of a radioimmunoassay for Tamm-Horsfall glycoprotein in serum | A. DAWNAY, C. MCLEAN & W. R. CATTELL | 679-687 |

INDEXES

| | |
|---------------|---------|
| Author Index | 791-792 |
| Subject Index | 793-798 |