

## Cellular Aspects

### Contents of Volume 210, No. 3, 1983

#### Protein Biosynthesis/Molecular Genetics

|  |  |         |
|--|--|---------|
| The effect of insulin infusion and food intake on muscle protein synthesis in postabsorptive rats  | P. J. GARLICK, M. FERN & V. R. PREEDY    | 669-676 |
| Synthesis and secretion of C-reactive protein by rabbit primary hepatocyte cultures                | S. S. MACINTYRE, D. SCHULTZ & I. KUSHNER | 707-715 |
| Comparison of cadmium-metallothionein synthesis in parenchymal and non-parenchymal rat liver cells | K. CAIN & D. N. SKILLETER                | 769-773 |

#### Protein Turnover

|   |                                |         |
|---|--------------------------------|---------|
| Inhibition of pyruvate carboxylase degradation and total protein breakdown by lysosomotropic agents in 3T3-L1 cells | C. S. CHANDLER & F. J. BALLARD | 845-853 |
|---|--------------------------------|---------|

#### Metabolism, Regulation and Control Processes

|  |   |         |
|--|---|---------|
| The role of substrate supply in the regulation of cholesterol biosynthesis in rat hepatocytes  | C. R. PULLINGER & G. F. GIBBONS                                   | 625-632 |
| Changes with starvation in the rat of the lipoprotein lipase activity and hydrolysis of triacylglycerols from triacylglycerol-rich lipoproteins in adipose tissue preparations | M. A. LASUNCIÓN & E. HERRERA                                      | 639-643 |
| Control of alanine metabolism in rat liver by transport processes or cellular metabolism   | P. FAFOURNOUX, C. RÉMÉSY & C. DEMIGNÉ                             | 645-652 |
| Changes in glycosaminoglycan biosynthesis during differentiation <i>in vitro</i> of human monocytes  | S. O. KOLSET, L. KJELLÉN, R. SELJELID & U. LINDAHL                | 661-667 |
| Water- and solute-accessible spaces of purified peroxisomes. Evidence that peroxisomes are permeable to NAD <sup>+</sup>   | P. VAN VELDHoven, L. J. DEBEER & G. P. MANNAERTS                  | 685-693 |
| The relationship between glutamate deamination and gluconeogenesis in kidney   | R. T. BOGUSKY, L. M. LOWENSTEIN & T. T. AOKI                      | 695-698 |
| Studies on the role of zinc in parturition in the rat  | G. E. BUNCE, G. R. WILSON, C. F. MILLS & A. KLOPPER               | 761-767 |
| Effects of glucose on phosphorylase and glycogen synthase in hepatocytes from diabetic rats  | M. BOLLEN, L. HUE & W. STALMANS                                   | 783-787 |
| Inhibition by cyanate of the processing of lysosomal enzymes   | A. HASILIK, R. POHLMANN & K. VON FIGURA                           | 795-802 |
| The contribution of phenylalanine to tyrosine metabolism <i>in vivo</i> . Studies in the post-absorptive and phenylalanine-loaded rat  | L. L. MOLDAWER, I. KAWAMURA, B. R. BISTRAN & G. L. BLACKBURN      | 811-817 |
| Stimulation of mitochondria functions by glucagon treatment. Evidence that effects are not artifacts of mitochondrial isolation  | C. B. JENSEN, F. D. SISTARE, H. C. HAMMAN & R. C. HAYNES, JR.     | 819-827 |
| Deoxycholic acid degradation by a <i>Pseudomonas</i> sp. Acidic intermediates with A-ring unsaturation   | R. A. LEPPIK  | 829-836 |
| Control of activation of liver RNA polymerase I occurring after re-feeding of protein-depleted mice  | L. HAIM, S. IAPALUCCI-ESPINOZA, R. CONDE & M. T. FRANZE-FERNÁNDEZ | 837-844 |
| The role of 5-aminolaevulinic synthase, haem oxygenase and ligand formation in the mechanism of maintenance of cytochrome P-450 concentration in hepatocyte culture            | L. J. HOCKIN & A. J. PAINE  | 855-857 |
| $\alpha$ -Adrenergic stimulation of respiration in isolated rat hepatocytes  | A. BINET & M. CLARET  | 867-873 |

|   |  |         |
|---|--|---------|
| Induction of peroxisomal enzymes in livers of neonatal rats exposed to lactating mothers treated with hypolipidaemic drugs. Role of drug metabolite transfer in milk  | W. E. FAHL, N. D. LALWANI,<br>M. K. REDDY & J. K. REDDY  | 875-883 |
| Mechanism of 3-phenylpyruvate-induced insulin release. Secretory, ionic and oxidative aspects   | A. SENER, M. WELSH, P. LEBRUN,<br>P. GARCIA-MORALES, M. SACEDA,<br>F. MALAISSE-LAGAE, A. HERCHUELZ,<br>I. VALVERDE, C. HELLERSTRÖM &<br>W. J. MALAISSE | 913-919 |
| Mechanism of 3-phenylpyruvate-induced insulin release. Metabolic aspects  | W. J. MALAISSE, A. SENER, M. WELSH,<br>F. MALAISSE-LAGAE, C. HELLERSTRÖM<br>& J. CHRISTOPHE  | 921-927 |
| Interactions between insulin and thyroid hormone in the control of lipogenesis  | M. C. SUGDEN, S. E. STEARE,<br>D. I. WATTS & T. N. PALMER  | 937-944 |
| Polyamine starvation causes accumulation of cadaverine and its derivatives in a polyamine-dependent strain of Chinese-hamster ovary cells   | E. HÖLTTÄ & P. POHJANPELTO   | 945-948 |
| Relationship between fatty-acyl composition of diacylgalactosylglycerol and turnover of chloroplast phosphatidate   | S. E. GARDINER & P. G. ROUGHAN   | 949-952 |
| Increased sensitivity of carnitine palmitoyltransferase I activity to malonyl-CoA inhibition after preincubation of intact rat liver mitochondria with macromolar concentrations of malonyl-CoA <i>in vitro</i> | V. A. ZAMMIT   | 953-956 |
| Hormonal stimulation of mitochondrial glutaminase. Effects of vasopressin, angiotensin II, adrenaline and glucagon  | S. CORVERA & J. A. GARCÍA-SÁINZ  | 957-960 |
| <b>Cell Surfaces and Receptors</b>  |  |         |
| Acceptor sites for the oestrogen receptor in hen oviduct chromatin  | T. S. RUH & T. C. SPELSBERG  | 905-912 |
| <b>Developmental Biochemistry</b>   |  |         |
| The effect of a vitamin B-6 antagonist, 4-deoxypyridoxine, on the cross-linking of collagen in the developing chick embryo  | T. A. BIRD & C. I. LEVENE  | 633-638 |
| Identification and origin of oestradiol-binding protein in immature rat skin. Demonstration of oestrophilic $\alpha$ -foetoprotein synthesis and secretion by developing rat skin explants in culture           | M. ALI, K. MUJOO & M. K. SAHIB   | 653-660 |
| Glucose metabolism and its regulation in pieces of perirenal adipose tissue from foetal lambs   | J. P. ROBERTSON, A. FAULKNER &<br>R. G. VERNON   | 677-683 |
| Characterization and partial purification of a haemopoietic cell growth factor in WEHI-3 cell conditioned medium  | G. W. BAZILL, M. HAYNES,<br>J. GARLAND & T. M. DEXTER  | 747-759 |
| <b>Membranes, Transport, Bioenergetics and Photosynthesis</b>   |  |         |
| The association of proton movement with galactose transport into subcellular membrane vesicles of <i>Escherichia coli</i>   | P. HORNE & P. J. F. HENDERSON  | 699-705 |
| Photochemical action spectra indicate that cytochrome <i>a/a</i> <sub>3</sub> is the predominant haemoprotein terminal oxidase in <i>Acanthamoeba castellanii</i>   | R. I. SCOTT & D. LLOYD   | 721-725 |
| A kinetic study of the interaction between mitochondrial F <sub>1</sub> adenosine triphosphatase and adenylyl imidodiphosphate and guanylyl imidodiphosphate  | F. J. F. BELDA, F. G. CARMONA,<br>F. G. CÁNOVAS, J. C. GÓMEZ-<br>FERNÁNDEZ & J. A. LOZANO  | 727-735 |
| Rat mast cells permeabilized with Sendai virus secrete histamine in response to Ca <sup>2+</sup> buffered in the micromolar range   | B. D. GOMPERTS, J. M. BALDWIN &<br>K. J. MICKLEM   | 737-745 |

|   |   |         |
|---|---|---------|
| Transport of <i>myo</i> -inositol into endoneurial preparations of sciatic nerve from normal and streptozotocin-diabetic rats           | K. R. W. GILLON & J. N. HAWTHORNE                                       | 775-781 |
| Specificity and properties of the nucleotide carrier in chromaffin granules from bovine adrenal medulla                                 | A. WEBER, E. W. WESTHEAD & H. WINKLER                                   | 789-794 |
| 5-Hydroxytryptamine transport in cells and secretory granules from a transplantable rat insulinoma                                      | J. C. HUTTON, M. PESHAVARIA & N. E. TOOKE                               | 803-810 |
| An immunological study of the uncoupling protein of brown adipose tissue mitochondria   | D. RICQUIER, J.-P. BARLET, J.-M. GAREL, M. COMBES-GEORGE & M. P. DUBOIS | 859-866 |
| Early plasma-membrane-potential changes during stimulation of lymphocytes by concanavalin A   | S. M. FELBER & M. D. BRAND  | 885-891 |
| Concanavalin A causes an increase in sodium permeability and intracellular sodium content of pig lymphocytes                            | S. M. FELBER & M. D. BRAND  | 893-897 |
| Glutathione and ascorbic acid in spinach ( <i>Spinacia oleracea</i> ) chloroplasts. The effect of hydrogen peroxide and of Paraquat     | M. Y. LAW, S. A. CHARLES & B. HALLIWELL                                 | 899-903 |
| Cellular and lysosomal uptake of methylamine in isolated rat hepatocytes  | A. E. SOLHEIM & P. O. SEGLEN  | 929-936 |
| <b>General Methods</b>  |   |         |
| A simple assay method for determination of the specific radioactivity of the $\gamma$ -phosphate group of $^{32}\text{P}$ -labelled ATP | P. T. HAWKINS, R. H. MICHELL & C. J. KIRK                               | 717-720 |
| Measurement of intracellular potassium ion concentrations by n.m.r.   | P. J. BROPHY, M. K. HAYER & F. G. RIDDELL                               | 961-963 |