

# CONTENTS

## Volume 225, No. 3, 1985

### FULL AND RAPID PAPERS

#### Proteins

|  |   |         |
|--|---|---------|
| Relations between high-affinity binding sites of markers for binding regions on human serum albumin  | U. KRAGH-HANSEN                           | 629-638 |
| Evidence for glucose-mediated covalent cross-linking of collagen after glycosylation <i>in vitro</i> | M. J. C. KENT, N. D. LIGHT & A. J. BAILEY | 745-752 |

#### Enzymes

|   |   |         |
|---|---|---------|
| Investigation of the effect of metal ions on the reactivity of thiol groups in human 5-aminolaevulinatase dehydratase   | P. N. B. GIBBS, M. G. GORE & P. M. JORDAN                 | 573-580 |
| Increased dihydrofolate reductase activity in methotrexate-resistant human promyelocytic-leukaemia (HL-60) cells. Lack of correlation between increased activity and overproduction | S. DEDHAR, D. HARTLEY & J. H. GOLDIE                      | 609-617 |
| Translocation of mycobactin synthetase in <i>Bacillus subtilis</i>  | N. K. MUKHOPADHYAY, S. K. GHOSH, S. MAJUMDER & S. K. BOSE | 639-643 |
| Contribution of different protein phosphatases to the dephosphorylation of 6-phosphofructo-1-kinase and 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase in rat liver           | G. MIESKES & H.-D. SÖLING                                 | 665-670 |
| Phospholipid biosynthesis in human platelets. The acylation of lyso-platelet-activating factor  | M. L. MCKEAN & M. J. SILVER                               | 723-729 |
| Resolution and reconstitution of bovine kidney branched-chain 2-oxo acid dehydrogenase complex  | K. G. COOK, A. P. BRADFORD & S. J. YEAMAN                 | 731-735 |
| Determination of acetylcholinesterase activity by a new chemiluminescence assay with the natural substrate  | S. BIRMAN   | 825-828 |

#### Nucleic acids and molecular genetics

|   |   |         |
|---|---|---------|
| The use of monoclonal antibodies to study the proteins specified by the transforming region of human adenoviruses | M. E. BLAIR ZAJDEL, M. D. BARKER, S. C. DIXON & G. E. BLAIR | 649-655 |
| A mouse gene family associated with a major submaxillary-gland glycoprotein                                       | A. G. CRAIG, J. J. MULLINS, P. MCTURK & W. J. BRAMMAR       | 657-663 |
| Molecular cloning of a bovine cathepsin   | N. J. GAY & J. E. WALKER                                    | 707-712 |
| Design of a new DNA-polyintercalating drug, a bisacridinyl peptidic analogue of Triostin A                        | N. HELBECQUE, J.-L. BERNIER & J.-P. HÉNICHART               | 829-832 |

#### Synthesis and degradation of macromolecules

|   |  |         |
|---|--|---------|
| Comparison of the behaviour <i>in vivo</i> of two molecular forms of antithrombin III   | T. H. CARLSON, C. A. ATENCIO & T. L. SIMON   | 557-564 |
| Extracellular-matrix synthesis by skeletal muscle in culture. Major secreted collagenous proteins of clonal myoblasts   | R. L. BEACH, J. S. RAO & B. W. FESTOFF   | 619-627 |
| The effects of 17 $\alpha$ -ethynyl-estradiol and of acute inflammation on the plasma concentration of rat $\alpha_1$ -acid glycoprotein and on the induction of its hepatic mRNA | M. DIARRA-MEHRPOUR, J. BOURGUIGNON, I. LEROUX-NICOLLET, D. MARKO-VERCAIGNE, D. BIOU, M. HIRON & J.-P. LEBRETON | 681-687 |
| Inhibition of hyaluronate synthesis   | P. PREHM   | 699-705 |

#### Intermediary metabolism and metabolic control

|  |   |         |
|--|---|---------|
| Increased loss and decreased synthesis of hepatic glutathione after acute ethanol administration. Turnover studies | H. SPEISKY, A. MACDONALD, G. GILES, H. ORREGO & Y. ISRAEL                     | 565-572 |
| Calcium and calmodulin in the regulation of human thyroid adenylate cyclase activity                               | T. LAKEY, S. MAC NEIL, H. HUMPHRIES, S. W. WALKER, D. S. MUNRO & S. TOMLINSON | 581-589 |

## Contents

|   |   |         |
|---|---|---------|
| Stimulation of creatine kinase BB activity by parathyroid hormone and by prostaglandin E <sub>2</sub> in cultured bone cells  | D. SÖMJEN, A. M. KAYE & I. BINDERMAN                          | 591-596 |
| Accumulation of ornithine decarboxylase-antizyme complex in HMO <sub>A</sub> cells  | Y. MURAKAMI, K. FUJITA, T. KAMEJI & S. HAYASHI                | 689-697 |
| Branched-chain amino acid metabolism and alanine formation in rat muscles <i>in vitro</i> . Mitochondria-cytosolic interrelationships   | K. SNELL & D. A. DUFF   | 737-743 |
| Quantitative analysis of intermediary metabolism in hepatocytes incubated in the presence and absence of glucagon with a substrate mixture containing glucose, ribose, fructose, alanine and acetate                                | M. RABKIN & J. J. BLUM  | 761-786 |
| <i>Appendix</i> : Statistical considerations pertaining to analysis of metabolic models   | D. S. BURDICK   | 783-785 |
| <b>Bioenergetics and transport</b>  |   |         |
| Phospholipid association with the bovine cardiac mitochondrial adenosine triphosphatase   | R. E. BROWN, R. I. MONTGOMERY, P. I. SPACH & C. C. CUNNINGHAM | 597-608 |
| Mechanism of depolarization of rat cortical synaptosomes at submicromolar external Ca <sup>2+</sup> activity. The use of Ca <sup>2+</sup> buffers to control the synaptosomal membrane potential                                    | G. SCHMALZING   | 671-680 |
| Tetraphenylphosphonium ion is a true indicator of negative plasma-membrane potential in the yeast <i>Rhodotorula glutinis</i> . Experiments under osmotic stress and at low external pH values                                      | M. HÖFER & A. KÜNEMUND  | 815-819 |
| Purification of F <sub>1</sub> -ATPase from cuckoo-pint ( <i>Arum maculatum</i> ) mitochondria. A comparison of subunit composition with that of rat liver F <sub>1</sub> -ATPase   | P. P. J. DUNN, A. R. SLABAS & A. L. MOORE                     | 821-824 |
| <b>Membranes and receptors</b>  |   |         |
| Gangliosides and sialosylglycoproteins in coated vesicles from bovine brain   | D. GRAVOTTA & H. J. F. MACCIONI                               | 713-721 |
| Affinity-labelling of the thyrotropin receptor. Characterization of the photoactive ligand  | P. R. BUCKLAND, R. D. HOWELLS, C. R. RICKARDS & B. REES SMITH | 753-760 |
| Redistribution of membrane-bound and cytosolic actin in rabbit polymorphonuclear leucocytes during phagocytosis   | D. I. H. STEWART & N. CRAWFORD                                | 807-814 |
| <b>Cell structure and function</b>  |   |         |
| Effect of glycy-L-phenylalanine 2-naphthylamide on invertase endocytosed by rat liver   | M. JADOT & R. WATTIAUX  | 645-648 |
| <b>Methods</b>  |   |         |
| Reverse-phase h.p.l.c. separation, quantification and preparation of bilirubin and its conjugates from native bile. Quantitative analysis of the intact tetrapyrroles based on h.p.l.c. of their ethyl anthranilate azo derivatives | W. SPIVAK & M. C. CAREY                                       | 787-805 |