

**Review**

Regulated exocytosis  
by R.D. Burgoyne and A. Morgan

305–316

**Research Communications**

Band 3 HT, a human red-cell variant associated with acanthocytosis and increased anion transport, carries the mutation Pro-868 → Leu in the membrane domain of band 3

L.J. Bruce, M.M.B. Kay, C. Lawrence and M.J.A. Tanner **317–320**

Peptide glyoxals: a novel class of inhibitor for serine and cysteine proteinases

B. Walker, N. McCarthy, A. Healy, T. Ye and M.A. McKervey **321–323**

Predominant expression of type-VI adenylate cyclase in C6-2B rat glioma cells may account for inhibition of cyclic AMP accumulation by calcium

M.A. Debernardi, R. Munshi, M. Yoshimura, D.M.F. Cooper and G. Brooker **325–328**

**Research Papers****Proteins**

Studies on secondary structure of caldesmon and its C-terminal fragments

E.A. Czurylo, S.Y. Venyaminov and R. Dąbrowska **363–368**

Binding of platelet-derived growth factor-BB and transforming growth factor- $\beta$ 1 to  $\alpha_2$ -macroglobulin *in vitro* and *in vivo*: comparison of receptor-recognized and non-recognized  $\alpha_2$ -macroglobulin conformations

K.P. Crookston, D.J. Webb, J. Lamarre and S.L. Gonias **443–450**

Calorimetric studies of the N-terminal half-molecule of transferrin and mutant forms modified near the Fe<sup>3+</sup>-binding site

L.-N. Lin, A.B. Mason, R.C. Woodworth and J. F. Brandts **517–522**

<sup>1</sup>H-n.m.r. study of the solution properties and secondary structure of neurotoxin III from the sea anemone *Anemonia sulcata*

R.S. Norton, K. Cross, V. Braach-Maksvytis and E. Wachter **545–551**

Purification, characterization and binding interactions of the Chinese-cobra (*Naja naja atra*) serum antitoxic protein CSAP

J. Shao, H. Shen and B. Havsteen **559–566**

**Enzymes**

Unusual reactivity of Tyr-7 of GSH transferase P1-1

D.J. Meyer, C. Xia, B. Coles, H. Chen, P. Reinemer, R. Huber and B. Ketterer **351–356**

Chemical modification of GSH transferase P1-1 confirms the presence of Arg-13, Lys-44 and one carboxylate group in the GSH-binding domain of the active site

C. Xia, D. J. Meyer, H. Chen, P. Reinemer, R. Huber and B. Ketterer **357–362**

Influence of cellobiose oxidase on peroxidases from *Phanerochaete chrysosporium*

P. Ander, G. Sena-Martins and J.C. Duarte **431–435**

Activation of procathepsin B in human hepatoma cells: the conversion into the mature enzyme relies on the action of cathepsin B itself

L. Mach, H. Schwihla, K. Stüwe, A.D. Rowan, J.S. Mort and J. Glössl **437–442**

Digestion of damaged DNA by the T7 DNA polymerase-exonuclease L(+) -Mandelate dehydrogenase from *Rhodotorula graminis*: purification, partial characterization and identification as a flavocytochrome *b*

D.R. Koehler and P.C. Hanawalt **451–453**

M. Yasin and C.A. Fewson **455–460**

*Trichomonas vaginalis*: characterization of ornithine decarboxylase

N. Yarlett, B. Goldberg, M.A. Moharrami and C.J. Bacchi **487–493**

Inactivation of the endogenous argininosuccinate lyase activity of duck  $\delta$ -crystallin by modification of an essential histidine residue with diethyl pyrocarbonate

H.-J. Lee, S.-H. Chiou and G.-G. Chang **537–544**

Binding of a substrate analogue can induce co-operative structure in the plasmin serine-proteinase domain

A.J. Teuten, A. Cooper, R.A.G. Smith and C.M. Dobson **567–572**

Polarimetry and <sup>13</sup>C n.m.r. show that the hydrolyses of  $\beta$ -D-glucopyranosyl fluoride by  $\beta$ (1 → 3)-glucanases from *Phanerochaete chrysosporium* and *Sporotrichum dimorphosporum* have opposite stereochemistries

J.L. Copa-Patiño, Y. Zhang, B. Padmaperuma, I. Marsden, P. Broda and M.L. Sinnott **591–594**

Mg <sup>2+</sup> affects the binding of ADP but not ATP to 3-phosphoglycerate kinase. Correlation between equilibrium dialysis binding and enzyme kinetic data	M. Molnár and M. Vas	595–599
<b>Carbohydrates and lipids</b>		
Biochemical characterization of a rat oncofetal colonic antigen defined by a monoclonal antibody raised against gastric surface epithelium	C. Decaens, J. Nardelli, J. Bara and P. Burtin	531–536
<b>Gene structure and expression</b>		
Degenerate 87-base-pair tandem repeats create hydrophilic/hydrophobic alternating domains in human mucin peptides mapped to 11p15	J. Dufosse, N. Porchet, J.-P. Audie, V. Guyonnet Duperat, A. Laine, I. Van-Seuningen, S. Marrakchi, P. Degand and J.-P. Aubert	329–337
A type I collagen reporter gene construct for protein engineering studies. Functional equivalence of transfected reporter <i>COL1A1</i> and endogenous gene products during biosynthesis and <i>in vitro</i> extracellular matrix accumulation	S.R. Lamandé and J.F. Bateman	387–394
Cytosolic and nuclear spermidine acetyltransferases in growing NIH 3T3 fibroblasts stimulated with serum or polyamines: relationship to polyamine-biosynthetic decarboxylases and histone acetyltransferase	M.A. Desiderio, S. Mattei, G. Biondi and M.P. Colombo	475–479
Characterization of a full-length cDNA encoding human liver S-adenosylmethionine synthetase: tissue-specific gene expression and mRNA levels in hepatopathies	L. Alvarez, F. Corrales, A. Martín-Duce and J.M. Mato	481–486
Transgenic mice over-expressing the human spermidine synthase gene	L. Kauppinen, S. Myöhänen, M. Halmekytö, L. Alhonen and J. Jänne	513–516
<b>Regulation of metabolism</b>		
Rapid activation of hepatic glutaminase in rats fed on a single high-protein meal	H.S. Ewart and J.T. Brosnan	339–344
Use of superfused rat skeletal muscle for metabolic studies: assessment of pH by <sup>31</sup> P n.m.r.	D. Meynial-Denis, M. Mignon, L. Foucat, Y. Bonnet, G. Biélicki, J.-P. Renou, P. Lacourt, A. Lacourt and M. Arnal	399–405
Calcium measurements with a new high-affinity n.m.r. indicator in the isolated perfused heart	H.L. Kirschenlohr, A.A. Grace, S.D. Clarke, Y. Shachar-Hill, J.C. Metcalfe, P.G. Morris and G.A. Smith	407–411
<sup>15</sup> N n.m.r. measurement of the <i>in vivo</i> rate of glutamine synthesis and utilization at steady state in the brain of the hyperammonaemic rat	K. Kanamori and B.D. Ross	461–468
Hepatocyte heterogeneity in response to extracellular adenosine	Y. Morimoto, M. Wettstein and D. Häussinger	573–581
<b>Membranes and bioenergetics</b>		
The erythrocyte calcium pump is inhibited by non-enzymic glycation: studies <i>in situ</i> and with the purified enzyme	F.L. González Flecha, P.R. Castello, A.J. Caride, J.J. Gagliardino and J.P.F.C. Rossi	369–375
The Ca <sup>2+</sup> -transporting ATPases of rabbit and trout exhibit different pH- and temperature-dependences	E.N. Chini, F.G.S. de Toledo, M.C. Albuquerque and L. de Meis	469–473
Cloning, <i>in vitro</i> mitochondrial import and membrane assembly of the 17.8 kDa subunit of complex I from <i>Neurospora crassa</i>	J.E. Azevedo, J. Abrolat-Scharff, C. Eckerskorn and S. Werner	501–506
<b>Receptors and signal transduction</b>		
Influence of Mg <sup>2+</sup> on detection of somatogenic and lactogenic components of growth-hormone-binding protein in mammalian sera	T. Amit, Z. Hochberg and R.J. Barkey	345–349
Endothelins stimulate tyrosine phosphorylation and activity of p42/mitogen-activated protein kinase in astrocytes	S. Cazaubon, P.J. Parker, A.D. Strosberg and P.-O. Couraud	381–386
Two independently regulated Ca <sup>2+</sup> entry mechanisms coexist in Jurkat T cells during T cell receptor antigen activation	S.C. Chow, G.E.N. Kass and S. Orrenius	395–398
Multiple mechanisms by which protein kinase A potentiates inositol 1,4,5-trisphosphate-induced Ca <sup>2+</sup> mobilization in permeabilized hepatocytes	G. Hajnóczky, E. Gao, T. Nomura, J.B. Hoek and A.P. Thomas	413–422

---

Mobilization of Ca <sup>2+</sup> by thapsigargin and 2,5-di-(t-butyl)-1,4-benzohydroquinone in permeabilized insulin-secreting RINm5F cells: evidence for separate uptake and release compartments in inositol 1,4,5-trisphosphate-sensitive Ca <sup>2+</sup> pool	M.S. Islam and P.-O. Berggren	<b>423–429</b>
Enhanced degradation of the phosphoinositidase C-linked guanine-nucleotide-binding protein G <sub>q</sub> $\alpha$ /G <sub>11</sub> $\alpha$ following activation of the human M1 muscarinic acetylcholine receptor expressed in CHO cells	F.M. Mitchell, N.J. Buckley and G. Milligan	<b>495–499</b>
Non-catalytic activation of phospholipase C- $\gamma$ 1 <i>in vitro</i> by epidermal growth factor receptor	S.M.T. Hernández-Sotomayor and G. Carpenter	<b>507–511</b>
4,4'-Di-isothiocyanatostilbene-2,2'-disulphonic acid ('DIDS') activates protein kinase C and Na <sup>+</sup> /H <sup>+</sup> exchange in human platelets via $\alpha_{2A}$ -adrenergic receptors	R. Nieuwland, G. Van Willigen and J.-W.N. Akkerman	<b>523–530</b>
Turnover of inositol pentakisphosphates, inositol hexakisphosphate and diphosphoinositol polyphosphates in primary cultured hepatocytes	M.C. Glennon and S.B. Shears	<b>583–590</b>
<b>Cell biology and development</b>		
Parathyroid hormone transport effects and hormonal processing in primary cultured rat proximal tubular cells	R.M. O'Donovan, C.C. Widnell, T.C. Chen and J.B. Puschett	<b>377–380</b>
Effect of betaine on HSP70 expression and cell survival during adaptation to osmotic stress	P.G. Petronini, E.M. De Angelis, A.F. Borghetti and K.P. Wheeler	<b>553–558</b>