

## Review

Transcriptional control of genes that regulate glycolysis and gluconeogenesis in adult liver

by F.P. Lemaigre and G.G. Rousseau

1-14

Biochemical Journal

Volume 303, part 1

1 October 1994

## Research Communications

Wortmannin inhibits the effects of insulin and serum on the activities of glycogen synthase kinase-3 and mitogen-activated protein kinase

The inhibition of glycogen synthase kinase-3 by insulin or insulin-like growth factor 1 in the rat skeletal muscle cell line L6 is blocked by wortmannin, but not by rapamycin: evidence that wortmannin blocks activation of the mitogen-activated protein kinase pathway in L6 cells between Ras and Raf

The mechanism by which epidermal growth factor inhibits glycogen synthase kinase 3 in A431 cells

Prolactin increases hepatic Na<sup>+</sup>/taurocholate co-transport activity and messenger RNA *post partum*

Divergent sequences in the 5' region of cDNA suggest alternative splicing as a mechanism for the generation of carnitine acetyltransferases with different subcellular localizations

G.I. Welsh, E.J. Foulstone, S.W. Young, J.M. Tavaré and C.G. Proud

15-20

D.A.E. Cross, D.R. Alessi, J.R. Vandenheede, H.E. McDowell, H.S. Hundal and P. Cohen

21-26

Y. Saito, J.R. Vandenheede and P. Cohen

27-32

T.C. Ganguly, Y. Liu, J.F. Hyde, B. Hagenbuch, P.J. Meier and M. Vore

33-36

O. Corti, S. DiDonato and G. Finocchiaro

37-41

## Research Papers

### Proteins

Oxidized mucus proteinase inhibitor: a fairly potent neutrophil elastase inhibitor

Increased thermal stability of proteins in the presence of amino acids

Characterization of monoclonal antibodies against *Naja naja oxiana* neurotoxin I

The non-phospholipase A<sub>2</sub> subunit of  $\beta$ -bungarotoxin plays an important role in the phospholipase A<sub>2</sub>-independent neurotoxic effect: characterization of three isotoxins with a common phospholipase A<sub>2</sub> subunit

Human seminal plasma contains somatostatin-64

C. Boudier and J.G. Bieth

61-68

S. Taneja and F. Ahmad

147-153

B.G. Stiles, F.W. Sexton, S.B. Guest, M.A. Olson and D.C. Hack

163-170

C.-C. Chu, S.-T. Chu, S.-W. Chen and Y.-H. Chen

171-176

L. Ødum and A.H. Johnsen

263-268

### Enzymes

Chemical modification of *Penicillium* 1,2- $\alpha$ -D-mannosidase by water-soluble carbodi-imide: identification of a catalytically important aspartic acid residue

Major proteins of bovine seminal plasma inhibit phospholipase A<sub>2</sub>

Thermal stability of methanol dehydrogenase is altered by the replacement of enzyme-bound Ca<sup>2+</sup> with Sr<sup>2+</sup>

Identification of one- and two-chain forms of trypsinogen 1 produced by a human gastric adenocarcinoma cell line

Investigation of the substrate specificity of a cloned expressed human bilirubin UDP-glucuronosyltransferase: UDP-sugar specificity and involvement in steroid and xenobiotic glucuronidation

Inactivation precedes changes in allosteric properties and conformation of D-glyceraldehyde-3-phosphate dehydrogenase and fructose-1,6-bisphosphatase during denaturation by guanidinium chloride

Microheterogeneity of cytosolic and membrane-bound hexokinase II in Morris hepatoma 3924A

Thalporphine selectively inhibits expression of the inducible, but not the constitutive, nitric oxide synthase

T. Yoshida, K. Maeda, M. Kobayashi and E. Ichishima

97-103

P. Manjunath, S. Soubeyrand, L. Chandonnet and K.D. Roberts

121-128

T.K. Harris and V.L. Davidson

141-145

N. Koshikawa, H. Yasumitsu, Y. Nagashima, M. Umeda and K. Miyazaki

187-190

S.B. Senafi, D.J. Clarke and B. Burchell

233-240

R.-F. Jiang and C.-L. Tsou

241-245

A. Rempel, P. Bannasch and D. Mayer

269-274

S.M. Yu

289-294

Incorporation of bisphosphonates into adenine nucleotides by amoebae of the cellular slime mould <i>Dictyostelium discoideum</i>	M.J. Rogers, X. Ji, R.G.G. Russell, G.M. Blackburn, M.P. Williamson, A.V. Bayless, F.H. Ebetino and D.J. Watts	<b>303–311</b>
Phosphinic peptide analogues as potent inhibitors of <i>Corynebacterium rathayii</i> bacterial collagenase	A. Yiotakis, A. Lecoq, A. Nicolaou, J. Labadie and V. Dive	<b>323–327</b>
<b>Carbohydrates and lipids</b>		
Sulphated and undersulphated heparan sulphate proteoglycans in a Chinese hamster ovary cell mutant defective in <i>N</i> -sulpho-transferase	K.J. Bame, L. Zhang, G. David and J.D. Esko	<b>81–87</b>
Effect of chronic incubation of CaCo-2 cells with eicosapentaenoic acid (20:5, <i>n</i> –3) and oleic acid (18:1, <i>n</i> –9) on triacylglycerol production	T. Ranheim, A. Gedde-Dahl, A.C. Rustan and C.A. Drevon	<b>155–161</b>
<b>Gene structure and expression</b>		
Expression of a biologically active plant cytochrome <i>b<sub>5</sub></i> in <i>Escherichia coli</i>	M.A. Smith, J.A. Napier, S. Stymne, A.S. Tatham, P.R. Shewry and A.K. Stobart	<b>73–79</b>
Heterogeneity of vacuolar H <sup>+</sup> -ATPase: differential expression of two human subunit B isoforms	B. van Hille, H. Richener, P. Schmid, I. Puetner, J.R. Green and G. Bilbe	<b>191–198</b>
Regulation of the expression of alkaline phosphatase in a human breast-cancer cell line	T.-C. Chang, J.-K. Wang, M.-W. Hung, C.-H. Chiao, L.-C. Tsai and G.-G. Chang	<b>199–205</b>
Cloning and characterization of the human osteopontin gene and its promoter	N. Hijjiya, M. Setoguchi, K. Matsuura, Y. Higuchi, S. Akizuki and S. Yamamoto	<b>255–262</b>
Plasma-membrane calcium-pump isoforms in human and rat liver	A. Howard, N.F. Barley, S. Legon and J.R.F. Walters	<b>275–279</b>
Human link protein gene: structure and transcription pattern in chondrocytes	J. Dudhia, M.T. Bayliss and T.E. Hardingham	<b>329–333</b>
<b>Regulation of metabolism</b>		
Pancreastatin activates protein kinase C by stimulating the formation of 1,2-diacylglycerol in rat hepatocytes	V. Sánchez-Margalet, M. Lucas and R. Goberna	<b>51–54</b>
Effects of the <i>S</i> -adenosylmethionine decarboxylase inhibitor, 5'-[[ <i>Z</i> ]-4-amino-2-butenyl]methylamino}-5'-deoxyadenosine, on cell growth and polyamine metabolism and transport in Chinese hamster ovary cell cultures	T.L. Byers, R.S. Wechter, R.-H. Hu and A.E. Pegg	<b>89–96</b>
Zinc, vanadate and selenate inhibit the tri-iodothyronine-induced expression of fatty acid synthase and malic enzyme in chick-embryo hepatocytes in culture	Y. Zhu, A.G. Goodridge and S.R. Stapleton	<b>213–216</b>
Interactions of peroxynitrite with human plasma and its constituents: oxidative damage and antioxidant depletion	A. Van Der Vliet, D. Smith, C.A. O'Neill, H. Kaur, V. Darley-Usmar, C.E. Cross and B. Halliwell	<b>295–301</b>
<b>Membranes and bioenergetics</b>		
Inhibition of electron flow through complex I of the mitochondrial respiratory chain of Ehrlich ascites carcinoma cells by methylglyoxal	S. Ray, S. Dutta, J. Halder and M. Ray	<b>69–72</b>
Partial purification and reconstitution of the sarcolemmal L-lactate carrier from rat skeletal muscle	P.J. Allen and G.A. Brooks	<b>207–212</b>
<b>Receptors and signal transduction</b>		
Characterization of phorbol ester-stimulated serine phosphorylation of the human insulin receptor	E.P. Feener, T. Shiba, K.-Q. Hu, P.A. Wilden, M.F. White and G.L. King	<b>43–50</b>
Lysophosphatidic acid activation of phosphatidylcholine-hydrolysing phospholipase D and actin polymerization by a pertussis toxin-sensitive mechanism	K.-S. Ha, E.-J. Yeo and J.H. Exton	<b>55–59</b>
Expression, purification and characterization of recombinant mitogen-activated protein kinase kinases	P. Dent, Y.-H. Chow, J. Wu, D.K. Morrison, R. Jove and T.W. Sturgill	<b>105–112</b>
The hemispheric functional expression of the thyrotropin-releasing-hormone receptor is not determined by the receptors' physical distribution	N. Matus-Leibovitch, D.R. Nussenzweig, M.C. Gershengorn and Y. Oron	<b>129–134</b>
Purification of the G-protein G <sub>13</sub> from rat brain membranes	R. Harhammer, B. Nürnberg, K. Spicher and G. Schultz	<b>135–140</b>

---

Agonist-induced internalization of the substance P (NK <sub>1</sub> ) receptor expressed in epithelial cells	A.M. Garland, E.F. Grady, D.G. Payan, S.R. Vigna and N.W. Bunnett	<b>177–186</b>
Role of prostaglandin-mediated cyclic AMP formation in protein kinase C-dependent secretion of atrial natriuretic peptide in rat cardiomyocytes	D.J. Church, V. Van Der Bent, M.B. Vallotton and U. Lang	<b>217–225</b>
Characterization of an atypical lipoprotein-binding protein in human aortic media membranes by ligand blotting	Y.S. Kuzmenko, V.N. Bochkov, M.P. Philippova, V.A. Tkachuk and T.J. Resink	<b>281–287</b>
Switching from $\alpha_1$ - to $\beta$ -subtypes in adrenergic response during primary culture of adult-rat hepatocytes as affected by the cell-to-cell interaction through plasma membranes	Y. Kajiyama and M. Ui	<b>313–321</b>
<b>Cell biology and development</b>		
Effects of aniso-osmolarity and hydroperoxides on intracellular pH in isolated rat hepatocytes as assessed by (2',7')-bis(carboxyethyl)-5(6)-carboxyfluorescein and fluorescein isothiocyanate–dextran fluorescence	R. Schreiber, B. Stoll, F. Lang and D. Häussinger	<b>113–120</b>
Thrombin induces endothelial cell growth via both a proteolytic and a non-proteolytic pathway	J.-M. Herbert, E. Dupuy, M.-C. Laplace, J.-M. Zini, R. Bar Shavit and G. Tobelem	<b>227–231</b>
Role of phospholipase A <sub>2</sub> in expression of the scavenger pathway in cultured aortic smooth muscle cells stimulated with phorbol 12-myristate 13-acetate	N. Morisaki, K. Yokote, M. Otabe, Y. Saito, S. Yoshida and S. Ueda	<b>247–253</b>
<b>BJ Letters</b>		
PH-20 and sperm hyaluronidase: a conceptual conundrum in mammalian fertilization	P. Gacesa, N.D. Civill and R.A.P. Harrison	<b>335–336</b>