| Regulation of ribosomal gene transcription by S.T. Jacob 617–626  | Volume 306, part 3<br>15 March 1995  |         |
|---|--|---------|
| Research Communication  |  |         |
| Formation <i>in vitro</i> of the 3,4,6-trihydroxyphenylalanine quinone cofactor   | S.P. Hanlon, K. Carpenter, A. Hassan and R.A. Cooper   | 627–630 |
| Proteins  |  |         |
| Detection of acyl-CoA-binding protein in human red blood cells and investigation of its role in membrane phospholipid renewal   | H. Fyrst, J. Knudsen, M.A. Schott, B.H. Lubin and F.A. Kuypers   | 793–799 |
| In vitro fibrillogenesis of collagen II from pig vitreous humour  | C. Yang, H. Notbohm, Y. Açil, R. Heifeng, S. Bierbaum and P.K. Müller                                  | 871–875 |
| Enzymes   |  |         |
| Recombinant 10-formyltetrahydrofolate dehydrogenase catalyses both dehydrogenase and hydrolase reactions utilizing the synthetic substrate 10-formyl-5,8-dideazafolate  | S.A. Krupenko, C. Wagner and R.J. Cook   | 651–655 |
| Inhibition of specific binding of okadaic acid to protein phosphatase 2A by microcystin-LR, calyculin-A and tautomycin: method of analysis of interactions of tight-binding ligands with target protein                       | A. Takai, K. Sasaki, H. Nagai, G. Mieskes, M. Isobe,<br>K. Isono and T. Yasumoto                       | 657–665 |
| Solubilization, molecular forms, purification and substrate specificity of two acetylcholinesterases in the medicinal leech ( <i>Hirudo medicinalis</i> )   | V. Talesa, M. Grauso, E. Giovannini, G. Rosi and<br>JP. Toutant  | 687–692 |
| Stability of Japanese-lacquer-tree ( <i>Rhus vernicifera</i> ) laccase to thermal and chemical denaturation: comparison with ascorbate oxidase  | E. Agostinelli, L. Cervoni, A. Giartosio and L. Morpurgo   | 697–702 |
| Interaction of component enzymes with the peripheral subunit-<br>binding domain of the pyruvate dehydrogenae multienzyme<br>complex of <i>Bacillus stearothermophilus</i> : stoichiometry and<br>specificity in self-assembly | I.A.D. Lessard and R.N. Perham   | 727–733 |
| Irreversible inactivation of lactoperoxidase by mercaptomethylimidazole through generation of a thiyl radical: its use as a probe to study the active site  | U. Bandyopadhyay, D.Kr. Bhattacharyya, R. Chatterjee and R.K. Banerjee                                 | 751–757 |
| A stopped-flow apparatus for infrared spectroscopy of aqueous solutions   | A.J. White, K. Drabble and C.W. Wharton  | 843-849 |
| The properties of a cloned human high-molecular-mass cytosolic phospholipase $A_2$ investigated using a continuous fluorescence displacement assay: evidence for enzyme clustering on phospholipid vesicles                   | A. Creaney, D.J. Masters, M.B. Needham, R.D. Gordon, R. Mott and D.C. Wilton                           | 857–864 |
| Carbohydrates and lipids  |  |         |
| Decrease in cardiac phosphatidylglycerol in streptozotocin-induced diabetic rats does not affect cardiolipin biosynthesis: evidence for distinct pools of phosphatidylglycerol in the heart                                   | G.M. Hatch, S.G. Cao and A. Angel  | 759–764 |
| Occurrence of a novel fucose-containing pentaglycosylceramide with blood-group-B active determinant in <i>Xenopus</i> blastula cells: its possible involvement in cell-cell adhesion  | K. Nomura, N. Nakajo, K.IP.J. Hidari, H. Nomura,<br>M. Murata, M. Suzuki, K. Yamana and Y. Hirabayashi | 821–827 |
| Gene structure and expression   |  |         |
| Cholinergic regulation of amylase gene expression in the rat parotid gland. Inhibition by two distinct post-transcriptional mechanisms  | Y. Liu, P.Y. Woon, S.C. Lim, K. Jeyaseelan and P. Thiyagarajah   | 637–642 |

Review

**Biochemical Journal** 

| Regulation of metabolism  |  |                      |
|---|--|----------------------|
| Microcystin-LR induced an inhibition of protein synthesis in isolated   | S. Claeyssens, A. Francois, A. Chedeville and A. Lavoinne  | 693–696              |
| rat hepatocytes Glutamine metabolism in chick enterocytes: absence of pyrroline-5- carboxylase synthase and citrulline synthesis  | G. Wu, N.E. Flynn, W. Yan and D.G. Barstow, Jr.  | 717–721              |
| Cell wall polysaccharide biosynthesis and related metabolism in elicitor-stressed cells of French bean ( <i>Phaseolus vulgaris</i> L.)  | D. Robertson, B.A. McCormack and G.P. Bolwell  | 745–750              |
| Differential selenium-dependent expression of type I 5'-deiodinase and glutathione peroxidase in the porcine epithelial kidney cell line LLC-PK <sub>1</sub>  | M. Gross, M. Oertel and J. Köhrle  | 851–856              |
| Membranes and bioenergetics   |  |                      |
| Primary structure of rat plasma membrane Ca <sup>2+</sup> -ATPase isoform 4 and analysis of alternative splicing patterns at splice site A  | T.P. Keeton and G.E. Shull   | 779–785              |
| Localization and identification of Ca <sup>2+</sup> ATPases in highly purified human platelet plasma and intracellular membranes. Evidence that the monoclonal antibody PL/IM 430 recognizes the SERCA 3 Ca <sup>2+</sup> ATPase in human platelets | S. Bokkala, S.S. El-Daher, V.V. Kakkar, F. Wuytack and K.S. Authi  | 837–842              |
| Receptors and signal transduction   |  |                      |
| Receptor for <i>myo</i> -inositol trisphosphate from the microsomal fraction of <i>Vigna radiata</i>  | S. Biswas, B. Dalal, M. Sen and B.B. Biswas  | 631–636              |
| Physicochemical characterization of the cytoplasmic domain of the epidermal growth factor receptor and evidence for conformational changes associated with its activation by ammonium sulphate  | M. Gregoriou, P.F. Jones, J.F. Timms, J.J. Yang, S.E. Radford and A.R. Rees  | 667678               |
| Effects of caffeine on cytoplasmic free $Ca^{2+}$ concentration in pancreatic $\beta$ -cells are mediated by interaction with ATP-sensitive $K^+$ channels and L-type voltage-gated $Ca^{2+}$ channels but not the ryanodine receptor               | Md.S. Islam, O. Larsson, T. Nilsson and PO. Berggren   | 679–686              |
| Effects of CoA and acyl-CoA on Ca <sup>2+</sup> -permeability of endoplasmic-reticulum membranes from rat liver   | G.T. Rich, J.G. Comerford, S. Graham and A.P. Dawson   | 703–708              |
| Intracellular cross-talk between thyrotropin receptor and A <sub>1</sub> adenosine receptor in regulation of phospholipase C and adenylate cyclase in COS-7 cells transfected with their receptor genes   | F. Okajima, H. Tomura, K. Sho, M. Akbar, M.A. Majid and<br>Y. Kondo  | 70 <del>9</del> –715 |
| Adenylate cyclase, cyclic AMP and extracellular-signal-regulated kinase-2 in airway smooth muscle: modulation by protein kinase C and growth serum  | N. Moughal, P.A. Stevens, D. Kong, S. Pyne and N.J. Pyne   | 723–726              |
| The cross-regulation of $G_{\rm i}$ -protein by cholera toxin involves a phosphorylation by protein kinase A  | R. Levistre, M. Berguerand, G. Bereziat and J. Masliah   | 765–769              |
| Identification and characterization of the type-IVA cyclic AMP-<br>specific phosphodiesterase RD1 as a membrane-bound protein<br>expressed in cerebellum  | Y. Shakur, M. Wilson, L. Pooley, M. Lobban, S.L. Griffiths,<br>A.M. Campbell, J. Beattie, C. Daly and M.D. Houslay | 801—809              |
| Mutagenic structure/function analysis of the cytoplasmic cysteines of the insulin receptor  | S.L. Macaulay, M. Polites, M.J. Frenkel, D.R. Hewish and C.W. Ward   | 811–820              |
| Glucan receptor and zymosan-induced lysosomal enzyme secretion in macrophages   | H. Tapper and R. Sundler   | 829–835              |
| Free calcium transients in chemotactic and non-chemotactic strains of <i>Escherichia coli</i> determined by using recombinant aequorin  | N.J. Watkins, M.R. Knight, A.J. Trewavas and A.K. Campbell   | 865–869              |
| Cell biology and development  |  |                      |
| Lack of glycosyl-phosphatidylinositol anchoring leads to precursor retention by a unique mechanism in <i>Dictyostelium discoideum</i>   | P.C. Pauly and C. Klein  | 643–650              |
| The proximal pathway of metabolism of the chlorinated signal molecule differentiation-induced factor-1 (DIF-1) in the cellular slime mould <i>Dictyostelium</i>   | P. Morandini, J. Offer, D. Traynor, O. Nayler, D. Neuhaus, G.W. Taylor and R.R. Kay                                | 735–743              |
| Nitrone spin traps and a nitroxide antioxidant inhibit a common pathway of thymocyte apoptosis  | A.F.G. Slater, C.S.I. Nobel, E. Maellaro, J. Bustamante,<br>M. Kimland and S. Orrenius                             | 771–778              |

|  | Contents   | xiii    |
|--|--|---------|
| Mechanism of anti-proliferation caused by YC-1, an indazole derivative, in cultured rat A10 vascular smooth-muscle cells   | S.M. Yu, Z.J. Cheng, J.H. Guh, F.Y. Lee and S.C. Kuo                               | 787–792 |
| <b>BJ Letters</b> The two-gene model of the RH blood-group locus   | JP. Cartron, C. Le Van Kim, B. Cherif-Zahar, I. Mouro,<br>C. Rouillac and Y. Colin | 877–878 |
| Corrections  |  |         |
| Regulation of fos—lacZ fusion gene expression in primary mouse epidermal keratinocytes isolated from transgenic mice by W.B. Bollag, Y. Xiong, J. Ducote and C.S. Harmon (volume 300, pages 263–270, 1994)           |  | 879     |
| Multiple effects of haemin binding on protease susceptibility of bovine serum albumin and a novel isolation procedure for its large fragment by WS. Shin, H. Yamashita and M. Hirose (volume 304, pages 81–86, 1994) |  | 879     |
| Characterization of cDNA for murine tripeptidyl-peptidase II reveals alternative splicing by B. Tomkinson (volume 304, pages 517–523, 1994)  |  | 880     |