Review

Cell cycle regulation in *Aspergillus* by two protein kinases by S.A. Osmani and X.S. Ye

633-641

Biochemical Journal Volume 317, part 3 1 August 1996

| Research Communications | | |
|--|---|---------------------|
| Inhibition of protein synthesis in frog (<i>Xenopus laevis</i>) egg extracts by an antibody against phosphatidylinositol 4,5- bisphosphate | T.J. Keating, K. Fukami and K.R. Robinson | 643–64 |
| Alternative processing of the sarco/endoplasmic reticulum Ca ²⁺ -ATPase transcripts during muscle differentiation is a specifically regulated process | L. Van Den Bosch, L. Mertens, Y. Cavaloc, M. Peterson, F. Wuytack and J. Eggermont | 647–65 |
| Research Papers | | |
| Proteins | | |
| Cytokeratin 8 released by breast carcinoma cells <i>in vitro</i> binds plasminogen and tissue-type plasminogen activator and promotes plasminogen activation | T.A. Hembrough, K.R. Kralovich, L. Li and S.L. Gonias | 763–769 |
| Lack of integrin $lpha$ -chain endoproteolytic cleavage in furin-deficient human colon adenocarcinoma cells LoVo | M. Lehmann, V. Rigot, N.G. Seidah, J. Marvaldi and JC. Lissitzky | 803–809 |
| Calcium-dependent ADP-ribosylation of high-mobility-group I (HMGI) proteins | V. Giancotti, A. Bandiera, C. Sindici, L. Perissin and C. Crane-Robinson | 865–870 |
| Interactions between cytosolic components of the NADPH oxidase: $p40^{\rho hox}$ interacts with both $p67^{\rho hox}$ and $p47^{\rho hox}$ | F.B. Wientjes, G. Panayotou, E. Reeves and A.W. Segal | 919–92 |
| Human surfactant protein A with two distinct oligomeric structures which exhibit different capacities to interact with alveolar type II cells | A. Hattori, Y. Kuroki, H. Sohma, Y. Ogasawara and T. Akino | 939–944 |
| Enzymes | | |
| Baculovirus expression of two protein disulphide isomerase isoforms from $Caenorhabditis$ elegans and characterization of prolyl 4-hyydroxylases containing one of these polypeptides as their β subunit | J. Veijola, P. Annunen, P. Koivunen, A.P. Page, T. Pihlajaniemi and K.I. Kivirikko | 721–729 |
| Identification of the cysteine residues implicated in the formation of α_2 and α/β dimers of rat meprin | S. Chevallier, J. Ahn, G. Boileau and P. Crine | 731–73 |
| Endothelial-cell-stimulating angiogenesis factor (ESAF) activates progelatinase A (72 kDa type IV collagenase), prostromelysin 1 and procollagenase and reactivates their complexes with tissue inhibitors of metalloproteinases: a role for ESAF in non-inflammatory angiogenesis | B. McLaughlin and J.B. Weiss | 739–74 |
| Mechanism of the reaction catalysed by cytosolic 5'-nucleotidase/phosphotransferase: formation of a phosphorylated intermediate | C. Baiocchi, R. Pesi, M. Camici, R. Itoh and M. Grazia Tozzi | 797–80 ⁻ |
| Antibiotic interactions with the hammerhead ribozyme: tetracyclines as a new class of hammerhead inhibitor | J.B. Murray and J.R.P. Arnold | 855–860 |
| Characterization of pig liver glutathione S-transferases using HPLC-electrospray-ionization mass spectrometry | P. Rouimi, P. Anglade, L. Debrauwer and J. Tulliez | 879–884 |
| Role of an aprotinin-sensitive protease in the activation of Ca ²⁺ -ATPase by superoxide radical (O ₂ ⁻¹) in microsomes of pulmonary vascular smooth muscle | T. Chakraborti, S.K. Ghosh, J.R. Michael and S. Chakraborti | 885–890 |
| pH-dependent hysteretic behaviour of human myeloblastin (leucocyte proteinase 3) | A. Baici, S.E. Szedlacsek, H. Früh and B.A. Michel | 901–90 |
| Carbohydrates and lipids | | |
| Structural domains of heparan sulphate for specific recognition of the C-terminal heparin-binding domain of human plasma fibronectin (HEPII) | A. Walker and J.T. Gallagher | 871–877 |

| Gene structure and expression | | |
|---|---|---------|
| Heat shock selectively inhibits ribosomal RNA gene transcription and down-regulates E ₁ BF/Ku in mouse lymphosarcoma cells | K. Ghoshal and S.T. Jacob | 689–695 |
| Identification of distal silencing elements in the murine interferon- A11 gene promoter | P. Roffet, S. Lopez, S. Navarro, MT. Bandu, C. Coulombel, M. Vignal, J. Doly and G. Vodjdani | 697–706 |
| Identification of the transcription factors NF-YA and NF-YB as factors A and B that bound to the promoter of the major histocompatibility complex class II gene I-A β | A. Celada, S.R. McKercher and R.A. Maki | 771–778 |
| Differential expression of fibromodulin mRNA associated with tendon fibril growth: isolation and characterization of a chicken fibromodulin cDNA | M.V. Nurminskaya and D.E. Birk | 785–789 |
| Overproduction of stable ornithine decarboxylase and antizyme in the difluoromethylornithine-resistant cell line DH23b | J.L.A. Mitchell, Cy. Choe, G.G. Judd, D.J. Daghfal, R.J. Kurzeja and A. Leyser | 811–816 |
| Regulation of metabolism | | |
| Phosphorylation of phosphoenolpyruvate carboxykinase in plants. Studies in plants with C_4 photosynthesis and Crassulacean acid metabolism and in germinating seeds | R.P. Walker and R.C. Leegood | 653–658 |
| Uncoupling effect of polyunsaturated fatty acid deficiency in isolated rat hepatocytes: effect on glycerol metabolism | MA. Piquet, E. Fontaine, B. Sibille, C. Filippi, C. Keriel and X.M. Leverve | 667–674 |
| Modulation of intrahepatic cholesterol trafficking: evidence by in vivo antisense treatment for the involvement of sterol carrier protein-2 in newly synthesized cholesterol transport into rat bile | L. Puglielli, A. Rigotti, L. Amigo, L. Nuñez, A.V. Greco, M.J. Santos and F. Nervi | 681–687 |
| Flux control exerted by mitochondrial outer membrane carnitine palmitoyltransferase over β -oxidation, ketogenesis and tricarboxylic acid cycle actiity in hepatocytes isolated from rats in different metabolic states | L. Drynan, P.A. Quant and V.A. Zammit | 791–795 |
| Enzyme-to-enzyme channelling of Krebs cycle metabolic intermediates in Caco-2 cells exposed to [2-13C]propionate | W.J. Malaisse, TM. Zhang, I. Verbruggen and R. Willem | 861–863 |
| Inhibition of liver RNA breakdown during acute inflammation in the rat | A. Saadane, N. Neveux, G. Feldmann, B. Laredeux and F. Bleiberg-Daniel | 907–912 |
| Regulation of adenine nucleotide translocase and glycerol 3-phosphate dehydrogenase expression by thyroid hormones in different rat tissues | K. Dümmler, S. Müller and H.J. Seitz | 913–918 |
| Membranes and bioenergetics | | |
| Protection by chlorpromazine, albumin and bivalent cations against haemolysis induced by melittin, [Ala-14]melittin and whole bee venom | S.V. Rudenko and E.E. Nipot | 747–754 |
| Nucleoside uptake in rat liver parenchymal cells | J. Mercader, M. Gomez-Angelats, B. del Santo, F.J. Casado, A. Felipe and M. Pastor Anglada | 835–842 |
| Hypoxanthine enters human vascular endothelial cells (ECV 304) via the nitrobenzylthioinosine-insensitive equilibrative nucleoside transporter | N. Osses, J.D. Pearson, D.L. Yudilevich and S.M. Jarvis | 843–848 |
| Phosphatidylethanol stimulates the plasma-membrane calcium pump from human erythrocytes | M. Suju, M. Davila, G. Poleo, R. Docampo and G. Benaim | 933–938 |
| Receptors and signal transduction | | |
| Calcineurin is essential for DNA synthesis in Swiss 3T3 fibroblasts | M. Tomono, K. Toyoshima, M. Ito and H. Amano | 675–681 |
| The properties of a subtype of the inositol 1,4,5-trisphosphate receptor resulting from alternative splicing of the mRNA in the ligand-binding domain | JP. Lièvremont, H. Lancien, M. Hilly and JP. Mauger | 755–762 |
| Amplification of the thapsigargin-evoked increase in the cytosolic free Ca ²⁺ concentration by acetylcholine in acutely isolated mouse submandibular acinar cells | P.M. Smith and H.E. Reed | 779–783 |
| Immunological characterization of eristostatin and echistatin binding sites on $\alpha_{\rm llb}\beta_3$ and $\alpha_{\rm v}\beta_3$ integrins | C. Marcinkiewicz, L.A. Rosenthal, D.M. Mosser, T.J. Kunicki and S. Niewiarowski | 817–825 |

| Kinetics of the non-specific calcium leak from non-mitochondrial calcium stores in permeabilized A7r5 cells | L. Missiaen, H. De Smedt, J.B. Parys, L. Raeymaekers, G. Droogmans, L. Van Den Bosch and R. Casteels | 849–853 |
|---|---|---------|
| Purification of a rat neurotensin receptor expressed in <i>Escherichia coli</i> | J. Tucker and R. Grisshammer | 891–899 |
| Regulation of $\beta\text{-}\mathrm{adrenoceptor}$ density and mRNA levels in the rat heart cell-line H9c2 | V. Dangel, J. Giray, D. Ratge and H. Wisser | 925–931 |
| Insulin-responsive tissues contain the core complex protein SNAP-25 (synaptosomal-associated protein 25) A and B isoforms in addition to syntaxin 4 and synaptobrevins 1 and 2 | M.N. Jagadish, C.S. Fernandez, D.R. Hewish, S.L. Macaulay, K.H. Gough, J. Grusovin, A. Verkuylen, L. Cosgrove, A. Alafaci, M.J. Frenkel and C.W. Ward | 945–954 |
| Cell biology and development | | |
| Intracellular dissociation and reassembly of prolyl 4-hydroxylase: the α -subunits associate with the immunoglobulin-heavy-chain binding protein (BiP) allowing reassembly with the β -subunit | D.C.A. John and N.J. Bulleid | 659–665 |
| Reversal of the temperature-shift-induced growth restriction of a temperature-sensitive simian virus 40 T-antigen-transformed human fibroblast cell line by treatment with retinoic acid | YP. Tsao, SF. Li, SW. Kuo, JC. Liu and SL. Chen | 707–711 |
| Cytokine regulation of glutamate decarboxylase biosynthesis in isolated rat islets of Langerhans | R-S. Schmidli, B.E. Faulkner-Jones, L.C. Harrison, R.F.L. James and H.J. DeAizpurua | 713–719 |
| In vitro and in vivo induction of brown adipocyte uncoupling protein (thermogenin) by retinoic acid | P. Puigserver, F. Vázquez, M.L. Bonet, C. Picó and A. Palou | 827–833 |
| BJ Letters | | |
| Oil-bodies from sunflower (Helianthus annuus L.) seeds | F. Beisson, N. Ferte and G. Noat | 955–956 |
| Oleosins and oil bodies in plant seeds have postulated structures | C. Ratnayake and A.H.C. Huang | 956–958 |
| Corrections | | |
| Identification of putative ligand-binding sites of the integrin $\alpha 4\beta 1$ (VLA-4, CD49d/CD29) | | 959 |
| by T. Kamata, W. Puzon and T. Takada (volume 305, pages 945–951, 1995) | | |
| Purification of the synaptosomal plasma membrane $(Ca^{2+} + Mg^{2+})$ | | 959 |
| ATPase from pig brain | | |
| by J.M. Salvador and A.M. Mata (volume 315, pages 183–187, 1996) | | |
| CT-1 and Oct-2 DNA-binding site specificity is regulated in vitro | | 959 |
| by different kinases | | |

by S.J. Grenfell, D.S. Latchman and N.S.B. Thomas (volume 315, pages 889–893, 1996)

Contents

XIII