

Research Communications

- | | | |
|---|--|------|
| Functional characterization by heterologous expression of a novel cloned tachykinin peptide receptor | L.F. Donaldson, C.A. Haskell and M.R. Hanley | 1–5 |
| Evidence for two types of phosphate translocators in sweet-pepper (<i>Capsicum annuum</i> L.) fruit chromoplasts | W.P. Quick and H.E. Neuhaus | 7–10 |

Research Papers

Proteins

- | | | |
|---|--|---------|
| Stoichiometry of the subunits of flavocytochrome b_{558} of the NADPH oxidase of phagocytes | T.M. Wallach and A.W. Segal | 33–38 |
| Physical and kinetic characterization of recombinant human cholesteryl ester transfer protein | D.T. Connolly, J. McIntyre, D. Heuvelman, E.E. Remsen, R.E. McKinnie, L. Vu, M. Melton, R. Monsell, E.S. Krul and K. Glenn | 39–47 |
| LP2, a differentiation-associated lipid-binding protein expressed in bovine lens | C. Jaworski and G. Wistow | 49–54 |
| A 19 kDa protein secreted by the endometrium of the mare is a novel member of the lipocalin family | B. Crossett, T. Allen and F. Stewart | 137–143 |
| Biochemical characterization of the 49 kDa penicillin-binding protein of <i>Mycobacterium smegmatis</i> | T. Mukherjee, D. Basu, S. Mahapatra, C. Goffin, J. van Beeumen and J. Basu | 197–200 |

Enzymes

- | | | |
|---|--|---------|
| Proinsulin processing in the rat insulinoma cell line INS after overexpression of the endoproteases PC2 or PC3 by recombinant adenovirus | J.-C. Irminger, K. Meyer and P. Halban | 11–15 |
| Analysis of the interactions between an enzyme and multiple inhibitors using combination plots | E. Asante-Appiah and W.W.-C. Chan | 17–26 |
| Interaction of plant lipids with 14 kDa phospholipase A_2 enzymes | B.S. Vishwanath, W. Eichenberger, F.J. Frey and B.M. Frey | 93–99 |
| Production of the R2 subunit of ribonucleotide reductase from herpes simplex virus with prokaryotic and eukaryotic expression systems: higher activity of R2 produced by eukaryotic cells related to higher iron-binding capacity | N. Lamarche, G. Matton, B. Massie, M. Fontecave, M. Atta, F. Dumas, P. Gaudreau and Y. Langelier | 129–135 |
| New type of hexameric ornithine carbamoyltransferase with arginase activity in the cephamycin producers <i>Streptomyces clavuligerus</i> and <i>Nocardia lactamdurans</i> | J.L. de la Fuente, J.F. Martín and P. Liras | 173–179 |
| Identification of an active site cysteine residue in human type I $\text{Ins}(1,4,5)P_3$ 5-phosphatase by chemical modification and site-directed mutagenesis | D. Communi and C. Erneux | 181–186 |
| Kinetics of inactivation of bovine pancreatic ribonuclease A by bromopyruvic acid | M.-H. Wang, Z.-X. Wang and K.-Y. Zhao | 187–192 |
| Identification of the 4-amino analogue of tetrahydrobiopterin as a dihydropteridine reductase inhibitor and a potent pteridine antagonist of rat neuronal nitric oxide synthase | E.R. Werner, E. Pitters, K. Schmidt, H. Wachter, G. Werner-Felmayer and B. Mayer | 193–196 |
| Purification and biochemical characterization of a vacuolar serine endopeptidase induced by glucose starvation in maize roots | F. James, R. Brouquisse, C. Suire, A. Pradet and P. Raymond | 283–292 |
| Subcellular distribution of glycosylphosphatidylinositol-specific phospholipase D in rat liver | T. Hari, H. Kunze, E. Bohn, U. Brodbeck and P. Bütikofer | 315–319 |

Regulation of γ -glutamylcysteine synthetase by protein phosphorylation	W.-S. Sun, Z.-Z. Huang and S.C. Lu	321–328
Carbohydrates and lipids		
Molecular cloning and further characterization of rat peroxisomal trihydroxycoprostanoyl-CoA oxidase	E. Baumgart, J.C.T. Vanhooren, M. Fransen, F. van Leuven, H.D. Fahimi, P.P. van Veldhoven and G.P. Mannaerts	115–121
Regulation of metabolism		
Hypusine modification in eukaryotic initiation factor 5A in rodent cells selected for resistance to growth inhibition by ornithine decarboxylase-inhibiting drugs	M.E. Tome and E.W. Gerner	55–60
Impairment by interleukin 1β and tumour necrosis factor α of the glucagon-induced increase in phosphoenolpyruvate carboxykinase gene expression and gluconeogenesis in cultured rat hepatocytes	B. Christ and A. Nath	161–166
Construction of a P450c27 fusion enzyme: a useful tool for analysis of vitamin D ₃ 25-hydroxylase activity	F.J. Dilworth, S.M. Black, Y.-D. Guo, W.L. Miller and G. Jones	261–271
Ca ²⁺ stimulates both the respiratory and phosphorylation subsystems in rat heart mitochondria	V. Mildaziene, R. Baniene, Z. Nauciene, A. Marcinkeviciute, R. Morkuniene, V. Borutaite, B. Kholodenko and G.C. Brown	329–334
Membranes and bioenergetics		
Stimulation of the Ca ²⁺ -ATPase of sarcoplasmic reticulum by disulfiram	A.P. Starling, J.M. East and A.G. Lee	101–105
MOAT4, a novel multispecific organic-anion transporter for glucuronides and mercapturates in mouse L1210 cells and human erythrocytes	M. Saxena and G.B. Henderson	273–281
Effects of phosphatidylethanolamines on the activity of the Ca ²⁺ -ATPase of sarcoplasmic reticulum	A.P. Starling, K.A. Dalton, J.M. East, S. Oliver and A.G. Lee	309–314
Receptors and signal transduction		
Interaction between the two signal transduction systems of the histamine H2 receptor: desensitizing and sensitizing effects of histamine stimulation on histamine-dependent cAMP production in Chinese hamster ovary cells	Y. Fukushima, T. Asano, H. Katagiri, M. Aihara, T. Saitoh, M. Anai, M. Funaki, T. Ogihara, K. Inukai, N. Matsuhashi, Y. Oka, Y. Yazaki and K. Sugano	27–32
Association between phosphatidylinositol-3 kinase, Cbl and other tyrosine phosphorylated proteins in colony-stimulating factor-1-stimulated macrophages	V. Kanagasundaram, A. Jaworowski and J.A. Hamilton	69–77
A constitutively active mutant of the α_{1B} -adrenergic receptor can cause greater agonist-dependent down-regulation of the G-proteins G _q α and G ₁₁ α than the wild-type receptor	T.W. Lee, A. Wise, S. Cotecchia and G. Milligan	79–86
Bombesin and thrombin affect discrete pools of intracellular calcium through different G-proteins	J.-L. Wang, S. Kalyanaraman, M. De Vivo and N. Gautam	87–91
Treatment of vascular smooth muscle cells with antisense phosphorothioate oligodeoxynucleotides directed against p42 and p44 mitogen-activated protein kinases abolishes DNA synthesis in response to platelet-derived growth factor	C.J.M. Robinson, P.H. Scott, A.B. Allan, T. Jess, G.W. Gould and R. Plevin	123–127
Activation of mitogen-activated protein kinase couples neurotensin receptor stimulation to induction of the primary response gene <i>Krox-24</i>	C. Poinot-Chazel, M. Portier, M. Bouaboula, N. Vita, F. Pecceu, D. Gully, J.G. Monroe, J.-P. Maffrand, G. Le Fur and P. Casellas	145–151
Regulation of natural killer cell-mediated cytotoxicity by serine/threonine phosphatases: identification of a calyculin A-sensitive serine/threonine kinase	A. Bajpai and Z. Brahmi	153–159
Calcium-dependent activation of Erk-1 and Erk-2 after hypo-osmotic astrocyte swelling	F. Schliess, R. Sinning, R. Fischer, C. Schmalenbach and D. Häussinger	167–171
Inhibition of insulin release by synthetic peptides shows that the H3 region at the C-terminal domain of syntaxin-1 is crucial for Ca ²⁺ but not for guanosine 5'-[γ -thio]triphosphate-induced secretion	F. Martin, E. Salinas, J. Vazquez, B. Soria and J.A. Reig	201–205

Characterization of two forms of protein kinase C α , with different substrate specificities, from skeletal muscle	C. Schmitz-Peiffer, C.L. Browne and T. J. Biden	207–214
Desensitization of P _{2U} receptor in neuronal cell line. Different control by the agonists ATP and UTP, as demonstrated by single-cell Ca ²⁺ responses	U. Czubayko and G. Reiser	215–219
Phosphorylation and activation of p42 and p44 mitogen-activated protein kinase are required for the P2 purinoceptor stimulation of endothelial prostacyclin production	V. Patel, C. Brown, A. Goodwin, N. Wilkie and M.R. Boarder	221–226
Agonist activation of p42 and p44 mitogen-activated protein kinases following expression of the mouse δ opioid receptor in Rat-1 fibroblasts: effects of receptor expression levels and comparisons with G-protein activation	A.R. Burt, I.C. Carr, I. Mullaney, N.G. Anderson and G. Milligan	227–235
Kinetic and biochemical correlation between sustained p44ERK1 (44 kDa extracellular signal-regulated kinase 1) activation and lysophosphatidic acid-stimulated DNA synthesis in Rat-1 cells	S.J. Cook and F. McCormick	237–245
Cleavage of the thrombin receptor: identification of potential activators and inactivators	M.A.A. Parry, T. Myles, J. Tschopp and S.R. Stone	335–341
Cell biology and development		
Importance of catalase in the adaptive response to hydrogen peroxide: analysis of acatalasaemic <i>Saccharomyces cerevisiae</i>	S. Izawa, Y. Inoue and A. Kimura	61–67
Changes in mRNA levels of the sarcoplasmic/endoplasmic-reticulum Ca ²⁺ -ATPase isoforms in the rat soleus muscle regenerating from notexin-induced necrosis	E. Zádor, L. Mendler, M. Ver Heyen, L. Dux and F. Wuytack	107–113
Evidence for a novel cytoplasmic tRNA–protein complex containing the KH-multidomain protein vigilin	C. Kruse, A. Grünweller, H. Notbohm, S. Kügler, W.G. Purschke and P.K. Müller	247–252
Extracellular high-mobility group 1 protein is essential for murine erythroleukaemia cell differentiation	B. Sparatore, M. Passalacqua, M. Patrone, E. Melloni and S. Pontremoli	253–256
Changes in the nuclear matrix of chicken erythrocytes that accompany maturation	H.Y. Chen, J.-M. Sun, M.J. Hendzel, J.B. Rattner and J.R. Davie	257–265
Internalization and catabolism of radiolabelled antibodies to the MHC class-II invariant chain by B-cell lymphomas	H.J. Hansen, G.L. Ong, H. Diril, A. Valdez, P.A. Roche, G.L. Griffiths, D.M. Goldenberg and M.J. Mattes	293–300
A steady-state labelling approach to the measurement of proteoglycan turnover <i>in vivo</i> and its application to glomerular proteoglycans	E.L. Akuffo, J.R. Hunt, J. Moss, D. Woodrow, M. Davies and R.M. Mason	301–308
BJ Letters		
Fatty acyl-CoA esters and the permeability of rat liver microsomal vesicles	G. Bánhegyi, M. Csala, J. Mandl, A. Burchell, B. Burchell, P. Marcolongo, R. Fulceri and A. Benedetti	343–344