

**Review**

Regulatory mechanisms that modulate signalling by G-protein-coupled receptors  
by S. Böhm, E.F. Grady and N.W. Bunnett

1–18

**Biochemical Journal****Volume 322, part 1****15 February 1997****Research Communications**

Activation of CPP32 and Mch3 $\alpha$  in wild-type p53-induced apoptosis

Electrospray mass-spectrometric, spectrophotometric and electrochemical methods do not provide evidence for the binding of nitric oxide by pyocyanine at pH 7

Diphospho-myo-inositol phosphates from *Dictyostelium* identified as D-6-diphospho-myo-inositol pentakisphosphate and D-5,6-bisdiphospho-myo-inositol tetrakisphosphate

J.M. Chandler, E.S. Alnemri, G.M. Cohen and M. MacFarlane

19–23

D.V. Vukomanovic, D.E. Zoutman, J.A. Stone, G.S. Marks, J.F. Brien and K. Nakatsu

25–29

T. Laussmann, K.M. Reddy, K.K. Reddy, J.R. Falck and G. Vogel

31–33

**Research Papers****Proteins**

Expression and characterization of *Pseudomonas aeruginosa* cytochrome c-551 and two site-directed mutants: role of tryptophan 56 in the modulation of redox properties

The variant-specific surface protein of *Giardia*, VSP4A1, is a glycosylated and palmitoylated protein

Difference in hepatic metallothionein content in Antarctic red-blooded and haemoglobinless fish: undetectable metallothionein levels in haemoglobinless fish is accompanied by accumulation of untranslated metallothionein mRNA

Biosynthesis of 'essential' amino acids by scleractinian corals

Identification of an N-capping box that affects the  $\alpha$ 6-helix propensity in glutathione S-transferase superfamily proteins: a role for an invariant aspartic residue

Highly conserved residue arginine-15 is required for the Ca<sup>2+</sup>-dependent properties of the  $\gamma$ -carboxyglutamic acid domain of human anticoagulation Protein C and activated Protein C

Quantification of malondialdehyde and 4-hydroxynonenal adducts to lysine residues in native and oxidized human low-density lipoprotein

Secretion of a type II integral membrane protein induced by mutation of the transmembrane segment

F. Cutruzzolà, I. Ciabatti, G. Rolli, S. Falcinelli, M. Arese, G. Ranghino, A. Anselmino, E. Zennaro and M.C. Silvestrini

35–42

P. Papanastasiou, M.J. McConville, J. Ralton and P. Köhler

49–56

R. Scudiero, V. Carginale, M. Riggio, C. Capasso, A. Capasso, P. Kille, G. di Prisco and E. Parisi

207–211

L.M. FitzGerald and A.M. Szmant

213–221

A. Aceto, B. Dragani, S. Melino, N. Allocati, M. Masulli, C. Di Ilio and R. Petruzzelli

229–234

A. Thariath and F.J. Castellino

309–315

J.R. Requena, M.X. Fu, M.U. Ahmed, A.J. Jenkins, T.J. Lyons, J.W. Baynes and S.R. Thorpe

317–325

I. Lemire, C. Lazare, P. Crine and G. Boileau

335–342

**Enzymes**

Glutathione-dependent activities of *Trypanosoma cruzi* p52 makes it a new member of the thiol:disulphide oxidoreductase family

Structural instability of mutant  $\beta$ -cell glucokinase: implications for the molecular pathogenesis of maturity-onset diabetes of the young (type-2)

Telokin (kinase-related protein) modulates the oligomeric state of smooth-muscle myosin light-chain kinase and its interaction with myosin filaments

Involvement of two basic residues (Lys-270 and Arg-276) of human liver 3 $\alpha$ -hydroxysteroid dehydrogenase in NAD(P)H binding and activation by sulphobromophthalein: site-directed mutagenesis and kinetic analysis

Epoxyethylglycyl peptides as inhibitors of oligosaccharyltransferase: double-labelling of the active site

M. Moutiez, E. Quéméneur, C. Sergheraert, V. Lucas, A. Tartar and E. Davioud-Charvet

43–48

P. Kesavan, L. Wang, E. Davis, A. Cuesta, I. Sweet, K. Niswender, M.A. Magnuson and F.M. Matschinsky

57–63

K. Nieznanski and A. Sobieszek

65–71

K. Matsuura, Y. Tamada, K. Sato, H. Iwasa, G. Miwa, Y. Deyashiki and A. Hara

89–93

E. Bause, M. Wesemann, A. Bartoschek and W. Breuer

95–102

---

Identification of a putative flexible loop in <i>Arabidopsis</i> glutathione synthetase	C.-L. Wang and D.J. Oliver	<b>241–244</b>
<b>Carbohydrates and lipids</b>		
Effects of charged lipids on the interaction of cholesteryl ester transfer protein with lipid microemulsions	O.V. Rajaram and W.H. Sawyer	<b>159–165</b>
Metabolism of 27-, 25- and 24-hydroxycholesterol in rat glial cells and neurons	J. Zhang, Y. Akwa, M. El-Etr, E.-E. Baulieu and J. Sjövall	<b>175–184</b>
<b>Gene structure and expression</b>		
The octamer-binding proteins Oct-1 and Oct-2 repress the HIV long terminal repeat promoter and its transactivation by Tat	Y.-Z. Liu and D.S. Latchman	<b>155–158</b>
Regulation of the human $\alpha 2(1)$ procollagen gene by sequences adjacent to the CCAAT box	M. Collins, V.D. Leaner, M. Madikizela and M.I. Parker	<b>199–206</b>
Regulation by fatty acids of angiotensinogen gene expression in preadipose cells	I. Safonova, J. Aubert, R. Negrel and G. Ailhaud	<b>235–239</b>
Structure of active chromatin: isolation and characterization of transcriptionally active chromatin from rat liver	K. Tikoo, S. Gupta, A.A. Hamid, V. Shah, B. Chatterjee and Z. Ali	<b>273–279</b>
Structure of active chromatin: covalent modifications of histones in active and inactive genes of control and hypothyroid rat liver	K. Tikoo and Z. Ali	<b>281–287</b>
Structure of active chromatin: higher-order folding of transcriptionally active chromatin in control and hypothyroid rat liver	K. Tikoo, Q.A. Hamid and Z. Ali	<b>289–296</b>
CCAAT-enhancer-binding protein $\alpha$ (C/EBP $\alpha$ ) is required for the thyroid hormone but not the retinoic acid induction of phosphoenolpyruvate carboxykinase (PEPCK) gene transcription	E.A. Park, S. Song, M. Olive and W.J. Roesler	<b>343–349</b>
<b>Regulation of metabolism</b>		
Mitochondrial creatine kinase isoform expression does not correlate with its mode of action	K. Anflous, V. Veksler, P. Mateo, F. Samson, V. Saks and R. Ventura-Clapier	<b>73–78</b>
Finite change analysis of glycolytic intermediates in tuber tissue of lines of transgenic potato ( <i>Solanum tuberosum</i> ) overexpressing phosphofructokinase	S. Thomas, P.J.F. Mooney, M.M. Burrell and D.A. Fell	<b>111–117</b>
Metabolic Control Analysis of glycolysis in tuber tissue of potato ( <i>Solanum tuberosum</i> ): explanation for the low control coefficient of phosphofructokinase over respiratory flux	S. Thomas, P.J.F. Mooney, M.M. Burrell and D.A. Fell	<b>119–127</b>
Transient inactivation of phosphatidylethanolamine <i>N</i> -methyltransferase-2 and activation of cytidine triphosphate:phosphocholine cytidylyltransferase during non-neoplastic liver growth	L. Tessitore, Z. Cui and D.E. Vance	<b>151–154</b>
Pathways of peroxynitrite oxidation of thiol groups	C. Quijano, B. Alvarez, R.M. Gatti, O. Augusto and R. Radi	<b>167–173</b>
Nitric oxide stimulates glucose transport and metabolism in rat skeletal muscle <i>in vitro</i>	M.E. Young, G.K. Radda and B. Leighton	<b>223–228</b>
CGP 48664, a potent and specific S-adenosylmethionine decarboxylase inhibitor: effects on regulation and stability of the enzyme	F. Svensson, H. Mett and L. Persson	<b>297–302</b>
Regulation of glycogen synthase and phosphorylase during recovery from high-intensity exercise in the rat	L. Bräu, L.D.M.C.B. Ferreira, S. Nikolovski, G. Raja, T.N. Palmer and P.A. Fournier	<b>303–308</b>
<b>Membranes and bioenergetics</b>		
Calcium-dependence of synxin binding may determine aggregation and fusion of lamellar bodies	N. Sen, A.R. Spitzer and A. Chander	<b>103–109</b>
Evidence for $P_2$ -purinoceptor-mediated uptake of $Ca^{2+}$ across a fish ( <i>Oreochromis mossambicus</i> ) intestinal brush border membrane	P.H.M. Klaren, S.E. Wendelaar Bonga and G. Flik	<b>129–134</b>
The plasma-membrane $Ca^{2+}$ -ATPase of <i>Leishmania donovani</i> is an extrusion pump for $Ca^{2+}$	D. Mandal, T. Mukherjee, S. Sarkar, S. Majumdar and A. Bhaduri	<b>251–257</b>

---

Functional expression and apical localization of the cystic fibrosis transmembrane conductance regulator in MDCK I cells	A. Mohamed, D. Ferguson, F.S. Seibert, H.-M. Cai, N. Kartner, S. Grinstein, J.R. Riordan and G.L. Lukacs	259–265
<b>Receptors and signal transduction</b>		
cAMP suppresses p21 <sup>ras</sup> and Raf-1 responses but not the Erk-1 response to granulocyte-colony-stimulating factor: possible Raf-1-independent activation of Erk-1	X.F. Csar, A.C. Ward, B.W. Hoffmann, G.G. Guy and J.A. Hamilton	79–87
Expression of leptin and $\beta_3$ -adrenergic receptors in rat adipose tissue in altered thyroid states	J.N. Fain, E.C. Coronel, M.J. Beauchamp and S.W. Bahouth	145–150
Regulation of the activity of MEK kinase (MEKK1) by autophosphorylation within the kinase activation domain	J.C. Deak and D.J. Templeton	185–192
Analysis of mitogen-activated protein kinase activation by naturally occurring splice variants of TrkB, the receptor for neurotrophin-3	F.J. Gunn-Moore, A.G. Williams and J.M. Tavaré	192–198
Involvement of the Ca <sup>2+</sup> /calmodulin-dependent protein kinase II pathway in the Ca <sup>2+</sup> -mediated regulation of the capacitative Ca <sup>2+</sup> entry in <i>Xenopus</i> oocytes	F. Matifat, F. Fournier, T. Lorca, J. P. Capony, G. Brûlé and T. Collin	267–272
Sphingosylphosphocholine modulates the ryanodine receptor/calcium-release channel of cardiac sarcoplasmic reticulum membranes	R. Betto, A. Teresi, F. Turcato, G. Salviati, R.A. Sabbadini, K. Krown, C.C. Glembotski, L.A. Kindman, C. Dettbarn, Y. Pereon, K. Yasui and P.T. Palade	327–333
The CD46 transmembrane domain is required for efficient formation of measles-virus-mediated syncytium	T. Seya, M. Kurita, K. Iwata, Y. Yanagi, K. Tanaka, K. Shida, M. Hatanaka, M. Matsumoto, S. Jun, A. Hirano, S. Ueda and S. Nagasawa	135–144
Chick osteocyte-derived protein inhibits osteoclastic bone resorption	A. Maejima-Ikeda, M. Aoki, K. Tsuritani, K. Kamioka, K. Hiura, T. Miyoshi, H. Hara, T. Takano-Yamamoto and M. Kumegawa	245–250
<b>BJ Letters</b>		
Enzyme–substrate interaction in the catalytic triad of serine proteases: increase in the $pK_a$ of Asp <sup>102</sup>	H. Neuvonen	351–352