

**Review**

Phospholipid transfer proteins revisited  
by K. W. A. Wirtz

353–360

**Research Communication**

Activation of pro-caspase-7 by serine proteases includes a non-canonical specificity

Q. Zhou and G.S. Salvesen

361–364

**Research Papers****Proteins**

Isolation of peptides from phage-displayed random peptide libraries that interact with the talin-binding domain of vinculin *N*<sup>ε</sup>-(Carboxyethyl)lysine, a product of the chemical modification of proteins by methylglyoxal, increases with age in human lens proteins

N.B. Adey and B.K. Kay

523–528

Influence of the conserved disulphide bond, exposed to the putative binding pocket, on the structure and function of the immunoglobulin-like molecular chaperone Caf1M of *Yersinia pestis*

V.P. Zav'yalov, T.V. Chernovskaya, D.A.G. Chapman, A.V. Karlyshev, S. MacIntyre, A.V. Zavalov, A.M. Vasiliev, A.I. Denesyuk, G.A. Zav'yalova, I.V. Dudich, T. Korpela and V.M. Abramov

571–578

Selective binding of *N*-acetylglucosamine to the chicken hepatic lectin

L. Burrows, S.T. Iobst and K. Drickamer

673–680

**Enzymes**

Inactivation kinetics of dihydrofolate reductase from Chinese hamster during urea denaturation

J.-W. Wu, Z.-X. Wang and J.-M. Zhou

395–401

Reactions of the oxidized organic cofactor in copper-depleted bovine serum amine oxidase

E. Agostinelli, G. De Matteis, A. Sinibaldi, B. Mondovì and L. Morpurgo

497–501

pH-dependence for binding a single nitrite ion to each type-2 copper centre in the copper-containing nitrite reductase of *Alcaligenes xylosoxidans*

Z.H.L. Abraham, B.E. Smith, B.D. Howes, D.J. Lowe and R.R. Eady

511–516

Structural features that make oligopeptides susceptible substrates for hydrolysis by recombinant thimet oligopeptidase

A.C.M. Camargo, M.D. Gomes, A.P. Reichl, E.S. Ferro, S. Jacchieri, I.Y. Hirata and L. Juliano

517–522

Effects of epidermal growth factor and insulin on the activity of *N*-acetylglucosaminyltransferase V

Q. Wang, D. Zhou, D. Shao, Z. Shen and J. Gu

543–545

Purification of a dichlorophenol-indophenol oxidoreductase from rat and bovine synaptic membranes: tight complex association of a glyceraldehyde-3-phosphate dehydrogenase isoform, TOAD64, enolase- $\gamma$  and aldolase C

C. Bulliard, R. Zurbriggen, J. Tornare, M. Faty, Z. Dastoor and J.-L. Dreyer

555–563

Structural and functional analysis of a glutathione S-transferase from *Ascaris suum*

E. Liebau, V.H.O. Eckelt, G. Wildenburg, P. Teesdale-Spittle, P.M. Brophy, R.D. Walter and K. Henkle-Dührsen

659–666

**Carbohydrates and lipids**

Phospholipid metabolism of serine in *Plasmodium*-infected erythrocytes involves phosphatidylserine and direct serine decarboxylation

N. Elabbadi, M.L. Ancelin and H.J. Vial

435–445

Nitric oxide degradation of heparin and heparan sulphate

R.E. Vilar, D. Ghael, M. Li, D.D. Bhagat, L.M. Arrigo, M.K. Cowman, H.S. Dweck and L. Rosenfeld

473–479

**Gene structure and expression**

Existence of two translation initiation sites leading to the expression of two proteins from the rat high-affinity neurotensin-receptor cDNA: possible regulation by the 5' end non-coding region

J.-M. Botto, J.-P. Vincent and J. Mazella

389–393

Two genes encoding an endoglucanase and a cellulose-binding protein are clustered and co-regulated by a TTA codon in *Streptomyces halstedii* JM8

A.L. Garda, J.M. Fernández-Abalos, P. Sánchez, A. Ruiz-Arribas and R.I. Santamaría

403–411

Differential functional significance of AP-1 binding sites in the promoter of the gene encoding mouse tissue inhibitor of metalloproteinases-3	H. Kim, W.D. Pennie, Y. Sun and N.H. Colburn	<b>547–553</b>
Transcriptional activity of the human tissue inhibitor of metalloproteinases 1 (TIMP-1) gene in fibroblasts involves elements in the promoter, exon 1 and intron 1	I.M. Clark, A.D. Rowan, D.R. Edwards, T. Bech-Hansen, D.A. Mann, M.J. Bahr and T.E. Cawston	<b>611–617</b>
A soluble 3-hydroxy-3-methylglutaryl-CoA reductase in the protozoan <i>Trypanosoma cruzi</i>	J. Peña-Díaz, A. Montalvetti, A. Camacho, C. Gallego, L.M. Ruiz-Perez and D. Gonzalez-Pacanowska	<b>619–626</b>
Genetic information 'created' by archaebacterial DNA polymerase	N. Ogata and T. Miura	<b>667–671</b>
<b>Regulation of metabolism</b>		
Expression of an exogenous isopentenyl diphosphate isomerase gene enhances isoprenoid biosynthesis in <i>Escherichia coli</i>	S. Kajiwara, P.D. Fraser, K. Kondo and N. Misawa	<b>421–426</b>
Recombinant fish neurotrophin-6 is a heparin-binding glycoprotein: implications for a role in axonal guidance	X. Li, J. Franz, F. Lottspeich and R. Götz	<b>461–466</b>
ADP-glucose drives starch synthesis in isolated maize endosperm amyloplasts: characterization of starch synthesis and transport properties across the amyloplast envelope	T. Möhlmann, J. Tjaden, G. Henrichs, W.P. Quick, R. Häusler and H.E. Neuhaus	<b>503–509</b>
Triacsin C blocks <i>de novo</i> synthesis of glycerolipids and cholesterol esters but not recycling of fatty acid into phospholipid: evidence for functionally separate pools of acyl-CoA	R.A. Igal, P. Wang and R.A. Coleman	<b>529–534</b>
Insulin and insulin-like growth factor 1 antagonize the stimulation of <i>ob</i> gene expression by dexamethasone in cultured rat adipose tissue	B.A. Reul, L.N. Ongemba, A.-M. Pottier, J.-C. Henquin and S.M. Brichard	<b>605–610</b>
<b>Membranes and bioenergetics</b>		
Resolution of two ADP-ribosylation factor 1 GTPase-activating proteins from rat liver	P.A. Randazzo	<b>413–419</b>
Cloning of the monocarboxylate transporter isoform MCT2 from rat testis provides evidence that expression in tissues is species-specific and may involve post-transcriptional regulation	V.N. Jackson, N.T. Price, L. Carpenter and A.P. Halestrap	<b>447–453</b>
Expression of the surface antigen 4F2hc affects system-L-like neutral-amino acid-transport activity in mammalian cells	S. Bröer, A. Bröer and B. Hamprecht	<b>535–541</b>
Asymmetrical distribution of cardiolipin in yeast inner mitochondrial membrane triggered by carbon catabolite repression	P.F. Gallet, J.-M. Petit, A. Maftah, A. Zachowski and R. Julien	<b>627–634</b>
Properties of a cysteine-free proton-pumping nicotinamide nucleotide transhydrogenase	J. Meuller, J. Zhang, C. Hou, P.D. Bragg and J. Rydström	<b>681–687</b>
<b>Receptors and signal transduction</b>		
Activation of mitogen-activated protein kinase and p70S6 kinase is not correlated with cerebellar granule cell survival	F.J. Gunn-Moore, A.G. Williams, N.J. Toms and J.M. Tavaré	<b>365–369</b>
Synthesis and properties of the very-low-density-lipoprotein receptor and a comparison with the low-density-lipoprotein receptor	D.D. Patel, R.A. Forder, A.K. Soutar and B.L. Knight	<b>371–377</b>
Interaction between cAMP-dependent and insulin-dependent signal pathways in tyrosine phosphorylation in primary cultures of rat hepatocytes	Y. Ito, Y. Uchijima, M. Ariga, T. Seki, A. Takenaka, F. Hakuno, S.-I. Takahashi, T. Ariga and T. Noguchi	<b>379–388</b>
Latent transforming growth factor- $\beta$ complex in Chinese hamster ovary cells contains the multifunctional cysteine-rich fibroblast growth factor receptor, also termed E-selectin-ligand or MG-160	A. Olofsson, U. Hellman, P. Ten Dijke, S. Grimsby, H. Ichijo, A. Morén, K. Miyazono and C.-H. Heldin	<b>427–434</b>
Tight links between adenine and guanine nucleotide pools in mouse pancreatic islets: a study with mycophenolic acid	P. Detimary, C. Xiao and J.-C. Henquin	<b>467–471</b>
Sphingosine 1-phosphate stimulates Rho-mediated tyrosine phosphorylation of focal adhesion kinase and paxillin in Swiss 3T3 fibroblasts	F. Wang, C.D. Nobes, A. Hall and S. Spiegel	<b>481–488</b>
Lipid kinase and protein kinase activities of G-protein-coupled phosphoinositide 3-kinase $\gamma$ : structure–activity analysis and interactions with wortmannin	S. Stoyanova, G. Bulgarelli-Leva, C. Kirsch, T. Hanck, R. Klingler, R. Wetzker and M.P. Wymann	<b>489–495</b>

---

Membrane association, localization and topology of rat inositol 1,4,5-trisphosphate 3-kinase B: implications for membrane traffic and $\text{Ca}^{2+}$ homeostasis	S. Soriano, S. Thomas, S. High, G. Griffiths, C. D'Santos, P. Cullen and G. Banting	<b>579–589</b>
Interaction of truncated human interferon $\gamma$ variants with the interferon $\gamma$ receptor: crucial importance of Arg-129	J. Haelewyn, L. Michiels, P. Verhaert, M.F. Hoylaerts, R. Witters and M. De Ley	<b>591–595</b>
Dynamic equilibrium between calcineurin and kinase activities regulates the phosphorylation state and localization of the nuclear factor of activated T-cells	J.E. Scott, V.A. Ruff and K.L. Leach	<b>597–603</b>
Phospholipase C inhibitor, U73122, releases intracellular $\text{Ca}^{2+}$ , potentiates $\text{Ins}(1,4,5)\text{P}_3$ -mediated $\text{Ca}^{2+}$ release and directly activates ion channels in mouse pancreatic acinar cells	H. Mogami, C. Lloyd Mills and D.V. Gallacher	<b>645–651</b>
Interleukin 1-induced calcium signalling in chondrocytes requires focal adhesions	L. Luo, T. Cruz and C. McCulloch	<b>653–658</b>
Cross-linking analysis of the ryanodine receptor and $\alpha_1$ -dihydropyridine receptor in rabbit skeletal muscle triads	B.E. Murray and K. Ohlendieck	<b>689–696</b>
<b>Cell biology and development</b>		
Heterologous expression of rab4 reduces glucose transport and GLUT4 abundance at the cell surface in oocytes	S. Mora, I. Monden, A. Zorzano and K. Keller	<b>455–459</b>
Subcellular structure containing mRNA for $\beta$ subunit of mitochondrial $\text{H}^+$ -ATP synthase in rat hepatocytes is translationally active	J. Ricart, G. Egea, J.M. Izquierdo, C. San Martín and J.M. Cuezva	<b>635–643</b>