

**Biochemical Journal**  
**Volume 328, part 1**  
**15 November 1997**

**Review**

**The DNA damage-recognition problem in human and other eukaryotic cells: the XPA damage binding protein**  
*by J.E. Cleaver and J.C. States*

1–12

**Research Papers****Proteins**

- |   |   |         |
|---|---|---------|
| Metallothionein accretion in human hepatic cells is linked to cellular proliferation  | R. Studer, C.P. Vogt, M. Cavigelli, P.E. Hunziker and J.H.R. Kägi   | 63–67   |
| Calnexin and calreticulin bind to enzymically active tissue-type plasminogen activator during biosynthesis and are not required for folding to the native conformation  | S. Allen and N.J. Bulleid   | 113–119 |
| Recombinant two-iron rubredoxin of <i>Pseudomonas oleovorans</i> : overexpression, purification and characterization by optical, CD and $^{13}\text{C}$ NMR spectroscopies                                    | H.J. Lee, L.-Y. Lian and N.S. Scrutton  | 131–136 |
| N-terminal stretch Arg <sup>2</sup> , Arg <sup>3</sup> , Arg <sup>4</sup> and Arg <sup>5</sup> of human lactoferrin is essential for binding to heparin, bacterial lipopolysaccharide, human lysozyme and DNA | P.H.C. van Berkel, M.E.J. Geerts, H.A. van Veen, M. Mericskay, H.A. de Boer and J.H. Nuijens                  | 145–151 |
| Intracellular coupling of bikunin and the heavy chain of rat pre- $\alpha$ -inhibitor in COS-1 cells  | A.M. Blom, M. Thuveson and E. Fries   | 185–191 |
| Location and functional characterization of myosin contact sites in smooth muscle caldesmon   | A.V. Vorotnikov, S.B. Marston and P.A.J. Huber  | 211–218 |
| Primary structure of a copper-binding metallothionein from mantle tissue of the terrestrial gastropod <i>Helix pomatia</i> L.   | B. Berger, R. Dallinger, P. Gehrig and P.E. Hunziker  | 219–224 |
| Expression of hydrophilic surfactant proteins by mesentery cells in rat and man   | B. Chailley-Heu, S. Rubio, J.-P. Rougier, R. Ducroc, A.-M. Barlier-Mur, P. Ronco and J.R. Bourbon             | 251–256 |
| Evolution of phosphagen kinase. Isolation, characterization and cDNA-derived amino acid sequence of two-domain arginine kinase from the sea anemone <i>Anthopleura japonicus</i>                              | T. Suzuki, Y. Kawasaki and T. Furukohri   | 301–306 |
| A four-disulphide-bridged toxin, with high affinity towards voltage-gated K <sup>+</sup> channels, isolated from <i>Heterometrus spinifer</i> (Scorpionidae) venom  | B. Lebrun, R. Romi-Lebrun, M.-F. Martin-Eauclaire, A. Yasuda, M. Ishiguro, Y. Oyama, O. Pongs and T. Nakajima | 321–327 |

**Enzymes**

- |   |  |         |
|---|--|---------|
| Cloning and overexpression of rat kidney biliverdin IX $\alpha$ reductase as a fusion protein with glutathione S-transferase: stereochemistry of NADH oxidation and evidence that the presence of the glutathione S-transferase domain does not effect BVR-A activity | O. Ennis, R. Maytum and T.J. Mantle  | 33–36   |
| Molecular cloning and expression of a rat hepatic multiple inositol polyphosphate phosphatase   | A. Craxton, J.J. Caffrey, W. Burkhardt, S.T. Safrany and S.B. Shears                 | 75–81   |
| Kinetic mechanism of the glycogen-phosphorylase-catalysed reaction in the direction of glycogen synthesis: co-operative interactions of AMP and glucose 1-phosphate during catalysis  | E.A. Sergienko and D.K. Srivastava   | 83–91   |
| Expression and processing of vertebrate acetylcholinesterase in the yeast <i>Pichia pastoris</i>  | N. Morel and J. Massoulié  | 121–129 |
| Expression, purification, and characterization of recombinant human glutamine synthetase  | C.D. Listrom, H. Morizono, B.S. Rajagopal, M.T. McCann, M. Tuchman and N.M. Allewell | 159–163 |
| Identification of essential histidine residues in UDP-N-acetyl-D-galactosamine: polypeptide N-acetylgalactosaminyltransferase-T1  | S. Wragg, F.K. Hagen and L.A. Tabak  | 193–197 |
| 14,15-Dehydroleukotriene A <sub>4</sub> : a specific substrate for leukotriene C <sub>4</sub> synthase  | A. Sala, M. Garcia, S. Zarini, J.C. Rossi, G. Folco and T. Durand                    | 225–229 |
| Mutagenesis of residue 157 in the active site of human glyoxalase I   | M. Ridderström, A.D. Cameron, T.A. Jones and B. Mannervik                            | 231–235 |

Adenosine 5'-tetraphosphate phosphohydrolase from yellow lupin seeds: purification to homogeneity and some properties	A. Guranowski, E. Starzyńska, P. Brown and G.M. Blackburn	<b>257–262</b>
Relaxed enantioselectivity of human mitochondrial thymidine kinase and chemotherapeutic uses of L-nucleoside analogues	A. Verri, G. Priori, S. Spadari, L. Tondelli and F. Focher	<b>317–320</b>
<b>Carbohydrates and lipids</b>		
NMR solution conformation of heparin-derived hexasaccharide	D. Mikhailov, R.J. Linhardt and K.H. Mayo	<b>51–61</b>
Substrate specificity of the <i>N</i> -acetylglucosaminyl-phosphatidylinositol de-N-acetylase of glycosylphosphatidylinositol membrane anchor biosynthesis in African trypanosomes and human cells	D.K. Sharma, T.K. Smith, A. Crossman, J.S. Brimacombe and M.A.J. Ferguson	<b>171–177</b>
<b>Gene structure and expression</b>		
Degradation of the inducible cAMP early repressor (ICER) by the ubiquitin–proteasome pathway	E.J. Folco and G. Koren	<b>37–43</b>
Stress-induced transcription of the <i>clusterin/apoJ</i> gene	D. Michel, G. Chatelain, S. North and G. Brun	<b>45–50</b>
Regulation of human prohormone convertase 2 promoter activity by the transcription factor EGR-1	E. Jansen, T.A.Y. Ayoubi, S.M.P. Meulemans and W.J.M. Van de Ven	<b>69–74</b>
Regulation of human $\gamma$ -glutamylcysteine synthetase: co-ordinate induction of the catalytic and regulatory subunits in HepG2 cells	D.C. Galloway, D.G. Blake, A.G. Shepherd and L.I. McLellan	<b>99–104</b>
Species-specific alternative splicing generates a catalytically inactive form of human hormone-sensitive lipase	H. Laurell, J. Grober, C. Vindis, T. Lacombe, M. Dauzats, C. Holm and D. Langin	<b>137–143</b>
Expression of Menkes disease gene in mammary carcinoma cells	M.L. Ackland, E.J. Cornish, J.A. Paynter, A. Grimes, A. Michalczyk and J.F.B. Mercer	<b>237–243</b>
Thioredoxin from <i>Bacillus acidocaldarius</i> : characterization, high-level expression in <i>Escherichia coli</i> and molecular modelling	S. Bartolucci, A. Guagliardi, E. Pedone, D. De Pascale, R. Cannio, L. Camardella, M. Rossi, G. Nicastro, C. de Chiara, P. Facci, G. Mascetti and C. Nicolini	<b>277–285</b>
DNA-binding protein Pur $\alpha$ and transcription factor YY1 function as transcription activators of the neuron-specific FE65 gene promoter	N. Zambrano, S. De Renzis, G. Minopoli, R. Faraonio, V. Donini, A. Scaloni, F. Cimino and T. Russo	<b>293–300</b>
<b>Regulation of metabolism</b>		
A novel pathway for the conversion of homocysteine to methionine in eukaryotes	C.M. Antonio, M.C. Nunes, H. Refsum and A.K. Abraham	<b>165–170</b>
Characterization of homocysteine metabolism in the rat kidney	J.D. House, M.E. Brosnan and J.T. Brosnan	<b>287–292</b>
Rapid induction of apoptosis by deregulated uptake of polyamine analogues	R.-H. Hu and A.E. Pegg	<b>307–316</b>
<b>Membranes and bioenergetics</b>		
Short-term impairment of energy production in isolated rat liver mitochondria by hypoxia/reoxygenation: involvement of oxidative protein modification	L. Schild, T. Reinheckel, I. Wiswedel and W. Augustin	<b>205–210</b>
<b>Receptors and signal transduction</b>		
Protein kinase C-dependent and $\text{Ca}^{2+}$ -dependent mechanisms of secretion from streptolysin O-permeabilized platelets: effects of leakage of cytosolic proteins	D.C. Sloan and R.J. Haslam	<b>13–21</b>
Analysis of the N-terminal binding domain of $\text{G}_\alpha$	L. Busconi and B.M. Denker	<b>23–31</b>
Inositol 1,4,5-trisphosphate receptor subtypes differentially recognize regiosomers of D-myo-inositol 1,4,5-trisphosphate	M. Hirata, H. Takeuchi, A.M. Riley, S.J. Mills, Y. Watanabe and B.V.L. Potter	<b>93–98</b>
Role of G-protein $\beta\gamma$ subunits in the augmentation of $\text{P}_2\text{Y}_2$ ( $\text{P}_{2U}$ ) receptor-stimulated responses by neuropeptide Y $\text{Y}_1$ , $\text{G}_{i/o}$ -coupled receptors	L.A. Selbie, N.V. King, J.M. Dickenson and S.J. Hill	<b>153–158</b>
Interleukin-1-induced nuclear factor $\kappa\text{B}$ activation is inhibited by overexpression of phospholipid hydroperoxide glutathione peroxidase in a human endothelial cell line	R. Brigelius-Flohé, B. Friedrichs, S. Maurer, M. Schultz and R. Streicher	<b>199–203</b>
Characterization of $\alpha$ -conotoxin interactions with the nicotinic acetylcholine receptor and monoclonal antibodies	J.D. Ashcom and B.G. Stiles	<b>245–250</b>

---

Thrombin produces phosphorylation of cytosolic phospholipase A <sub>2</sub> by a mitogen-activated protein kinase kinase-independent mechanism in the human astrocytoma cell line 1321N1	M. Hernández, Y. Bayón, M. Sánchez Crespo and M.L. Nieto	263–269
The sphingomyelin–ceramide pathway participates in cytokine regulation of C-reactive protein and serum amyloid A, but not $\alpha$ -fibrinogen	G. Lozanski, F. Berthier and I. Kushner	271–275

**Cell biology and development**

Mammalian actin-related protein 2/3 complex localizes to regions of lamellipodial protrusion and is composed of evolutionarily conserved proteins	L.M. Machesky, E. Reeves, F. Wientjes, F.J. Mattheyse, A. Grogan, N.F. Totty, A.L. Burlingame, J.J. Hsuan and A.W. Segal	105–112
Adrenergic activation of vascular endothelial growth factor mRNA expression in rat brown adipose tissue: implication in cold-induced angiogenesis	A. Asano, M. Morimatsu, H. Nikami, T. Yoshida and M. Saito	179–183