Biochemical Journal Volume 330, part 1 15 February 1998

| Research Communication | | |
|--|---|---------|
| Novel evidence for an ecto-phospholipid methyltransferase in isolated rat hepatocytes | F. Bontemps and G. Van den Berghe | 1–4 |
| Research Papers | | |
| Proteins | | |
| Purification and characterization of N-glycanase, a concanavalin A binding protein from jackbean (Canavalia ensiformis) | P.S. Sheldon, J.N. Keen and D.J. Bowles | 13–20 |
| cDNA isolation of Alzheimer's amyloid precursor protein from cholinergic nerve terminals of the electric organ of the electric ray | Ki. lijima, DS. Lee, Ji. Okutsu, S. Tomita, N. Hirashima, Y. Kirino and T. Suzuki | 29–33 |
| Mutagenesis of the aspartic acid ligands in human serum transferrin: lobe-lobe interaction and conformation as revealed by antibody, receptor-binding and iron-release studies | A.B. Mason, QY. He, B.M. Tam, R.T.A. MacGillivray and R.C. Woodworth | 35–40 |
| Annexins I and II bind to lipid A: a possible role in the inhibition of endotoxins | D.A. Eberhard and S.R. Vandenberg | 67–72 |
| Purification and functional characterization of a low-molecular-mass Ca ²⁺ , Mg ²⁺ - and Ca ²⁺ -ATPase modulator protein from rat brain cytosol | D. Bhattacharyya and P.C. Sen | 95–101 |
| Presence of dopa and amino acid hydroperoxides in proteins modified with advanced glycation end products (AGEs): amino acid oxidation products as a possible source of oxidative stress induced by AGE proteins | S. Fu, MX. Fu, J.W. Baynes, S.R. Thorpe and R.T. Dean | 233–239 |
| Transient overexpression of human H- and L-ferritin chains in COS cells | B. Corsi, F. Perrone, M. Bourgeois, C. Beaumont, M.C. Panzeri, A. Cozzi, R. Sangregorio, P. Santambrogio, A. Albertini, P. Arosio and S. Levi | 315–320 |
| Cloning, expression and map assignment of chicken prosaposin | N. Azuma, HC. Seo, Ø. Lie, Q. Fu, R.M. Gould, M. Hiraiwa, D.W. Burt, I.R. Paton, D.R. Morrice, J.S. O'Brien and Y. Kishimoto | 321–327 |
| Ageing and zonal variation in post-translational modification of collagen in normal human articular cartilage. The age-related increase in non-enzymatic glycation affects biomechanical properties of cartilage | R.A. Bank, M.T. Bayliss, F.P.J.G. Lafeber, A. Maroudas and J.M. Tekoppele | 345–351 |
| Characterization of mini-protein S, a recombinant variant of protein S that lacks the sex hormone binding globulin-like domain | M. van Wijnen, J.G. Stam, G.T.G. Chang, J.C.M. Meijers, P.H. Reitsma, R.M. Bertina and B.N. Bouma | 389–396 |
| The small GTPase Gsp1p binds to the repeat domain of the nucleoporin Nsp1p | U. Stochaj, M. Héjazi and P. Belhumeur | 421–427 |
| Neutralizing epitope mapping of six β_1 -bungarotoxin monoclonal antibodies and its application in β_1 -bungarotoxin peptide vaccine design | C.C. Yang and H.L. Chan | 497–503 |
| Fe65L2: a new member of the Fe65 protein family interacting with the intracellular domain of the Alzheimer's β -amyloid precursor protein | A. Duilio, R. Faraonio, G. Minopoli, N. Zambrano and T. Russo | 513–519 |
| Enzymes | | |
| Substrate-binding domains of glycanases from <i>Streptomyces lividans</i> : characterization of a new family of xylan-binding domains | C. Dupont, M. Roberge, F. Shareck, R. Morosoli and D. Kluepfel | 41–45 |
| Recombinant pseudorabies virus DNase exhibits a RecBCD-like catalytic function | CY. Hsiang, TY. Ho, CH. Hsiang and TJ. Chang | 55–59 |

| A novel peptide-processing activity of insect peptidyl-dipeptidase A (angiotensin I-converting enzyme): the hydrolysis of lysyl-arginine and arginyl-arginine from the C-terminus of an insect prohormone peptide | R.E. Isaac, L. Schoofs, T.A. Williams, D. Veelaert, M. Sajid, P. Corvol and D. Coates | 61–65 |
|---|--|---------|
| Down-regulation by prostaglandins of type-II phospholipase A ₂ expression in guinea-pig alveolar macrophages: a possible involvement of cAMP | D. Vial, L. Arbibe, N. Havet, C. Dumarey, B.B. Vargaftig and L. Touqui | 89–94 |
| Chicken FK506-binding protein, FKBP65, a member of the FKBP family of peptidylprolyl <i>cis-trans</i> isomerases, is only partially inhibited by FK506 | B. Zeng, J.R. Macdonald, J.G. Bann, K. Beck, J.E. Gambee, B.A. Boswell and H.P. Bächinger | 109–114 |
| Effect of mutations in the tranmethylase and dehydrogenase/chelatase domains of sirohaem synthase (CysG) on sirohaem and cobalamin biosynthesis | S.C. Woodcock, E. Raux, F. Levillayer, C. Thermes, A. Rambach and M.J. Warren | 121–129 |
| Human glutathione transferase A4-4: an Alpha class enzyme with high catalytic efficiency in the conjugation of 4-hydroxynonenal and other genotoxic products of lipid peroxidation | I. Hubatsch, M. Ridderström and B. Mannervik | 175–179 |
| Mechanistic consequences of replacing the active-site nucleophile Glu-358 in $Agrobacterium$ sp. β -glucosidase with a cysteine residue | S.L. Lawson, R.A.J. Warren and S.G. Withers | 203–209 |
| Solvent kinetic isotope effects of human placental alkaline phosphatase in reverse micelles | TM. Huang, HC. Hung, TC. Chang and GG. Chang | 267–275 |
| Identification of the structural similarity in the functionally related amidohydrolases acting on the cyclic amide ring | GJ. Kim and HS. Kim | 295–302 |
| Redox- and anion-linked protonation sites in horseradish peroxidase: analysis of distal haem pocket mutants | B. Meunier, J.N. Rodriguez-Lopez, A.T. Smith, R.N.F. Thorneley and P.R. Rich | 303–309 |
| Chemical mechanism of p-amino acid oxidase from <i>Rhodotorula</i> gracilis: pH dependence of kinetic parameters | F. Ramón, M.P. Castillón, I. de la Mata and C. Acebal | 311–314 |
| Reconstitution of Cu ²⁺ -depleted bovine serum amine oxidase with Co ²⁺ | E. Agostinelli, G. de Matteis, B. Mondovi and L. Morpurgo | 383–387 |
| Activation of a phospholipase $C\beta 2$ deletion mutant by limited proteolysis | P. Schnabel and M. Camps | 461–468 |
| Substrate specificity of sheep liver sorbitol dehydrogenase | R.I. Lindstad, P. Köll and J.S. McKinley-McKee | 479–487 |
| Characterization of a cellobiose dehydrogenase from Humicola insolens | C. Schou, M.H. Christensen and M. Schülein | 565–571 |
| Carbohydrates and lipids | | |
| Purification, cDNA cloning and expression of 15-oxoprostaglandin 13-reductase from pig lung | C.M. Ensor, H. Zhang and HH. Tai | 103–108 |
| Regulation of expression of human intestinal bile acid-binding protein in Caco-2 cells | T. Kanda, L. Foucand, Y. Nakamura, I. Niot, P. Besnard, M. Fujita, Y. Sakai, K. Hatakeyama, T. Ono and H. Fujii | 261–265 |
| Structural analysis of a new series of oligosaccharide-alditols released by reductive β -elimination from oviducal mucins of Rana utricularia | W. Morelle and G. Strecker | 469–478 |
| Gene structure and expression | | |
| Ischaemia and reperfusion injury of rat liver increases expression of glutathione S-transferase A1/A2 in zone 3 of the hepatic lobule | G.D. Branum, N. Selim, X. Liu, R. Whalen and T.D. Boyer | 73–79 |
| DNA recognition by quinoxaline antibiotics: use of base- modified DNA molecules to investigate determinants of sequence- specific binding of triostin A and TANDEM | C. Bailly and M.J. Waring | 81–87 |
| Harpin and hydrogen peroxide both initiate programmed cell death but have differential effects on defence gene expression in <i>Arabidopsis</i> suspension cultures | R. Desikan, A. Reynolds, J.T. Hancock and S.J. Neill | 115–120 |
| Transcriptional regulation of MHC class I gene expression in rat oligodendrocytes | G. Mavria, K.T. Hall, R. A. Jones and G.E. Blair | 155–161 |
| Molecular cloning of a murine homologue of membrane cofactor protein (CD46): preferential expression in testicular germ cells | A. Tsujimura, K. Shida, M. Kitamura, M. Nomura, J. Takeda, H. Tanaka, M. Matsumoto, K. Matsumiya, A. Okuyama, Y. Nishimune, M. Okabe and T. Seya | 163–168 |

| Evidence that the E4 and FE4 esterase genes responsible for | | |
|---|---|---|
| insecticide resistance in the aphid <i>Myzus persicae</i> (Sulzer) are part of a gene family | L.M. Field and A.L. Devonshire | 169–173 |
| The nuclear factor interleukin-6 (NF-IL6) and signal transducer and activator of transcription-3 (STAT-3) signalling pathways co-operate to mediate the activation of the $hsp90\beta$ gene by interleukin-6 but have opposite effects on its inducibility by heat shock | A. Stephanou, D.A. Isenberg, S. Akira, T. Kishimoto and D.S. Latchman | 189–195 |
| Cloning and characterization of the promoter for the liver isoform of the rat carnitine palmitoyltransferase I (L-CPT I) gene | E.A. Park, M.L. Steffen, S. Song, V.M. Park and G.A. Cook | 217–224 |
| Functional analysis of the endothelial cell-specific Tie2/Tek promoter identifies unique protein-binding elements | B.M. Fadel, S.C. Boutet and T. Quertermous | 335–343 |
| Two distinct types of fatty acid-binding protein are expressed in heart ventricle of Antarctic teleost fishes | M.E. Vayda, R.L. Londraville, R.E. Cashon, L. Costello and B.D. Sidell | 375–382 |
| Spacing requirements for interactions between the C-terminal domain of the α subunit of <i>Escherichia coli</i> RNA polymerase and the cAMP receptor protein | G.S. Lloyd, S.J.W. Busby and N.J. Savery | 413–420 |
| Enhanced AP-1 and NF- κ B activities and stability of interleukin 8 (IL-8) transcripts are implicated in IL-8 mRNA superinduction in lung epithelial H292 cells | T. Roger, T.A. Out, N. Mukaida, K. Matsushima, H.M. Jansen and R. Lutter | 429–435 |
| Characterization of the rat salivary-gland B1-immunoreactive proteins | L. Mirels, A.J. Miranda and W.D. Ball | 437–444 |
| The molecular basis for UV vision in birds: spectral characteristics, cDNA sequence and retinal localization of the UV-sensitive visual pigment of the budgerigar (<i>Melopsittacus undulatus</i>) | S.E. Wilkie, P.M.A.M. Vissers, D. Das, W.J. DeGrip, J.K. Bowmaker and D.M. Hunt | 541–548 |
| Effects of nutrient deprivation and differentiation on the expression of growth-arrest genes (gas and gadd) in F9 embryonal carcinoma cells | J.V. Fleming, S.M. Hay, D.N. Harries and W.D. Rees | 573–579 |
| Regulation of metabolism | | |
| | | |
| Uroporphyria produced in mice by iron and 5-aminolaevulinic acid does not occur in $Cyp1a2(-I-)$ null mutant mice | P.R. Sinclair, N. Gorman, T. Dalton, H.S. Walton, W.J. Bement, J.F. Sinclair, A.G. Smith and D.W. Nebert | 149–153 |
| | | 149–153 181–187 |
| acid does not occur in $Cyp1a2(-I-)$ null mutant mice | W.J. Bement, J.F. Sinclair, A.G. Smith and D.W. Nebert | |
| acid does not occur in $Cyp1a2(-I-)$ null mutant mice In vitro analysis of UV-damage-induced inhibition of replication Tyrosine kinase-dependent modulation of 3-hydroxy-3-methyl-glutaryl-CoA reductase in human breast adenocarcinoma | W.J. Bement, J.F. Sinclair, A.G. Smith and D.W. Nebert R. Driss and SH. Lee | 181–187 |
| acid does not occur in $Cyp1a2(-I-)$ null mutant mice In vitro analysis of UV-damage-induced inhibition of replication Tyrosine kinase-dependent modulation of 3-hydroxy-3-methylglutaryl-CoA reductase in human breast adenocarcinoma SKBR-3 cells Signalling pathways of an insulin-mimetic | W.J. Bement, J.F. Sinclair, A.G. Smith and D.W. Nebert R. Driss and SH. Lee R. Asslan, A. Pradines, G. Favre and F. Le Gaillard | 181–187 241–246 |
| acid does not occur in <i>Cyp1a2(-I-)</i> null mutant mice <i>In vitro</i> analysis of UV-damage-induced inhibition of replication Tyrosine kinase-dependent modulation of 3-hydroxy-3-methylglutaryl-CoA reductase in human breast adenocarcinoma SKBR-3 cells Signalling pathways of an insulin-mimetic phosphoinositolglycan-peptide in muscle and adipose tissue Effects of noradrenaline on the cell-surface glucose transporters in cultured brown adipocytes: novel mechanism for selective | W.J. Bement, J.F. Sinclair, A.G. Smith and D.W. Nebert R. Driss and SH. Lee R. Asslan, A. Pradines, G. Favre and F. Le Gaillard A. Kessler, G. Müller, S. Wied, A. Crecelius and J. Eckel Y. Shimizu, S. Satoh, H. Yano, Y. Minokoshi, | 181–187 241–246 277–286 |
| acid does not occur in <i>Cyp1a2(-I-)</i> null mutant mice <i>In vitro</i> analysis of UV-damage-induced inhibition of replication Tyrosine kinase-dependent modulation of 3-hydroxy-3-methylglutaryl-CoA reductase in human breast adenocarcinoma SKBR-3 cells Signalling pathways of an insulin-mimetic phosphoinositolglycan-peptide in muscle and adipose tissue Effects of noradrenaline on the cell-surface glucose transporters in cultured brown adipocytes: novel mechanism for selective activation of GLUT1 glucose transporters Possible role for gp160 in constitutive but not insulin-stimulated | W.J. Bement, J.F. Sinclair, A.G. Smith and D.W. Nebert R. Driss and SH. Lee R. Asslan, A. Pradines, G. Favre and F. Le Gaillard A. Kessler, G. Müller, S. Wied, A. Crecelius and J. Eckel Y. Shimizu, S. Satoh, H. Yano, Y. Minokoshi, S.W. Cushman and T. Shimazu | 181–187 241–246 277–286 397–403 |
| acid does not occur in <i>Cyp1a2(-I-)</i> null mutant mice <i>In vitro</i> analysis of UV-damage-induced inhibition of replication Tyrosine kinase-dependent modulation of 3-hydroxy-3-methylglutaryl-CoA reductase in human breast adenocarcinoma SKBR-3 cells Signalling pathways of an insulin-mimetic phosphoinositolglycan-peptide in muscle and adipose tissue Effects of noradrenaline on the cell-surface glucose transporters in cultured brown adipocytes: novel mechanism for selective activation of GLUT1 glucose transporters Possible role for gp160 in constitutive but not insulin-stimulated GLUT4 trafficking: dissociation of gp160 and GLUT4 localization Effect of 3,5-di-iodo-L-thyronine on the mitochondrial | W.J. Bement, J.F. Sinclair, A.G. Smith and D.W. Nebert R. Driss and SH. Lee R. Asslan, A. Pradines, G. Favre and F. Le Gaillard A. Kessler, G. Müller, S. Wied, A. Crecelius and J. Eckel Y. Shimizu, S. Satoh, H. Yano, Y. Minokoshi, S.W. Cushman and T. Shimazu A. Filippis, S. Clark and J. Proietto A. Lombardi, A. Lanni, M. Moreno, M.D. Brand and | 181–187 241–246 277–286 397–403 405–411 |
| acid does not occur in <i>Cyp1a2(-I-)</i> null mutant mice <i>In vitro</i> analysis of UV-damage-induced inhibition of replication Tyrosine kinase-dependent modulation of 3-hydroxy-3-methylglutaryl-CoA reductase in human breast adenocarcinoma SKBR-3 cells Signalling pathways of an insulin-mimetic phosphoinositolglycan-peptide in muscle and adipose tissue Effects of noradrenaline on the cell-surface glucose transporters in cultured brown adipocytes: novel mechanism for selective activation of GLUT1 glucose transporters Possible role for gp160 in constitutive but not insulin-stimulated GLUT4 trafficking: dissociation of gp160 and GLUT4 localization Effect of 3,5-di-iodo-L-thyronine on the mitochondrial energy-transduction apparatus | W.J. Bement, J.F. Sinclair, A.G. Smith and D.W. Nebert R. Driss and SH. Lee R. Asslan, A. Pradines, G. Favre and F. Le Gaillard A. Kessler, G. Müller, S. Wied, A. Crecelius and J. Eckel Y. Shimizu, S. Satoh, H. Yano, Y. Minokoshi, S.W. Cushman and T. Shimazu A. Filippis, S. Clark and J. Proietto A. Lombardi, A. Lanni, M. Moreno, M.D. Brand and F. Goglia | 181–187 241–246 277–286 397–403 405–411 521–526 |
| acid does not occur in <i>Cyp1a2(-I-)</i> null mutant mice <i>In vitro</i> analysis of UV-damage-induced inhibition of replication Tyrosine kinase-dependent modulation of 3-hydroxy-3-methylglutaryl-CoA reductase in human breast adenocarcinoma SKBR-3 cells Signalling pathways of an insulin-mimetic phosphoinositolglycan-peptide in muscle and adipose tissue Effects of noradrenaline on the cell-surface glucose transporters in cultured brown adipocytes: novel mechanism for selective activation of GLUT1 glucose transporters Possible role for gp160 in constitutive but not insulin-stimulated GLUT4 trafficking: dissociation of gp160 and GLUT4 localization Effect of 3,5-di-iodo-L-thyronine on the mitochondrial energy-transduction apparatus Estimation of total rate of formation of nitric oxide in the rat | W.J. Bement, J.F. Sinclair, A.G. Smith and D.W. Nebert R. Driss and SH. Lee R. Asslan, A. Pradines, G. Favre and F. Le Gaillard A. Kessler, G. Müller, S. Wied, A. Crecelius and J. Eckel Y. Shimizu, S. Satoh, H. Yano, Y. Minokoshi, S.W. Cushman and T. Shimazu A. Filippis, S. Clark and J. Proietto A. Lombardi, A. Lanni, M. Moreno, M.D. Brand and F. Goglia | 181–187 241–246 277–286 397–403 405–411 521–526 |
| acid does not occur in <i>Cyp1a2(-I-)</i> null mutant mice <i>In vitro</i> analysis of UV-damage-induced inhibition of replication Tyrosine kinase-dependent modulation of 3-hydroxy-3-methylglutaryl-CoA reductase in human breast adenocarcinoma SKBR-3 cells Signalling pathways of an insulin-mimetic phosphoinositolglycan-peptide in muscle and adipose tissue Effects of noradrenaline on the cell-surface glucose transporters in cultured brown adipocytes: novel mechanism for selective activation of GLUT1 glucose transporters Possible role for gp160 in constitutive but not insulin-stimulated GLUT4 trafficking: dissociation of gp160 and GLUT4 localization Effect of 3,5-di-iodo-L-thyronine on the mitochondrial energy-transduction apparatus Estimation of total rate of formation of nitric oxide in the rat Membranes and bioenergetics Binding of MARCKS (myristoylated alanine-rich C kinase substrate)-related protein (MRP) to vesicular phospholipid | W.J. Bement, J.F. Sinclair, A.G. Smith and D.W. Nebert R. Driss and SH. Lee R. Asslan, A. Pradines, G. Favre and F. Le Gaillard A. Kessler, G. Müller, S. Wied, A. Crecelius and J. Eckel Y. Shimizu, S. Satoh, H. Yano, Y. Minokoshi, S.W. Cushman and T. Shimazu A. Filippis, S. Clark and J. Proietto A. Lombardi, A. Lanni, M. Moreno, M.D. Brand and F. Goglia A. Sakinis and Å. Wennmalm | 181–187 241–246 277–286 397–403 405–411 521–526 527–532 |

| Effective electrostatic charge of coagulation factor X in solution and on phospholipid membranes: implications for activation mechanisms and structure—function relationships of the Gla domain | M.P. McGee, H. Teuschler and J. Liang | 533–539 |
|--|--|---------|
| Receptors and signal transduction | | |
| Angiotensin II potentiates adrenocorticotrophic hormone-induced cAMP formation in bovine adrenal glomerulosa cells through a capacitative calcium influx | M.M. Burnay, M.B. Vallotton, A.M. Capponi and M.F. Rossier | 21–27 |
| Internalization of the interleukin 6 signal transducer gp130 does not require activation of the Jak/STAT pathway | S. Thiel, I. Behrmann, E. Dittrich, L. Muys, J. Tavernier, J. Wijdenes, P.C. Heinrich and L. Graeve | 47–54 |
| Oscillating response to a purine nucleotide disrupted by mutation in <i>Paramecium tetraurelia</i> | J.L. Mimikakis, D. L. Nelson and R.R. Preston | 139–147 |
| Different pathways of colony-stimulating factor 1 degradation in macrophage populations revealed by wortmannin sensitivity | V. Kanagasundaram, E. Christy, J.A. Hamilton and A. Jaworowski | 197–202 |
| Dephosphorylation of Ser-137 in DARPP-32 by protein phosphatases 2A and 2C: different roles <i>in vitro</i> and in striatonigral neurons | F. Desdouits, J.C. Siciliano, A.C. Nairn, P. Greengard and JA. Girault | 211–216 |
| Identification and localization of myosin phosphatase in human platelets | A. Murányi, F. Erdődi, M. Ito, P. Gergely and D.J. Hartshorne | 225–231 |
| C1q-mediated chemotaxis by human neutrophils: involvement of gClqR and G-protein signalling mechanisms | L.E.A. Leigh, B. Ghebrehiwet, T.P.S. Perera, I.N. Bird, P. Strong, U. Kishore, K.B.M. Reid and P. Eggleton | 247–254 |
| Functional rescue of a constitutively desensitized $\beta_2 \mathrm{AR}$ through receptor dimerization | T.E. Hebert, T.P. Loisel, L. Adam, N. Ethier, S. St. Onge and M. Bouvier | 287–293 |
| Cholecystokinin octapeptide inhibits Ca ²⁺ -dependent amylase secretion from permeabilized pancreatic acini by blocking the MgATP-dependent priming of exocytosis | P.J. Padfield and N. Panesar | 329–334 |
| Nucleotide-binding properties of kinase-deficient epidermal-growth-factor-receptor mutants | K. Cheng and J.G. Koland | 353–359 |
| Roles for interleukin-1 β , phorbol ester and a post-transcriptional regulator in the control of bradykinin B1 receptor gene expression | X. Zhou, P. Polgar and L. Taylor | 361–366 |
| Sustained phospholipase D activation in response to angiotensin II but not carbachol in bovine adrenal glomerulosa cells | E. Jung, S. Betancourt-Calle, R. Mann-Blakeney, T. Foushee, C.M. Isales and W.B. Bollag | 445–451 |
| Rapid activation of sodium-proton exchange and extracellular signal-regulated protein kinase in fibroblasts by G protein-coupled 5-HT _{1A} receptor involves distinct signalling cascades | M.N. Garnovskaya, Y. Mukhin and J.R. Raymond | 489–495 |
| The effects of free [Ca ²⁺] on the cytosolic face of the inositol 1,4,5-trisphosphate receptor at the single channel level | E.C. Thrower, E.J.A. Lea and A.P. Dawson | 559–564 |
| Cell biology and development | | |
| Comparison of human red cell lysis by hypochlorous and hypobromous acids: insights into the mechanism of lysis | M.C.M. Vissers, A.C. Carr and A.L.P. Chapman | 131–138 |
| Biosynthesis of endotubin: an apical early endosomal glycoprotein from developing rat intestinal epithelial cells | K. Allen, K.E. Gokay, M.A. Thomas, B.A. Speelman and J.M. Wilson | 367–373 |
| Infectivity of <i>Trypanosoma cruzi</i> strains is associated with differential expression of surface glycoproteins with differential Ca ²⁺ signalling activity | R.C. Ruiz, S. Favoreto Jr., M.L. Dorta, M.E.M. Oshiro, A.T. Ferreira, P.M. Manque and N. Yoshida | 505–511 |
| Chondroadherin expression changes in skeletal development | Z. Shen, S. Gantcheva, B. Månsson, D. Heinegård and Y. Sommarin | 549–557 |