

Review

- The mitochondrial permeability transition pore and its role in cell death** 233–249
by M. Crompton

Research Communication

- The mechanism of activation of NADPH oxidase in the cell-free system: the activation process is primarily catalytic and not through the formation of a stoichiometric complex A.R. Cross, R.W. Erickson and J.T. Curnutte 251–255

Proteins

- Osteocalcin binds tightly to the γ -glutamylcarboxylase at a site distinct from that of the other known vitamin K-dependent proteins R.J.T.J. Houben, D. Jin, D.W. Stafford, P. Proost, R.H.M. Ebberink, C. Vermeer and B.A.M. Soute 265–269
- Identification of pituitary adenylate cyclase-activating polypeptide 1-38-binding factor in human plasma, as ceruloplasmin J.W. Tams, A.H. Johnsen and J. Fahrenkrug 271–276
- Cloning and characterization of gp36, a human mucin-type glycoprotein preferentially expressed in vascular endothelium G. Zimmer, F. Oeffner, V. von Messling, T. Tschernig, H.-J. Gröne, H.-D. Klenk and G. Herrler 277–284
- Expression, purification and biochemical characterization of recombinant murine secretory component: a novel tool in mucosal immunology P. Cottet, S. Cottet and B. Corthésy 299–306
- Molecular analysis of sialoside binding to sialoadhesin by NMR and site-directed mutagenesis P.R. Crocker, M. Vinson, S. Kelm and K. Drickamer 355–361

Enzymes

- Purification and enzymic properties of the fructosyltransferase of *Streptococcus salivarius* ATCC 25975 D.D. Song and N.A. Jacques 285–291
- Reductive half-reaction of the H172Q mutant of trimethylamine dehydrogenase: evidence against a carbanion mechanism and assignment of kinetically influential ionizations in the enzyme–substrate complex J. Basran, M.J. Sutcliffe, R. Hille and N.S. Scrutton 307–314
- Multiple-site phosphorylation of the 280 kDa isoform of acetyl-CoA carboxylase in rat cardiac myocytes: evidence that cAMP-dependent protein kinase mediates effects of β -adrenergic stimulation A.N. Boone, B. Rodrigues and R.W. Brownsey 347–354
- Characterization of trehalose phosphorylase from *Schizophyllum commune* C. Eis and B. Nidetzky 385–393
- Histidine-193 of rat glucosylceramide synthase resides in a UDP-glucose- and inhibitor (*D*-threo-1-phenyl-2-decanoylamino-3-morpholinopropan-1-ol)-binding region: a biochemical and mutational study K. Wu, D.L. Marks, R. Watanabe, P. Paul, N. Rajan and R.E. Pagano 395–400
- Peptidase activity of β -lactamases N. Rhazi, M. Galleni, M.I. Page and J.-M. Frère 409–413

Gene structure and expression

- Upstream stimulatory factor regulates *Pdx-1* gene expression in differentiated pancreatic β -cells J. Qian, E.N. Kaytor, H.C. Towle and L.K. Olson 315–322

p44/42 mitogen-activated protein kinase is involved in the expression of ornithine decarboxylase in leukaemia L1210 cells	F. Flamigni, A. Facchini, C. Capanni, C. Stefanelli, B. Tantini and C.M. Calderera	363–369
Polyunsaturated fatty acids inhibit fatty acid synthase and spot-14-protein gene expression in cultured rat hepatocytes by a peroxidative mechanism	M. Foretz, F. Foufelle and P. Ferré	371–376
Goblet-cell-specific transcription of mouse intestinal trefoil factor gene results from collaboration of complex series of positive and negative regulatory elements	H. Itoh, N. Inoue and D.K. Podolsky	461–472
Regulation of metabolism		
Lipoprotein cholesterol uptake mediates up-regulation of bile-acid synthesis by increasing cholesterol 7 α -hydroxylase but not sterol 27-hydroxylase gene expression in cultured rat hepatocytes	S.M. Post, J. Twisk, L. van der Fits, E.C.M. de Wit, M.F.M. Hoekman, W.H. Mager and H.M.G. Princen	339–346
Metabolic characteristics of a human hepatoma cell line stably transfected with hormone-sensitive lipase	R.J. Pease, D. Wiggins, E.D. Saggerson, J. Tree and G.F. Gibbons	453–460
Membranes and bioenergetics		
A shift in the equilibrium constant at the catalytic site of proton-translocating transhydrogenase: significance for a 'binding-change' mechanism	J.D. Venning and J.B. Jackson	329–337
Receptors and signal transduction		
Expression of I ₂ ^{PP2A} , an inhibitor of protein phosphatase 2A, induces c-Jun and AP-1 activity	S.W.K. Al-Murrani, J.R. Woodgett and Z. Damuni	293–298
39-kDa receptor-associated protein (RAP) facilitates secretion and ligand binding of extracellular region of very-low-density-lipoprotein receptor: implications for a distinct pathway from low-density-lipoprotein receptor	A. Sato, Y. Shimada, J. Herz, T. Yamamoto and H. Jingami	377–383
Evidence for the involvement of a small subregion of the endoplasmic reticulum in the inositol trisphosphate receptor-induced activation of Ca ²⁺ inflow in rat hepatocytes	R.B. Gregory, R.A. Wilcox, L.A. Berven, N.C.R. van Straten, G.A. van der Mare, J.H. van Boom and G.J. Barritt	401–408
Rapid internalization and surface expression of a functional, fluorescently tagged G-protein-coupled glutamate receptor	A.J. Doherty, V. Coutinho, G.L. Collingridge and J.M. Henley	415–422
Calexitin interaction with neuronal ryanodine receptors	T.J. Nelson, W.-Q. Zhao, S. Yuan, A. Favit, L. Pozzo-Miller and D.L. Alkon	423–433
Resynthesis of phosphatidylinositol in permeabilized neutrophils following phospholipase C β activation: transport of the intermediate, phosphatidic acid, from the plasma membrane to the endoplasmic reticulum for phosphatidylinositol resynthesis is not dependent on soluble lipid carriers or vesicular transport	J. Whatmore, C. Wiedemann, P. Somerharju, P. Swigart and S. Cockcroft	435–444
Cell biology and development		
Talin contains three similar vinculin-binding sites predicted to form an amphipathic helix	M.D. Bass, B.J. Smith, S.A. Prigent and D.R. Critchley	257–263
Cell polarization is required for ricin sensitivity in a Caco-2 cell line selected for ricin resistance	M.R. Jackman, J.A. Ellis, S.R. Gray, W. Shurety and J.P. Luzio	323–327
Endocytic delivery of intramolecularly quenched substrates and inhibitors to the intracellular yeast Kex2 protease	M.K. Henkel, G. Pott, A.W. Henkel, L. Juliano, C.-M. Kam, J.C. Powers and A. Franzusoff	445–452

