Biochemical Journal Volume 342, part 3 15 September 1999

First published on the Internet 8 September 1999

Review		
Repression of gene expression by oxidative stress <i>by</i> Y. Morel and R. Barouki		481–496
Research Communications		
High-efficiency Semliki Forest virus-mediated transduction in bovine adrenal chromaffin cells	R.R. Duncan, A.C. Don-Wauchope, S. Tapechum, M.J. Shipston, R.H. Chow and P. Estibeiro	497–501
Salicylate-enhanced activation of transcription factors induced by interferon- γ	LC. Chen, D. Kepka-Lenhart, T.M. Wright and S.M. Morris, Jr.	503–507
Research Papers		
Proteins		
Characterization and polyanion-binding properties of purified recombinant prion protein	D.B. Brimacombe, A.D. Bennett, F.S. Wusteman, A.C. Gill, J.C. Dann and C.J. Bostock	605–613
Production and functional activity of a recombinant von Willebrand factor-A domain from human complement factor B	S.C. Williams, J. Hinshelwood, S.J. Perkins and R.B. Sim	625–632
Iron-dependent regulation of transferrin receptor expression in <i>Trypanosoma brucei</i>	B. Fast, K. Kremp, M. Boshart and D. Steverding	691–696
Enzymes		
Ornithine and glutamate decarboxylases catalyse an oxidative deamination of their α -methyl substrates	M. Bertoldi, V. Carbone and C. Borri Voltattorni	509–512
Characterization of prenylated protein methyltransferase in <i>Leishmania</i>	MP. Hasne and F. Lawrence	513–518
GA2/GM2/GD2 synthase localizes to the trans-Golgi network of CHO-K1 cells	C.G. Giraudo, V.M. Rosales Fritz and H.J.F. Maccioni	633–640
Interaction with amylopectin influences the ability of granule-bound starch synthase I to elongate malto-oligosaccharides	K. Denyer, D. Waite, A. Edwards, C. Martin and A.M. Smith	647–653
Caspase-1 (interleukin-1 β -converting enzyme) is inhibited by the human serpin analogue proteinase inhibitor 9	R.R. Annand, J.R. Dahlen, C.A. Sprecher, P. de Dreu, D.C. Foster, J.A. Mankovich, R.V. Talanian, W. Kisiel and D.A. Giegel	655–665
Editing of non-cognate aminoacyl adenylates by peptide synthetases	M. Pavela-Vrancic, R. Dieckmann, H. von Döhren and H. Kleinkauf	715–719
Cloning and sequencing of the cDNA species for mammalian dimeric dihydrodiol dehydrogenases	E. Arimitsu, S. Aoki, S. Ishikura, K. Nakanishi, K. Matsuura and A. Hara	721–728
Carbohydrates and lipids		
Structural determination of a 5-acetamido-3,5,7,9-tetradeoxy-7-(3-hydroxybutyramido)-L-manno-nonulosonic acid-containing homopolysaccharide isolated from <i>Sinorhizobium fredii</i> HH103	A.M. Gil-Serrano, M.A. Rodriguez-Carvajal, P. Tejero- Mateo, J.L. Espartero, M. Menendez, J. Corzo, J.E. Ruiz-Sainz and A.M. Buendía-Clavería	527–535

The dihydrosphingosine-1-phosphate phosphatases of <i>Saccharomyces cerevisiae</i> are important regulators of cell proliferation and heat stress responses	C. Mao, J.D. Saba and L.M. Obeid	667–675
Gene structure and expression		
Expression of translationally controlled tumour protein is regulated by calcium at both the transcriptional and post-transcriptional level	A. Xu, A.R. Bellamy and J.A. Taylor	683–689
Regulation of metabolism		
Inhibition of glycogenolysis in primary rat hepatocytes by 1,4-dideoxy-1,4-imino-p-arabinitol	B. Andersen, A. Rassov, N. Westergaard and K. Lundgren	545–550
Acetate stimulates flux through the tricarboxylic acid cycle in rabbit renal proximal tubules synthesizing glutamine from alanine: a ¹³ C NMR study	S. Dugelay, MF. Chauvin, F. Megnin-Chanet, G. Martin, MC. Laréal, JM. Lhoste and G. Baverel	555–566
Model of 2,3-bisphosphoglycerate metabolism in the human erythrocyte based on detailed enzyme kinetic equations: <i>in vivo</i> kinetic characterization of 2,3-bisphosphoglycerate synthase/phosphatase using ¹³ C and ³¹ P NMR	P.J. Mulquiney, W.A. Bubb and P.W. Kuchel	567–580
Model of 2,3-bisphosphoglycerate metabolism in the human erythrocyte based on detailed enzyme kinetic equations: equations and parameter refinement	P.J. Mulquiney and P.W. Kuchel	581–596
Model of 2,3-bisphosphoglycerate metabolism in the human erythrocyte based on detailed enzyme kinetic equations: computer simulation and Metabolic Control Analysis	P.J. Mulquiney and P.W. Kuchel	597–604
Membranes and bioenergetics		
Defective kinetics of cytochrome c oxidase and alteration of mitochondrial membrane potential in fibroblasts and cytoplasmic hybrid cells with the mutation for myoclonus epilepsy with ragged-red fibres ('MERFF') at position 8344 nt	H. Antonická, D. Floryk, P. Klement, L. Stratilová, J. Heřmanská, H. Houšťková, M. Kalous, Z. Drahota, J. Zeman and J. Houštěk	537–544
Effects of disrupting the 21 kDa subunit of complex I from Neurospora crassa	F. Ferreirinha, M. Duarte, A.M.P. Melo and A. Videira	551–554
Localization of a carboxylic residue possibly involved in the inhibition of vacuolar H ⁺ -pyrophosphatase by <i>N,N'</i> -dicyclohexylcarbodi-imide	S.J. Yang, S.S. Jiang, S.Y. Kuo, S.H. Hung, M.F. Tam and R.L. Pan	641–646
Receptors and signal transduction		
Cellular stress in <i>Xenopus</i> kidney cells enhances the phosphorylation of eukaryotic translation initiation factor (eIF)4E and the association of eIF4F with poly(A)-binding protein	C.S. Fraser, V.M. Pain and S.J. Morley	519–526
Signal transduction by $\beta 1$ integrin receptors in human chondrocytes <i>in vitro</i> : collaboration with the insulin-like growth factor-1 receptor	M. Shakibaei, T. John, P. de Souza, R. Rahmanzadeh and HJ. Merker	615–623
Differential effects of fibroblast growth factor (FGF) 9 and FGF2 on proliferation, differentiation and terminal differentiation of chondrocytic cells <i>in vitro</i>	N.B. Weksler, G.P. Lunstrum, E.S. Reid and W.A. Horton	677–682
Distribution of the Src-homology-2-domain- containing inositol 5-phosphatase SHIP-2 in both non-haemopoietic and haemopoietic cells and possible involvement of SHIP-2 in negative signalling of B-cells	E. Muraille, X. Pesesse, C. Kuntz and C. Erneux	697–705

		Contents	xiii
Calcium-dependent properties of CIB binding to the integrin αIIb cytoplasmic domain and translocation to the platelet cytoskeleton	D.D. Shock, U.P. Naik, J.E. Brittain, S.K. Alahari, J. Sondek and L.V. Parise		729–735
Cell biology and development			
nSec-1 (munc-18) interacts with both primed and unprimed syntaxin 1A and associated in a dimeric complex on adrenal chromaffin granules	L.P. Haynes, A. Morgan and R.D. Burgoyne		707–714