

**Review****Phospholipase D: molecular and cell biology of a novel gene family**

by M. Liscovitch, M. Czarny, G. Fiucci and X. Tang

401–415

**Research Communication**

Cytoplasmic STAT proteins associate prior to activation

S. Haan, M. Kortylewski, I. Behrmann, W. Müller-Esterl, P.C. Heinrich and F. Schaper

417–421

**Research Papers****Proteins**

Analysis of free and protein-bound nitrotyrosine in human plasma by a gas chromatography/mass spectrometry method that avoids nitration artifacts

M.T. Frost, B. Halliwell and K.P. Moore

453–458

 $\alpha$ -Crystallin assists the renaturation of glyceraldehyde-3-phosphate dehydrogenase

E. Ganea and J.J. Harding

467–472

Collagen fragments in urine derived from bone resorption are highly racemized and isomerized: a biological clock of protein aging with clinical potential

P.A.C. Cloos and C. Fledelius

473–480

Racemization and isomerization of type I collagen C-telopeptides in human bone and soft tissues: assessment of tissue turnover

E. Gineyts, P.A.C. Cloos, O. Borel, L. Grimaud, P.D. Delmas and P. Garnero

481–485

L-Pipecolic acid oxidase, a human enzyme essential for the degradation of L-pipecolic acid, is most similar to the monomeric sarcosine oxidases

G. Dodt, D.G. Kim, S.A. Reimann, B.E. Reuber, K. McCabe, S.J. Gould and S.J. Mihalik

487–494

The recombinant N-terminal region of human salivary mucin MG2 (MUC7) contains a binding domain for oral *Streptococci* and exhibits candidacidal activity

B. Liu, S.A. Rayment, C. Gyurko, F.G. Oppenheim, G.D. Offner and R.F. Troxler

557–564

Disulphide-bond pattern and molecular modelling of the dimeric disintegrin EMF-10, a potent and selective integrin  $\alpha_5\beta_1$  antagonist from *Eristocophis macmahoni* venom

J.J. Calvete, M. Jürgens, C. Marcinkiewicz, A. Romero, M. Schrader and S. Niewiarowski

573–581

Heat shock cognate protein 70 chaperone-binding site in the co-chaperone murine stress-inducible protein 1 maps to within three consecutive tetratricopeptide repeat motifs

J. van der Spuy, B.D. Kana, H.W. Dirr and G.L. Blatch

645–651

LKB1, a novel serine/threonine protein kinase and potential tumour suppressor, is phosphorylated by cAMP-dependent protein kinase (PKA) and prenylated *in vivo*

S.P. Collins, J.L. Reoma, D.M. Gamm and M.D. Uhler

673–680

Isolation and characterization of glycosylphosphatidylinositol-anchored, mucin-like surface glycoproteins from bloodstream forms of the freshwater-fish parasite *Trypanosoma carassii*

A. Lischke, C. Klein, Y.-D. Stierhof, M. Hempel, A. Mehlert, I.C. Almeida, M.A.J. Ferguson and P. Overath

693–700

Differential recognition of snake venom proteins expressing specific Arg-Gly-Asp (RGD) sequence motifs by wild-type and variant integrin  $\alpha_{11b}\beta_3$ : further evidence for distinct sites of RGD ligand recognition exhibiting negative allosterity

S. Rahman, G. Flynn, A. Aitken, Y. Patel, F. Hussain, X. Lu, J.C. Loftus, D. French, E. Wijelath, K. Strand and G.F. Savidge

701–709

A possible origin of differences between calorimetric and equilibrium estimates of stability parameters of proteins	A. Sinha, S. Yadav, R. Ahmad and F. Ahmad	711–717
Direct and reversed amino acid sequence pattern analysis: structural reasons for activity of reversed sequence sites and results of kinase site mutagenesis	I. Torshin	733–740
<b>Enzymes</b>		
Preliminary evidence for the existence of specific functional assemblies between enzymes of the $\beta$ -oxidation pathway and the respiratory chain	A. Parker and P.C. Engel	429–435
The regulation of AMP-activated protein kinase by phosphorylation	S.C. Stein, A. Woods, N.A. Jones, M.D. Davison and D. Carling	437–443
Analysis of tissue inhibitor of metalloproteinases-2 effect on pro-matrix metalloproteinase-2 activation by membrane-type 1 matrix metalloproteinase using baculovirus/insect-cell expression system	Y. Jo, J. Yeon, H.-J. Kim and S.-T. Lee	511–519
Evidence that serine 304 is not a key ligand-binding residue in the active site of cytochrome P450 2D6	S.W. Ellis, G.P. Hayhurst, T. Lightfoot, G. Smith, J. Harlow, K. Rowland-Yeo, C. Larsson, J. Mahling, C.K. Lim, C.R. Wolf, M.G. Blackburn, M.S. Lennard and G.T. Tucker	565–571
Purification and characterization of a novel human acidic nuclease/intracyclobutyl-pyrimidine-dimer-DNA phosphodiesterase	K.S. Famulski, M. Liuzzi, S. Bashir, R. Mirzayans and M.C. Paterson	583–593
Leukotriene A <sub>4</sub> hydrolase: a critical role of glutamic acid-296 for the binding of bestatin	M. Andberg, A. Wetterholm, J.F. Medina and J.Z. Haeggström	621–625
Mechanistic studies of morphine dehydrogenase and stabilization against covalent inactivation	E.H. Walker, C.E. French, D.A. Rathbone and N.C. Bruce	687–692
<b>Carbohydrates and lipids</b>		
Evidence that the inhibition of cartilage proteoglycan breakdown by mannosamine is not mediated via inhibition of glycosylphosphatidylinositol anchor formation	H. Bryson, D.J. Buttle, L.D. Kozaci, R.N. Johnatty and R.A.D. Bunning	495–501
<b>Gene structure and expression</b>		
Mitochondrial aspartate aminotransferase: direction of a single protein with two distinct functions to two subcellular sites does not require alternative splicing of the mRNA	M.W. Bradbury and P.D. Berk	423–427
Four isoforms of serum response factor that increase or inhibit smooth-muscle-specific promoter activity	P.R. Kemp and J.C. Metcalfe	445–451
The gene for human TATA-binding-protein-associated factor (TAF <sub>II</sub> ) 170: structure, promoter and chromosomal localization	J.A. van der Knaap, V. van den Boom, J. Kuipers, M.J.T. van Eijk, P.C. van der Vliet and H.Th.M. Timmers	521–527
Mouse BTEB3, a new member of the basic transcription element binding protein (BTEB) family, activates expression from GC-rich minimal promoter regions	K.M. Martin, W.N. Cooper, J.C. Metcalfe and P.R. Kemp	529–533
Induction of Myc-intron-binding polypeptides MIBP1 and RFX1 during retinoic acid-mediated differentiation of haemopoietic cells	M. Zajac-Kaye, N. Ben-Baruch, E. Kastanos, F.J. Kaye and C. Allegra	535–541
A bone sialoprotein-binding protein from <i>Staphylococcus aureus</i> : a member of the staphylococcal Sdr family	H.-s. Tung, B. Guss, U. Hellman, L. Persson, K. Rubin and C. Rydén	611–619
<i>pgaA</i> and <i>pgaB</i> encode two constitutively expressed endopolygalacturonases of <i>Aspergillus niger</i>	L. Pařenicová, J.A.E. Benen, H.C.M. Kester and J. Visser	637–644
Characterization and gene structure of a novel retinoblastoma-protein-associated protein similar to the transcription regulator TFII-I	X. Yan, X. Zhao, M. Qian, N. Guo, X. Gong and X. Zhu	749–757

**Regulation of metabolism**

- Lipid metabolic changes caused by short-chain ceramides and the connection with apoptosis  
D. Allan **603–610**
- Dehydroascorbic acid uptake in a human keratinocyte cell line (HaCaT) is glutathione-independent  
I. Savini, S. Duflot and L. Avigliano **665–672**

**Membranes and bioenergetics**

- Human Nedd4 interacts with the human epithelial Na<sup>+</sup> channel: WW3 but not WW1 binds to Na<sup>+</sup>-channel subunits  
T.J. Farr, S.J. Coddington-Lawson, P.M. Snyder and F.J. McDonald **503–509**
- Androctonin, a hydrophilic disulphide-bridged non-haemolytic anti-microbial peptide: a plausible mode of action  
C. Hetru, L. Letellier, Z. Oren, J.A. Hoffmann and Y. Shai **653–664**
- Characterization of a second member of the subfamily of calcium-binding mitochondrial carriers expressed in human non-excitabile tissues  
A. del Arco, M. Agudo and J. Satrústegui **725–732**

**Receptors and signal transduction**

- G<sub>i</sub>-mediated translocation of GLUT4 is independent of p85/p110 $\alpha$  and p110 $\gamma$  phosphoinositide 3-kinases but might involve the activation of Akt kinase  
L. Wang, H. Hayashi, K. Kishi, L. Huang, A. Hagi, K. Tamaoka, P.T. Hawkins and Y. Ebina **543–555**
- Quantitative assessment of complex formation of nuclear-receptor accessory proteins  
K. Graumann and A. Jungbauer **627–636**
- Signalling via ADP-ribosylation factor 6 lies downstream of phosphatidylinositide 3-kinase  
K. Venkateswarlu and P.J. Cullen **719–724**
- 14-3-3 $\zeta$  interacts with the  $\alpha$ -chain of human interleukin 9 receptor  
D. Sliva, M. Gu, Y.X. Zhu, J. Chen, S. Tsai, X. Du and Y.-C. Yang **741–747**
- Platelet-derived growth factor (PDGF)-induced activation of signal transducer and activator of transcription (Stat) 5 is mediated by PDGF  $\beta$ -receptor and is not dependent on c-Src, Fyn, Jak1 or Jak2 kinases  
K. Paukku, S. Valgeirsdóttir, P. Saharinen, M. Bergman, C.-H. Heldin and O. Silvennoinen **759–766**

**Cell biology and development**

- Glucose persistence on high-mannose oligosaccharides selectively inhibits the macroautophagic sequestration of N-linked glycoproteins  
E. Ogier-Denis, C. Bauvy, F. Cluzeaud, A. Vandewalle and P. Codogno **459–466**
- Neuroserpin is expressed in the pituitary and adrenal glands and induces the extension of neurite-like processes in AtT-20 cells  
R.M. Hill, P.K. Parmar, L.C. Coates, E. Mezey, J.F. Pearson and N.P. Birch **595–601**
- Cellular adaptation to down-regulated iron transport into lymphoid leukaemic cells: effects on the expression of the gene for ribonucleotide reductase  
C.R. Chitambar, J.P. Wereley, T. Heiman, W.E. Antholine and W.J. O'Brien **681–685**

**Corrections**

- Up-regulation of liver glucose-6-phosphatase in rats fed with a P<sub>i</sub>-deficient diet  
by W. Xie, Y. Li, M.-C. Méchin and G. van de Werve (volume 343, pages 393–396, 1999) **767**
- Normal prion protein has an activity like that of superoxide dismutase  
by D.R. Brown, B.-S. Wong, F. Hafiz, C. Clive, S.J. Haswell and I.M. Jones (volume 344, pages 1–5, 1999) **767**
- Direct interaction between p47<sup>phox</sup> and protein kinase C: evidence for targeting of protein kinase C by p47<sup>phox</sup> in neutrophils  
by E.P. Reeves, L.V. Dekker, L.V. Forbes, F.B. Wientjes, A. Grogan, D.J.C. Pappin and A.W. Segal (volume 344, pages 859–866, 1999) **767**