

**Biochemical Journal**  
**Volume 346, part 2**  
**1 March 2000**

First published on the Internet  
22 February 2000

**Research Communications**

- |   |   |         |
|---|---|---------|
| On the preparation of $\beta$ -haematin   | G. Blauer and M. Akkawi   | 249–250 |
| The ShBle resistance determinant from <i>Streptoalloteichus hindustanus</i> is expressed in <i>Haloferax volcanii</i> and confers resistance to bleomycin | S.D. Nuttall, S.E. Deutscher, R.A. Irving,<br>J.A. Serrano-Gomicia and M.L. Dyall-Smith | 251–254 |

**Research Papers**

**Proteins**

- |   |   |         |
|---|---|---------|
| Mechanism of thermal denaturation of maltodextrin phosphorylase from <i>Escherichia coli</i>  | R. Grießler, S. D'Auria, R. Schinzel, F. Tanfani and B. Nidetzky  | 255–263 |
| Stimulation of cleavage of membrane proteins by calmodulin inhibitors   | E. Díaz-Rodríguez, A. Esparís-Ogando, J.C. Montero, L. Yuste and A. Pandiella   | 359–367 |
| Ferredoxin III of <i>Desulfovibrio africanus</i> : sequencing of the native gene and characterization of a histidine-tagged form                          | J.L.H. Busch, J.L. Breton, S.L. Davy, R. James, G.R. Moore, F.A. Armstrong and A.J. Thomson                                     | 375–384 |
| Vitamin C protects against and reverses specific hypochlorous acid- and chloramine-dependent modifications of low-density lipoprotein                     | A.C. Carr, T. Tijerina and B. Frei  | 491–499 |
| GTPase mechanism and function: new insights from systematic mutational analysis of the phosphate-binding loop residue Ala <sup>30</sup> of Rab5           | Z. Liang, T. Mather and G. Li   | 501–508 |
| Mutations in the reduced-folate carrier affect protein localization and stability   | H. Sadlish, R.C. Murray, F.M.R. Williams and W.F. Flintoff  | 509–518 |
| Crystal structure of the anti-(carcinoembryonic antigen) single-chain Fv antibody MFE-23 and a model for antigen binding based on intermolecular contacts | M.K. Boehm, A.L. Corper, T. Wan, M.K. Sohi, B.J. Sutton, J.D. Thornton, P.A. Keep, K.A. Chester, R.H.J. Begent and S.J. Perkins | 519–528 |

**Enzymes**

- |   |   |         |
|---|---|---------|
| HIV-2 protease is inactivated after oxidation at the dimer interface and activity can be partly restored with methionine sulphoxide reductase       | D.A. Davis, F.M. Newcomb, J. Moskovitz, P.T. Wingfield, S.J. Stahl, J. Kaufman, H.M. Fales, R.L. Levine and R. Yarchoan   | 305–311 |
| Enzyme activity and dynamics: xylanase activity in the absence of fast anharmonic dynamics  | R.V. Dunn, V. Réat, J. Finney, M. Ferrand, J.C. Smith and R.M. Daniel   | 355–358 |
| Identification of a DNA-binding domain and an active-site residue of pseudorabies virus DNase   | T.-Y. Ho, S.-L. Wu, C.-H. Hsiang, T.-J. Chang and C.-Y. Hsiang  | 441–445 |
| Characterization of peptidyl boronic acid inhibitors of mammalian 20 S and 26 S proteasomes and their inhibition of proteasomes in cultured cells   | R.C. Gardner, S.J. Assinder, G. Christie, G.G.F. Mason, R. Markwell, H. Wadsworth, M. McLaughlin, R. King, M.C. Chabot-Fletcher, J.J. Breton, D. Allsop and A.J. Rivett | 447–454 |
| Tandem mass spectrometric analysis of <i>Aspergillus niger</i> pectin methylesterase: mode of action on fully methyl-esterified oligogalacturonates | H.C.M. Kester, J.A.E. Benen, J. Visser, M.E. Warren, R. Orlando, C. Bergmann, D. Magaud, D. Anker and A. Doutheau   | 469–474 |
| Modulation of the glutathione S-transferase in <i>Ochrobactrum anthropi</i> : function of xenobiotic substrates and other forms of stress           | B. Favaloro, A. Tamburro, M.A. Trofino, L. Bologna, D. Rotilio and H.J. Heipieper   | 553–559 |

**Carbohydrates and lipids**

- Some polyphenols inhibit the formation of pentyl radical and octanoic acid radical in the reaction mixture of linoleic acid hydroperoxide with ferrous ions  
H. Iwahashi **265–273**
- Reagent or myeloperoxidase-generated hypochlorite affects discrete regions in lipid-free and lipid-associated human apolipoprotein A-I  
C. Bergt, K. Oettl, W. Keller, F. Andreae, H.J. Leis, E. Malle and W. Sattler **345–354**
- Expression of heparan sulphate L-iduronyl 2-O-sulphotransferase in human kidney 293 cells results in increased D-glucuronyl 2-O-sulphation  
J. Rong, H. Habuchi, K. Kimata, U. Lindahl and M. Kusche-Gullberg **463–468**

**Gene structure and expression**

- Target site search and effective inhibition of leukaemic cell growth by a covalently closed multiple anti-sense oligonucleotide to *c-myc*  
I.-J. Moon, Y. Lee, C.-S. Kwak, J.-H. Lee, K. Choi, A.D. Schreiber and J.-G. Park **295–303**
- Identification and expression of an allergen Asp f 13 from *Apergillus fumigatus* and epitope mapping using human IgE antibodies and rabbit polyclonal antibodies  
L.-P. Chow, S.-L. Liu, C.-J. Yu, H.-K. Liao, J.-J. Tsai and T.-K. Tang **423–431**
- Cloning and functional characterization of the 5'-flanking region of human methionine adenosyltransferase 1A gene  
Z. Zeng, Z.Z. Huang, C. Chen, H. Yang, Z. Mao and S.C. Lu **475–482**
- Interaction between the homeodomain proteins Cdx2 and HNF1 $\alpha$  mediates expression of the lactase-phlorizin hydrolase gene  
C. Mitchelmore, J.T. Troelsen, N. Spodsberg, H. Sjöström and O. Norén **529–535**

**Regulation of metabolism**

- Zinc-regulated ubiquitin conjugation signals endocytosis of the yeast ZRT1 zinc transporter  
R.S. Gitan and D.J. Eide **329–336**
- Thyroxine regulation of monolysocardiolipin acyltransferase activity in rat heart  
T. Mutter, V.W. Dolinsky, B.J. Ma, W.A. Taylor and G.M. Hatch **403–406**
- Continuous exposure to high concentrations of nitric oxide leads to persistent inhibition of oxygen consumption by J774 cells as well as extraction of oxygen by the extracellular medium  
A. Orsi, B. Beltrán, E. Clement, K. Hallén, M. Feelisch and S. Moncada **407–412**
- Investigation of the mechanism by which glucose analogues cause translocation of glucokinase in hepatocytes: evidence for two glucose binding sites  
L. Agius and M. Stubbs **413–421**
- Plasmodium falciparum*-infected red blood cells depend on a functional glutathione *de novo* synthesis attributable to an enhanced loss of glutathione  
K. Lüersen, R.D. Walter and S. Müller **545–552**

**Membranes and bioenergetics**

- Estimation of systolic and diastolic free intracellular Ca<sup>2+</sup> by titration of Ca<sup>2+</sup> buffering in the ferret heart  
H.L. Kirschenlohr, A.A. Grace, J.I. Vandenberg, J.C. Metcalfe and G.A. Smith **385–391**
- Ca<sup>2+</sup> buffering in the heart: Ca<sup>2+</sup> binding to and activation of cardiac myofibrils  
G.A. Smith, H.B.F. Dixon, H.L. Kirschenlohr, A.A. Grace, J.C. Metcalfe and J.I. Vandenberg **393–402**

**Receptors and signal transduction**

- Ca<sup>2+</sup> and calmodulin differentially modulate myo-inositol 1,4,5-trisphosphate (IP<sub>3</sub>)-binding to the recombinant ligand-binding domains of the various IP<sub>3</sub> receptor isoforms  
S. Vanlingen, H. Sipma, P. De Smet, G. Callewaert, L. Missiaen, H. De Smedt and J.B. Parys **275–280**

---

Pancreatic eukaryotic initiation factor-2 $\alpha$ kinase (PEK) homologues in humans, <i>Drosophila melanogaster</i> and <i>Caenorhabditis elegans</i> that mediate translational control in response to endoplasmic reticulum stress	R. Sood, A.C. Porter, K. Ma, L.A. Quilliam and R.C. Wek	281–293
Inhibition of atrial natriuretic peptide (ANP) C receptor expression by antisense oligodeoxynucleotides in A10 vascular smooth-muscle cells is associated with attenuation of ANP-C-receptor-mediated inhibition of adenylyl cyclase	A. Palaparti, Y. Li and M.B. Anand-Srivastava	313–320
Down-regulation of cyclic-nucleotide phosphodiesterase 3B in 3T3-L1 adipocytes induced by tumour necrosis factor $\alpha$ and cAMP	T.R. Landström, J. Mei, M. Karlsson, V. Manganiello and E. Degerman	337–343
Stress-induced generation of <i>N</i> -acylethanolamines in mouse epidermal JB6 P <sup>+</sup> cells	E.V. Berdyshev, P.C. Schmid, Z. Dong and H.H.O. Schmid	369–374
Protein phosphatase 2A is associated in an inactive state with microtubules through 2A1-specific interaction with tubulin	A. Hiraga and S. Tamura	433–439
Involvement of G $\alpha$ 2 in sodium butyrate-induced erythroblastic differentiation of K562 cells	M.G. Davis, Y. Kawai and I.J. Arinze	455–461
The N-terminal 34 residues of the 55 kDa regulatory subunits of phosphoinositide 3-kinase interact with tubulin	K. Inukai, M. Funaki, M. Nawano, H. Katagiri, T. Ogihara, M. Anai, Y. Onishi, H. Sakoda, H. Ono, Y. Fukushima, M. Kikuchi, Y. Oka and T. Asano	483–489
<b>Cell biology and development</b>		
Actin filaments play a critical role in insulin-induced exocytic recruitment but not in endocytosis of GLUT4 in isolated rat adipocytes	W. Omata, H. Shibata, L. Li, K. Takata and I. Kojima	321–328
Rapid replenishment of sphingomyelin in the plasma membrane upon degradation by sphingomyelinase in NIH3T3 cells overexpressing the phosphatidylinositol transfer protein $\beta$	C.M. van Tiel, C. Luberto, G.T. Snoek, Y.A. Hannun and K.W.A. Wirtz	537–543