821-827

Biochemical Journal Volume 347, part 3 1 May 2000

First published on the Internet 25 April 2000 http://www.BiochemJ.org

| | | http://www.BiochemJ.org |
|--|--|-------------------------|
| Review Article | | |
| Role of lipids in the translocation of proteins across membranes by F. van Voorst and B. de Kruijff | | 601–612 |
| Research Papers | | |
| Proteins | | |
| Large-scale synthesis and functional elements for the antimicrobial activity of defensins | P.A. Raj, K.J. Antonyraj and T. Karunakaran | 633–641 |
| Binding of α_2 -macroglobulin and limulin: regulation of the plasma haemolytic system of the American horseshoe crab, $Limulus$ | S. Swarnakar, R. Asokan, J.P. Quigley and P.B. Armstrong | 679–685 |
| Conserved sequences in the β subunit of archaeal and eukaryal translation initiation factor 2 (eIF2), absent from eIF5, mediate interaction with eIF2 γ | G.M. Thompson, E. Pacheco, E.O. Melo and B.A. Castilho | 703–709 |
| The domains of human fibronectin mediating the binding of α antigen, the most immunopotent antigen of mycobacteria that induces protective immunity against mycobacterial infection | M. Naito, T. Fukuda, K. Sekiguchi and T. Yamada | 725–731 |
| A novel 4 S $[^3H]\beta$ -naphthoflavone-binding protein in liver cytosol of female Sprague-Dawley rats treated with aryl hydrocarbon receptor agonists | D. Brauze and D. Malejka-Giganti | 787–795 |
| Alanine mutagenesis of high-mobility-group- protein-1 box B (HMG1-B) | S. Taudte, H. Xin and N.R. Kallenbach | 807–814 |
| Endonuclease activity in lipocalins | T.N. Yusifov, A.R. Abduragimov, O.K. Gasymov and B.J. Glasgow | 815–819 |
| Purification and cloning of phosphatidylinositol transfer proteins from <i>Dictyostelium discoideum</i> : homologues of both mammalian PITPs and <i>Saccharomyces cerevisiae</i> Sec14p are found in the same cell | P. Swigart, R. Insall, A. Wilkins and S. Cockcroft | 837–843 |
| Heteroduplex DNA and ATP induced conformational changes of a MutS mismatch repair protein from <i>Thermus aquaticus</i> | I. Biswas and R. Vijayvargia | 881–886 |
| Enzymes | | |
| Purification and characterization of a Baeyer-Villiger mono-oxygenase from Rhodococcus erythropolis DCL14 involved in three different monocyclic monoterpene degradation pathways | M.J. van der Werf | 693–701 |
| Shedding of somatic angiotensin-converting enzyme (ACE) is inefficient compared with testis ACE despite cleavage at identical stalk sites | Z.L. Woodman, S.Y. Oppong, S. Cook, N.M. Hooper, S.L.U. Schwager, W.F. Brandt, M.R.W. Ehlers and E.D. Sturrock | 711–718 |
| Identification of the catalytic residues of the first family of β (1-3)glucanosyltransferases identified in fungi | I. Mouyna, M. Monod, T. Fontaine, B. Henrissat, B. Léchenne and JP. Latgé | 741–747 |

S. Reddy, A.D. Jones, C.E. Cross, P.S.-Y. Wong and

A. van der Vliet

Inactivation of creatine kinase by

S-glutathionylation of the active-site cysteine residue

| Sugar-nucleotide-binding and autoglycosylating polypeptide(s) from nasturtium fruit: biochemical capacities and potential functions | A. Faik, D. Desveaux and G. Maclachlan | 857–864 |
|---|---|---------|
| Epoxyalkyl glycosides of D-xylose and xylo-oligosaccharides are active-site markers of xylanases from glycoside hydrolase family 11, not from family 10 | P. Ntarima, W. Nerinckx, K. Klarskov, B. Devreese, M.K. Bhat, J. van Beeumen and M. Claeyssens | 865–873 |
| Gene structure and expression | | |
| Differential adrenergic regulation of the gene expression of the β -adrenoceptor subtypes β_1 , β_2 and β_3 in brown adipocytes | T. Bengtsson, B. Cannon and J. Nedergaard | 643–651 |
| Prominent expression of the selenoprotein thioredoxin reductase in the medullary rays of the rat kidney and thioredoxin reductase mRNA variants differing at the 5' untranslated region | AK. Rundlöf, M. Carlsten, M.M.J. Giacobini and E.S.J. Arnér | 661–668 |
| cDNA cloning and characterization of human $\Delta^{\text{5}}\text{-desaturase}$ involved in the biosynthesis of arachidonic acid | A.E. Leonard, B. Kelder, E.G. Bobik, LT. Chuang, J.M. Parker-Barnes, J.M. Thurmond, P.E. Kroeger, J.J. Kopchick, YS. Huang and P. Mukerji | 719–724 |
| Involvement of NRAMP1 from <i>Arabidopsis thaliana</i> in iron transport | C. Curie, J.M. Alonso, M. Le Jean, J.R. Ecker and JF. Briat | 749–755 |
| Identification of abundant mRNAs from the third stage larvae of the parasitic nematode, <i>Ostertagia</i> ostertagi | J. Moore, L. Tetley and E. Devaney | 763–770 |
| Regulation of metabolism | | |
| Isolation and characterization of protoporphyrin glycoconjugates from rat Harderian gland by HPLC, capillary electrophoresis and HPLC/electrospray ionization MS | C.K. Lim, M.A. Razzaque, J. Luo and P.B. Farmer | 757–761 |
| Membranes and bioenergetics | | |
| Oxidative phosphorylation in cardiolipin-lacking yeast mitochondria | V. Koshkin and M.L. Greenberg | 687–691 |
| Intragenic and intergenic suppression of the Escherichia coli ATP synthase subunit a mutation of Gly-213 to Asn: functional interactions between residues in the proton transport site | P.H. Kuo and R.K. Nakamoto | 797–805 |
| Receptors and signal transduction | | |
| Cytosolic adaptor protein Dab2 is an intracellular ligand of endocytic receptor gp600/megalin | A.V. Oleinikov, J. Zhao and S.P. Makker | 613–621 |
| The human vasoactive intestinal peptide/pituitary adenylate cyclase-activating peptide receptor 1 (VPAC1) promoter: characterization and role in receptor expression during enterocytic differentiation of the colon cancer cell line Caco-2CI.20 | A. Couvineau, JJ. Maoret, C. Rouyer-Fessard, I. Carrero and M. Laburthe | 623–632 |
| Persistent activation of $G_s\alpha$ through limited proteolysis by calpain | K. Sato-Kusubata, Y. Yajima and S. Kawashima | 733–740 |
| Mutational analysis of the N-linked glycosylation sites of the human insulin receptor | T.C. Elleman, M.J. Frenkel, P.A. Hoyne, N.M. McKern, L. Cosgrove, D.R. Hewish, K.M. Jachno, J.D. Bentley, S.E. Sankovich and C.W. Ward | 771–779 |
| Protein kinase C activation by acidic proteins including 14-3-3 | P.C.J. van der Hoeven, J.C.M. van der Wal, P. Ruurs and W.J. van Blitterswijk | 781–785 |
| Conserved Ca ²⁺ -antagonist-binding properties and putative folding structure of a recombinant high-affinity dihydropyridine-binding domain | I. Huber, E. Wappl, A. Herzog, J. Mitterdorfer, H. Glossmann, T. Langer and J. Striessnig | 829–836 |

| | | Contents | xiii |
|---|--|----------|---------|
| A novel 50 kDa protein forms complexes with protein phosphatase 4 and is located at centrosomal microtubule organizing centres | C.J. Hastie, G.K. Carnegie, N. Morrice and P.T.W. Cohen | | 845–855 |
| Cell biology and development | | | |
| Identification of a C-terminal cdc25 sequence required for promotion of germinal vesicle breakdown | E.A. Powers, D.P. Thompson, P.A. Garner-Hamrick, W. He, A.W. Yem, C.A. Bannow, D.J. Staples, G.A. Waszak, C.W. Smith, M.R. Deibel, Jr. and C. Fisher | | 653–660 |
| Serine protease inhibitors suppress cytochrome c-mediated caspase-9 activation and apoptosis during hypoxia-reoxygenation | Z. Dong, P. Saikumar, Y. Patel, J.M. Weinberg and M.A. Venkatachalam | | 669–677 |
| Polyamines directly induce release of cytochrome c from heart mitochondria | C. Stefanelli, I. Stanic, M. Zini, F. Bonavita, F. Flamigni, L. Zambonin, L. Landi, C. Pignatti, C. Guarnieri and C.M. Caldarera | | 875–880 |
| Correction | | | |
| A pyruvate-proton symport and an H ⁺ -ATPase regulate the intracellular pH of <i>Trypanosoma brucei</i> at different stages of its life cycle by N. Vanderheyden, J. Wong and R. Docampo (volume 346, pages 53–62, 2000) | | | 887 |