

## Review Article

- Strategies for manipulating the p53 pathway in the treatment of human cancer** 1–17  
*by T.R. Hupp, D.P. Lane and K.L. Ball*

## Research Papers

- Changes in protein kinase C  $\epsilon$  phosphorylation status and intracellular localization as 3T3 and 3T6 fibroblasts grow to confluency and quiescence: a role for phosphorylation at Ser-729? K. England and M.G. Rumsby 19–26
- The mechanism of Fe<sup>2+</sup>-initiated lipid peroxidation in liposomes: the dual function of ferrous ions, the roles of the pre-existing lipid peroxides and the lipid peroxy radical L. Tang, Y. Zhang, Z. Qian and X. Shen 27–36
- Cloning, expression and gene organization of a human Neu5Ac $\alpha$ 2–3Gal $\beta$ 1–3GalNAc  $\alpha$ 2,6-sialyltransferase: hST6GalNAc IV A. Harduin-Lepers, D.C. Stokes, W.F.A. Steelant, B. Samyn-Petit, M.-A. Krzewinski-Recchi, V. Vallejo-Ruiz, J.-P. Zanetta, C. Augé and P. Delannoy 37–48
- Polydom: a secreted protein with pentraxin, complement control protein, epidermal growth factor and von Willebrand factor A domains D. Gilgès, M.-A. Vinit, I. Callebaut, L. Coulombel, V. Cacheux, P.-H. Romeo and I. Vigon 49–59
- Calcineurin regulates ryanodine receptor/ Ca<sup>2+</sup>-release channels in rat heart A. Bandyopadhyay, D.W. Shin, J.O. Ahn and D.H. Kim 61–70
- Role of glutathione in heat-shock-induced cell death of *Saccharomyces cerevisiae* K.-i. Sugiyama, A. Kawamura, S. Izawa and Y. Inoue 71–78
- Diabetes-induced changes in specific lipid molecular species in rat myocardium X. Han, D.R. Abendschein, J.G. Kelley and R.W. Gross 79–89
- Effect of alpha-1 antitrypsin Portland variant ( $\alpha$ 1-PDX) on HIV-1 replication B. Bahbouhi, M. Bendjennat, D. Guétard, N.G. Seidah and E. Bahraoui 91–98
- Thermodynamic mixing of molecular states of the epidermal growth factor receptor modulates macroscopic ligand binding affinity M.R. Holbrook, L.L. Slakey and D.J. Gross 99–108
- 3'-Untranslated regions of oxidative phosphorylation mRNAs function *in vivo* as enhancers of translation C.M. Di Liegro, M. Bellafiore, J.M. Izquierdo, A. Rantanen and J.M. Cuezva 109–115
- In vitro* membrane-inserted conformation of the cytochrome *b*<sub>5</sub> tail M.R. Hanlon, R.R. Begum, R.J. Newbold, D. Whitford and B.A. Wallace 117–124
- Macrophage cholesteryl ester hydrolases and hormone-sensitive lipase prefer specifically oxidized cholesteryl esters as substrates over their non-oxidized counterparts J. Belkner, H. Stender, H.-G. Holzhütter, C. Holm and H. Kühn 125–134
- Lysophosphatidic acid prevents apoptosis in fibroblasts via G<sub>i</sub>-protein-mediated activation of mitogen-activated protein kinase X. Fang, S. Yu, R. LaPushin, Y. Lu, T. Furui, L.Z. Penn, D. Stokoe, J.R. Erickson, R.C. Bast, Jr and G.B. Mills 135–143
- Synergistic activation of stress-activated protein kinase 1/c-Jun N-terminal kinase (SAPK1/JNK) isoforms by mitogen-activated protein kinase kinase 4 (MKK4) and MKK7 Y. Fleming, C.G. Armstrong, N. Morrice, A. Paterson, M. Goedert and P. Cohen 145–154
- Catalytic activation of mitogen-activated protein (MAP) kinase phosphatase-1 by binding to p38 MAP kinase: critical role of the p38 C-terminal domain in its negative regulation D. Hutter, P. Chen, J. Barnes and Y. Liu 155–163

|   |   |                |
|---|---|----------------|
| Identification of Rab6 as an <i>N</i> -ethylmaleimide-sensitive fusion protein-binding protein  | S.Y. Han, D.Y. Park, S.D. Park and S.H. Hong  | <b>165–173</b> |
| Divergence in the anti-apoptotic signalling pathways used by nerve growth factor and basic fibroblast growth factor (bFGF) in PC12 cells: rescue by bFGF involves protein kinase C $\delta$ | M.M. Wert and H.C. Palfrey  | <b>175–182</b> |
| Lentil seed aquaporins form a hetero-oligomer which is phosphorylated by a Mg <sup>2+</sup> -dependent and Ca <sup>2+</sup> -regulated kinase   | P. Harvengt, A. Vlerick, B. Fuks, R. Wattiez, J.-M. Ruyschaert and F. Homble                            | <b>183–190</b> |
| Bovine prion protein as a modulator of protein kinase CK2   | F. Meggio, A. Negro, S. Sarno, M. Ruzzene, A. Bertoli, M.C. Sorgato and L.A. Pinna                      | <b>191–196</b> |
| Angiotensin II stimulates cyclic ADP-ribose formation in neonatal rat cardiac myocytes  | H. Higashida, J.-S. Zhang, M. Hashii, M. Shintaku, C. Higashida and Y. Takeda                           | <b>197–202</b> |
| Dominance of G <sub>s</sub> in doubly G <sub>s</sub> /G <sub>i</sub> -coupled chimaeric A <sub>1</sub> /A <sub>2A</sub> adenosine receptors in HEK-293 cells                                | A.L. Tucker, L. Jia, D. Holeton, A.J. Taylor and J. Linden  | <b>203–210</b> |
| Regulation by glucagon (cAMP) and insulin of the promoter of the human phosphoenolpyruvate carboxykinase gene (cytosolic) in cultured rat hepatocytes and in human hepatoblastoma cells     | A.K. Rucktäschel, D.K. Granner and B. Christ  | <b>211–217</b> |
| Peroxynitrite activates the phosphoinositide 3-kinase/Akt pathway in human skin primary fibroblasts   | L.-O. Klotz, S.M. Schieke, H. Sies and N.J. Holbrook  | <b>219–225</b> |
| Dietary P <sub>i</sub> deprivation in rats affects liver cAMP, glycogen, key steps of gluconeogenesis and glucose production  | W. Xie, T.L. Tran, D.T. Finegood and G. van de Werve  | <b>227–232</b> |
| Endocytic uptake of advanced glycation end products by mouse liver sinusoidal endothelial cells is mediated by a scavenger receptor distinct from the macrophage scavenger receptor class A | K. Matsumoto, H. Sano, R. Nagai, H. Suzuki, T. Kodama, M. Yoshida, S. Ueda, B. Smedsrød and S. Horiuchi | <b>233–240</b> |