

**Biochemical Journal**  
**Volume 353, part 2**  
**15 January 2001**

First published on the Internet  
8 January 2001  
<http://www.BiochemJ.org>

**Subscribing organizations are encouraged to copy and distribute this table of contents for non-commercial purposes**

**Research Communications**

Chondrocyte expressed protein-68 (CEP-68), a novel human marker gene for cultured chondrocytes

E. Steck, K. Benz, H. Lorenz, M. Loew, T. Gress and W. Richter

**169–174**

Mitochondrial priming modifies  $\text{Ca}^{2+}$  oscillations and insulin secretion in pancreatic islets

E.K. Ainscow and G.A. Rutter

**175–180**

**Research Papers**

Haem-linked interactions in horseradish peroxidase revealed by spectroscopic analysis of the Phe-221 → Met mutant

B.D. Howes, N.C. Veitch, A.T. Smith, C.G. White and G. Smulevich

**181–191**

Dual DNA-binding specificity of peroxisome-proliferator-activated receptor  $\gamma$  controlled by heterodimer formation with retinoid X receptor  $\alpha$

M. Okuno, E. Arimoto, Y. Ikenobu, T. Nishihara and M. Imagawa

**193–198**

Identification of essential active-site residues in the cyanogenic  $\beta$ -glucosidase (linamarase) from cassava (*Manihot esculenta* Crantz) by site-directed mutagenesis

Z. Keresztessy, K. Brown, M.A. Dunn and M.A. Hughes

**199–205**

*Escherichia coli* flavohaemoglobin (Hmp) with equimolar FAD and haem contents has a low affinity for dioxygen in the absence or presence of nitric oxide

C.E. Mills, S. Sedelnikova, B. Søballe, M.N. Hughes and R.K. Poole

**207–213**

Phosphate exchange and ATP synthesis by DMSO-pretreated purified bovine mitochondrial ATP synthase

S. Beharry and P.D. Bragg

**215–222**

Negative cross-talk between interleukin-3 and interleukin-11 is mediated by suppressor of cytokine signalling-3 (SOCS-3)

F. Magrangeas, O. Boisteaup, S. Denis, Y. Jacques and S. Minvielle

**223–230**

Fatty-acyl-CoA thioesters inhibit recruitment of steroid receptor co-activator 1 to  $\alpha$  and  $\gamma$  isoforms of peroxisome-proliferator-activated receptors by competing with agonists

K. Murakami, T. Ide, T. Nakazawa, T. Okazaki, T. Mochizuki and T. Kadokawa

**231–238**

*Triticum aestivum* L. endoxylanase inhibitor (TAXI) consists of two inhibitors, TAXI I and TAXI II, with different specificities

K. Gebruers, W. Debysen, H. Goesaert, P. Proost, J. Van Damme and J.A. Delcour

**239–244**

Glutamine metabolism to glucosamine is necessary for glutamine inhibition of endothelial nitric oxide synthesis

G. Wu, T.E. Haynes, H. Li, W. Yan and C.J. Meininger

**245–252**

Peroxisomal bifunctional enzyme binds and activates the activation function-1 region of the peroxisome-proliferator-activated receptor  $\alpha$

C.E. Juge-Aubry, S. Kuenzli, J.-C. Sanchez, D. Hochstrasser and C.A. Meier

**253–258**

Catalytic and spectroscopic analysis of blue copper-containing nitrite reductase mutants altered in the environment of the type 2 copper centre: implications for substrate interaction

M. Prudêncio, R.R. Eady and G. Sawers

**259–266**

Insulin inhibits glucocorticoid-stimulated L-type 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase gene expression by activation of the c-Jun N-terminal kinase pathway

E. de los Pinos, S. Fernández de Mattos, M. Joaquin and A. Tauler

**267–273**

---

|   |   |                |
|---|---|----------------|
| Analysis of mitogen-activated protein kinase pathways used by interleukin 1 in tissues <i>in vivo</i> : activation of hepatic c-Jun N-terminal kinases 1 and 2, and mitogen-activated protein kinase kinases 4 and 7                | A. Finch, W. Davis, W.G. Carter and J. Saklatvala   | <b>275–281</b> |
| Cloning and functional expression of rat kidney dipeptidyl peptidase II   | K.M. Fukasawa, K. Fukasawa, K. Higaki, N. Shiina, M. Ohno, S. Ito, J. Otogoto and N. Ota                                | <b>283–290</b> |
| $\gamma$ -Interferon decreases the level of 26 S proteasomes and changes the pattern of phosphorylation   | S. Bose, P. Brooks, G.G.F. Mason and A.J. Rivett  | <b>291–297</b> |
| Resistance of bone marrow-derived macrophages to apoptosis is associated with the expression of X-linked inhibitor of apoptosis protein in primary cultures of bone marrow cells  | H. Lin, C. Chen and B.D.-M. Chen  | <b>299–306</b> |
| 5-Aminolaevulinate synthase gene promoter contains two cAMP-response element (CRE)-like sites that confer positive and negative responsiveness to CRE-binding protein (CREB)  | L.E. Giono, C.L. Varone and E.T. Cáñepa   | <b>307–316</b> |
| Cross-talk between steroid-receptor-mediated and cell-membrane-receptor-mediated signalling pathways results in the <i>in vivo</i> modulation of c-Met and ornithine decarboxylase gene expression in mouse kidney                  | M. Dudkowska, A. Stachurska, W. Chmurzyńska, B. Grzelakowska-Sztabert and M. Manteuffel-Cymborowska                     | <b>317–323</b> |
| Activation of blood coagulation factor IX by gingipains R, arginine-specific cysteine proteinases from <i>Porphyromonas gingivalis</i>  | T. Imamura, S. Tanase, T. Hamamoto, J. Potempa and J. Travis  | <b>325–331</b> |
| Inhibition of prolyl 4-hydroxylase <i>in vitro</i> and <i>in vivo</i> by members of a novel series of phenanthrolinones   | T.J. Franklin, W.P. Morris, P.N. Edwards, M.S. Large and R. Stephenson  | <b>333–338</b> |
| Endogenous glycosylphosphatidylinositol-specific phospholipase C releases renal dipeptidase from kidney proximal tubules <i>in vitro</i>  | S.W. Park, K. Choi, I.C. Kim, H.H.B. Lee, N.M. Hooper and H.S. Park   | <b>339–344</b> |
| Probing a novel potato lipoxygenase with dual positional specificity reveals primary determinants of substrate binding and requirements for a surface hydrophobic loop and has implications for the role of lipoxygenases in tubers | R.K. Hughes, S.I. West, A.R. Hornostaj, D.M. Lawson, S.A. Fairhurst, R.O. Sanchez, P. Hough, B.H. Robinson and R. Casey | <b>345–355</b> |
| Oxidation of pyridine nucleotides during Fas- and ceramide-induced apoptosis in Jurkat cells: correlation with changes in mitochondria, glutathione depletion, intracellular acidification and caspase 3 activation                 | M.-C. Gendron, N. Schrantz, D. Métivier, G. Kroemer, Z. Maciorowska, F. Sureau, S. Koester and P.X. Petit               | <b>357–367</b> |
| Overexpression of the human 2-oxoglutarate carrier lowers mitochondrial membrane potential in HEK-293 cells: contrast with the unique cold-induced mitochondrial carrier CGI-69   | X.X. Yu, D.A. Lewin, A. Zhong, J. Brush, P.W. Schow, S.W. Sherwood, G. Pan and S.H. Adams                               | <b>369–375</b> |
| Interaction of the $\alpha$ subunit of Na,K-ATPase with cofilin   | K. Lee, J. Jung, M. Kim and G. Guidotti   | <b>377–385</b> |
| Identification of Cys <sup>140</sup> in helix 4 as an exofacial cysteine residue within the substrate-translocation channel of rat equilibrative nitrobenzylthioinosine (NBMPR)-insensitive nucleoside transporter rENT2            | S.Y.M. Yao, M. Sundaram, E.G. Chomey, C.E. Cass, S.A. Baldwin and J.D. Young  | <b>387–393</b> |
| Studies on a single immunoglobulin-binding domain of protein L from <i>Peptostreptococcus magnus</i> : the role of tyrosine-53 in the reaction with human IgG   | J.A. Beckingham, N.G. Housden, N.M. Muir, S.P. Bottomley and M.G. Gore  | <b>395–401</b> |
| Characterization of monocot and dicot plant S-adenosyl-L-methionine decarboxylase gene families including identification in the mRNA of a highly conserved pair of upstream overlapping open reading frames                         | M. Franceschetti, C. Hanfrey, S. Scaramagli, P. Torrigiani, N. Bagni, D. Burtin and A.J. Michael                        | <b>403–409</b> |
| Mitochondrial respiratory chain-dependent generation of superoxide anion and its release into the intermembrane space   | D. Han, E. Williams and E. Cadena   | <b>411–416</b> |