

Biochemical Journal
Volume 355, part 1
1 April 2001

First published on the Internet
26 March 2001
<http://www.BiochemJ.org>

Subscribing organizations are encouraged to copy and distribute this table of contents for non-commercial purposes

Review

Recombinant aequorin and green fluorescent protein as valuable tools in the study of cell signalling

by A. Chiesa, E. Rapizzi, V. Tosello, P. Pinton,
M. de Virgilio, K.E. Fogarty and R. Rizzuto



1–12

Research Papers

Organization of uroplakin subunits:
transmembrane topology, pair formation and
plaque composition

Identification of the G13 (cAMP-response-element-binding protein-related protein) gene product related to activating transcription factor 6 as a transcriptional activator of the mammalian unfolded protein response

α -Melanocyte-related tripeptide, Lys-D-Pro-Val, ameliorates endotoxin-induced nuclear factor κ B translocation and activation: evidence for involvement of an interleukin-1 $\beta^{193-195}$ receptor antagonism in the alveolar epithelium

Kinetics of CO binding and CO photodissociation in *Pseudomonas stutzeri* cd₁ nitrite reductase: probing the role of extended N-termini in fast structural relaxation upon CO photodissociation

Characterization of the interaction between the transcription factors human polyamine modulated factor (PMF-1) and NF-E2-related factor 2 (Nrf-2) in the transcriptional regulation of the spermidine/spermine N¹-acetyltransferase (SSAT) gene

The *Escherichia coli* CcmG protein fulfils a specific role in cytochrome c assembly

Selective recognition of inositol phosphates by subtypes of the inositol trisphosphate receptor

Xenobiotic induction of cytochrome P450 2B1 (CYP2B1) is mediated by the orphan nuclear receptor constitutive androstane receptor (CAR) and requires steroid co-activator 1 (SRC-1) and the transcription factor Sp1

Long-lived glycosyl-enzyme intermediate mimic produced by formate re-activation of a mutant endoglucanase lacking its catalytic nucleophile

Nitric oxide and cGMP activate Ca²⁺-release processes in rat parotid acinar cells

Enzymic characterization of epidermis-derived 12-lipoxygenase isoenzymes

Differential regulation of expression of genes encoding uncoupling proteins 2 and 3 in brown adipose tissue during lactation in mice

F.-X. Liang, I. Riedel, F.-M. Deng, G. Zhou, C. Xu,
X.-R. Wu, X.-P. Kong, R. Moll and T.-T. Sun

13–18

K. Haze, T. Okada, H. Yoshida, H. Yanagi, T. Yura,
M. Negishi and K. Mori

19–28

J.J.E. Haddad, R. Lauterbach, N.E. Saadé,
B. Safieh-Garabedian and S.C. Land

29–38

E.K. Wilson, A. Bellelli, F. Cutruzzolà, W.G. Zumft,
A. Gutierrez and N.S. Scrutton

39–43

Y. Wang, W. Devereux, T.M. Stewart and R.A. Casero, Jr

45–49

E. Reid, J. Cole and D.J. Eaves

51–58

E.P. Nerou, A.M. Riley, B.V.L. Potter and C.W. Taylor

59–69

R. Muangmoonchai, D. Smirlis, S.-C. Wong, M. Edwards,
I.R. Phillips and E.A. Sheppard

71–78

J.-L. Viladot, F. Canals, X. Batllori and A. Planas

79–86

D.K. Looms, K. Tritsaris, B. Nauntofte and S. Dissing

87–95

M. Siebert, P. Krieg, W.D. Lehmann, F. Marks and
G. Fürstenberger

97–104

N. Pedraza, G. Solanes, R. Iglesias, M. Vázquez, M. Giralt
and F. Villarroya

105–111

| | | |
|--|--|----------------|
| Characterization of a novel phosphatidylinositol 3-phosphate-binding protein containing two FYVE fingers in tandem that is targeted to the Golgi | P.C.F. Cheung, L. Trinkle-Mulcahy, P. Cohen and J.M. Lucocq | 113–121 |
| Proinsulin C-peptide rapidly stimulates mitogen-activated protein kinases in Swiss 3T3 fibroblasts: requirement of protein kinase C, phosphoinositide 3-kinase and pertussis toxin-sensitive G-protein | T. Kitamura, K. Kimura, B.D. Jung, K. Makondo, S. Okamoto, X. Cañas, N. Sakane, T. Yoshida and M. Saito | 123–129 |
| Protein–coenzyme interactions in adenosylcobalamin-dependent glutamate mutase | M.S. Huhta, H.-P. Chen, C. Hemann, C.R. Hille and E.N.G. Marsh | 131–137 |
| Control of Ca^{2+} influx in human neutrophils by inositol 1,4,5-trisphosphate (IP_3) binding: differential effects of micro-injected IP_3 receptor antagonists | E.V. Davies-Cox, I. Laffafian and M.B. Hallett | 139–143 |
| Oxidative modification of H-ras: S-thiolation and S-nitrosylation of reactive cysteines | R.J. Mallis, J.E. Buss and J.A. Thomas | 145–153 |
| Pectate lyase 10A from <i>Pseudomonas cellulosa</i> is a modular enzyme containing a family 2a carbohydrate-binding module | I.E. Brown, M.H. Mallen, S.J. Charnock, G.J. Davies and G.W. Black | 155–165 |
| A new family of rhamnogalacturonan lyases contains an enzyme that binds to cellulose | V.A. McKie, J.-P. Vincken, A.G.J. Voragen, L.A.M. van den Broek, E. Stimson and H.J. Gilbert | 167–177 |
| Exonic Sp1 sites are required for neural-specific expression of the glycine receptor β subunit gene | H. Tintrup, M. Fischer, H. Betz and J. Kuhse | 179–187 |
| Sphingosylphosphocholine is a naturally occurring lipid mediator in blood plasma: a possible role in regulating cardiac function via sphingolipid receptors | K. Liliom, G. Sun, M. Büinemann, T. Virág, N. Nusser, D.L. Baker, D.-a. Wang, M.J. Fabian, B. Brandts, K. Bender, A. Eickel, K.U. Malik, D.D. Miller, D.M. Desiderio, G. Tigyi and L. Pott | 189–197 |
| Effect of polyamine depletion on caspase activation: a study with spermine synthase-deficient cells | C. Stefanelli, C. Pignatti, B. Tantini, M. Fattori, I. Stanic, C.A. Mackintosh, F. Flamigni, C. Guarneri, C.M. Calderara and A.E. Pegg | 199–206 |
| Enhancement of macrophage survival and DNA synthesis by oxidized-low-density-lipoprotein (LDL)-derived lipids and by aggregates of lightly oxidized LDL | J.A. Hamilton, W. Jessup, A.J. Brown and G. Whitty | 207–214 |
| Recombinant soluble betaglycan is a potent and isoform-selective transforming growth factor- β neutralizing agent | M.M. Vilchis-Landeros, J.L. Montiel, V. Mendoza, G. Mendoza-Hernández and F. López-Casillas | 215–222 |
| Structural and functional similarities between the central eukaryotic initiation factor (eIF)4A-binding domain of mammalian eIF4G and the eIF4A-binding domain of yeast eIF4G | D. Dominguez, E. Kislig, M. Altmann and H. Trachsel | 223–230 |
| The mitochondrial consequences of uncoupling intact cells depend on the nature of the exogenous substrate | B. Sible, C. Filippi, M.-A. Piquet, P. Leclercq, E. Fontaine, X. Ronot, M. Rigoulet and X. Leverve | 231–235 |
| (S)-Preferential detoxification of 4-hydroxy-2(<i>E</i>)-nonenal enantiomers by hepatic glutathione S-transferase isoforms in guinea-pigs and rats | A. Hiratsuka, K. Tobita, H. Saito, Y. Sakamoto, H. Nakano, K. Ogura, T. Nishiyama and T. Watabe | 237–244 |
| The efficiency of N-linked glycosylation of bovine DNase I depends on the Asn-Xaa-Ser/Thr sequence and the tissue of origin | A. Nishikawa and S. Mizuno | 245–248 |

