

**Review****Functional architecture in the cell nucleus**  
**by M. Dundr and T. Misteli**

297–310

**Research Communication**

Repression of the heavy ferritin chain increases the labile iron pool of human K562 cells

O. Kakhlon, Y. Gruenbaum and Z.I. Cabantchik

311–316

**Research Papers**

Down-regulation of intestinal scavenger receptor class B, type I (SR-BI) expression in rodents under conditions of deficient bile delivery to the intestine

P.J. Voshol, M. Schwarz, A. Rigotti, M. Krieger, A.K. Groen and F. Kuipers

317–325

Homology modelling and structural analysis of human arylamine *N*-acetyltransferase NAT1: evidence for the conservation of a cysteine protease catalytic domain and an active-site loop

F. Rodrigues-Lima, C. Deloménie, G.H. Goodfellow, D.M. Grant and J.-M. Dupret

327–334

Characterization of a differentially expressed protein that shows an unusual localization to intracellular membranes in *Leishmania major*

E. Knuepfer, Y.-D. Stierhof, P.G. McKean and D.F. Smith

335–344

Use of fluorescent Ca<sup>2+</sup> dyes with green fluorescent protein and its variants: problems and solutions

S. Bolsover, O. Ibrahim, N. O'Lunaigh, H. Williams and S. Cockcroft

345–352

Heat shock stimulation of a tilapia heat shock protein 70 promoter is mediated by a distal element

A. Molina, E. Di Martino, J.A. Martial and M. Muller

353–359

Effects of histatin 5 and derived peptides on *Candida albicans*

A.L.A. Ruissen, J. Groenink, E.J. Helmerhorst, E. Walgreen-Weterings, W. van't Hof, E.C.I. Veerman and A.V. Nieuw Amerongen

361–368

Uterocalin, a lipocalin provisioning the preattachment equine conceptus: fatty acid and retinol binding properties, and structural characterization

S. Suire, F. Stewart, J. Beauchamp and M.W. Kennedy

369–376

The neurofibromatosis 2 protein product merlin selectively binds F-actin but not G-actin, and stabilizes the filaments through a lateral association

M.F. James, N. Manchanda, C. Gonzalez-Agosti, J.H. Hartwig and V. Ramesh

377–386

A surface-associated retinol- and fatty acid-binding protein (Gp-FAR-1) from the potato cyst nematode *Globodera pallida*: lipid binding activities, structural analysis and expression pattern

A. Prior, J.T. Jones, V.C. Blok, J. Beauchamp, L. McDermott, A. Cooper and M.W. Kennedy

387–394

Metabolism of S-nitrosoglutathione in intact mitochondria

M. Steffen, T.M. Sarkela, A.A. Gybina, T.W. Steele, N.J. Trasseth, D. Kuehl and C. Giullivi

395–402

Rat glutathione S-transferase M4-4: an isoenzyme with unique structural features including a redox-reactive cysteine-115 residue that forms mixed disulphides with glutathione

H. Cheng, T. Tchaikovskaya, Y.-S.L. Tu, J. Chapman, B. Qian, W.-M. Ching, M. Tien, J.D. Rowe, Y.V. Patskovsky, I. Listowsky and C.-P.D. Tu

403–414

An integrated study of threonine-pathway enzyme kinetics in *Escherichia coli*

C. Chassagnole, B. Raïs, E. Quentin, D.A. Fell and J.-P. Mazat

415–423

Threonine synthesis from aspartate in *Escherichia coli* cell-free extracts: pathway dynamics

B. Raïs, C. Chassagnole, T. Letellier, D.A. Fell and J.-P. Mazat

425–432

Control of the threonine-synthesis pathway in <i>Escherichia coli</i> : a theoretical and experimental approach	C. Chassagnole, D.A. Fell, B. Raïs, B. Kudla and J.-P. Mazat	<b>433–444</b>
Regulation by retinoic acid of acylation-stimulating protein and complement C3 in human adipocytes	T. Scantlebury, A.D. Sniderman and K. Cianflone	<b>445–452</b>
Thermostability of <i>endo</i> -1,4- $\beta$ -xyylanase II from <i>Trichoderma reesei</i> studied by electrospray ionization Fourier-transform ion cyclotron resonance MS, hydrogen/deuterium-exchange reactions and dynamic light scattering	J. Jänis, J. Rouvinen, M. Leisola, O. Turunen and P. Vainiotalo	<b>453–460</b>
Mapping the Zap-70 phosphorylation sites on LAT (linker for activation of T cells) required for recruitment and activation of signalling proteins in T cells	P.E. Paz, S. Wang, H. Clarke, X. Lu, D. Stokoe and A. Abo	<b>461–471</b>
Distinct roles for extracellular-signal-regulated protein kinase (ERK) mitogen-activated protein kinases and phosphatidylinositol 3-kinase in the regulation of Mcl-1 synthesis	K.M. Schubert and V. Duronio	<b>473–480</b>
Bile acid hydrophobicity is correlated with induction of apoptosis and/or growth arrest in HCT116 cells	A.A. Powell, J.M. LaRue, A.K. Batta and J.D. Martinez	<b>481–486</b>
Pressure- and heat-induced inactivation of butyrylcholinesterase: evidence for multiple intermediates and the remnant inactivation process	A. Weingand-Ziade, F. Ribes, F. Renault and P. Masson	<b>487–493</b>
Role for phosphatidylinositol in nuclear envelope formation	B. Larijani, T.M. Barona and D.L. Poccia	<b>495–501</b>
Novel barnacle underwater adhesive protein is a charged amino acid-rich protein constituted by a Cys-rich repetitive sequence	K. Kamino	<b>503–507</b>
Intracellular ascorbic acid enhances the DNA single-strand breakage and toxicity induced by peroxyntirite in U937 cells	A. Guidarelli, R. De Sanctis, B. Cellini, M. Fiorani, M. Dachà and O. Cantoni	<b>509–513</b>
Triacylglycerol-rich lipoproteins alter the secretion, and the cholesterol-effluxing function, of apolipoprotein E-containing lipoprotein particles from human (THP-1) macrophages	E.M. Lindholm, A.M. Palmer and A. Graham	<b>515–523</b>
Inhibitor of nuclear factor- $\kappa$ B induction by cAMP antagonizes interleukin-1-induced human macrophage-colony-stimulating-factor expression	P.J. Kamthong and M.-c. Wu	<b>525–530</b>
Rat embryo fibroblasts require both the cell-binding and the heparin-binding domains of fibronectin for survival	J. Jeong, I. Han, Y. Lim, J. Kim, I. Park, A. Woods, J.R. Couchman and E.-S. Oh	<b>531–537</b>
The NR1 subunit of the <i>N</i> -methyl-D-aspartate receptor can be efficiently expressed alone in the cell surface of mammalian cells and is required for the transport of the NR2A subunit	M. García-Gallo, J. Renart and M. Díaz-Guerra	<b>539–547</b>
Apoptosis induced by exposure to a low steady-state concentration of H <sub>2</sub> O <sub>2</sub> is a consequence of lysosomal rupture	F. Antunes, E. Cadenas and U.T. Brunk	<b>549–555</b>
Modifications of eukaryotic initiation factor 4F (eIF4F) in adult cardiocytes by adenoviral gene transfer: differential effects on eIF4F activity and total protein synthesis rates	A.N. Saghir, W.J. Tuxworth, Jr, C.H. Hagedorn and P.J. McDermott	<b>557–566</b>
Distribution of protein disulphide isomerase in rat liver mitochondria	M.P. Rigobello, A. Donella-Deana, L. Cesaro and A. Bindoli	<b>567–570</b>
Functional analysis of the human annexin A5 gene promoter: a downstream DNA element and an upstream long terminal repeat regulate transcription	M.-T. Carcedo, J.-M. Iglesias, P. Bances, R.O. Morgan and M.-P. Fernandez	<b>571–579</b>
Identification of a novel neuroligin in humans which binds to PSD-95 and has a widespread expression	M.F. Bolliger, K. Frei, K.H. Winterhalter and S.M. Gloor	<b>581–588</b>

---

Proline residues in two tightly coupled helices of the sulphate transporter, SHST1, are important for sulphate transport	M.C. Shelden, P. Loughlin, M.L. Tierney and S.M. Howitt	<b>589–594</b>
Characterization of the gene family encoding alternative oxidase from <i>Candida albicans</i>	W.-K. Huh and S.-O. Kang	<b>595–604</b>
Signal transducer gp130: biochemical characterization of the three membrane-proximal extracellular domains and evaluation of their oligomerization potential	S. Pflanz, T. Kernebeck, B. Giese, A. Herrmann, M. Pachta-Nick, J. Stahl, A. Wollmer, P.C. Heinrich, G. Müller-Newen and J. Grötzinger	<b>605–612</b>
Human NADPH-P450 oxidoreductase modulates the level of cytochrome P450 CYP2D6 holoprotein via haem oxygenase-dependent and -independent pathways	S. Ding, D. Yao, Y.Y. Deeni, B. Burchell, C.R. Wolf and T. Friedberg	<b>613–619</b>
Tributyltin interacts with mitochondria and induces cytochrome <i>c</i> release	A. Nishikimi, Y. Kira, E. Kasahara, E.F. Sato, T. Kanno, K. Utsumi and M. Inoue	<b>621–626</b>
Importance of the 3′ untranslated region of ornithine decarboxylase mRNA in the translational regulation of the enzyme	E. Lövkvist Wallström, K. Takao, A. Wendt, C. Vargiu, H. Yin and L. Persson	<b>627–634</b>
The activation function-1 of hepatocyte nuclear factor-4 is an acidic activator that mediates interactions through bulky hydrophobic residues	E. Kistanova, H. Dell, P. Tsantili, E. Falvey, C. Cladaras and M. Hadzopoulou-Cladaras	<b>635–642</b>
Intracellular energetic units in red muscle cells	V.A. Saks, T. Kaambre, P. Sikk, M. Eimre, E. Orlova, K. Paju, A. Piirsoo, F. Appaix, L. Kay, V. Regitz-Zagrosek, E. Fleck and E. Seppet	<b>643–657</b>
Emergence during development of the white-adipocyte cell phenotype is independent of the brown-adipocyte cell phenotype	K. Moulin, N. Truel, M. André, E. Arnauld, M. Nibbelink, B. Cousin, C. Dani, L. Pénicaud and L. Casteilla	<b>659–664</b>