

**Review**

- Neuroendocrine secretory protein 7B2: structure, expression and functions** **329–342**  
by M. Mbikay, N.G. Seidah and M. Chrétien

**Research Papers**

- Variation in aspects of cysteine proteinase catalytic mechanism deduced by spectroscopic observation of dithioester intermediates, kinetic analysis and molecular dynamics simulations **343–352**  
J.D. Reid, S. Hussain, S.K. Sreedharan, T.S.F. Bailey, S. Pinitglang, E.W. Thomas, C.S. Verma and K. Brocklehurst
- Metalloprotease-disintegrin ADAM 12 interacts with  $\alpha$ -actinin-1 **353–361**  
Y. Cao, Q. Kang and A. Zolkiewska
- Functional coupling between the caffeine/ryanodine-sensitive  $Ca^{2+}$  store and mitochondria in rat aortic smooth muscle cells **363–371**  
O. Vallot, L. Combettes and A.-M. Lompré
- Macrophage migration inhibitory factor of the parasitic nematode *Trichinella spiralis* **373–383**  
T.H.P. Tan, S.A.V. Edgerton, R. Kumari, M.S.B. McAlister, S.M. Rowe, S. Nagl, L.H. Pearl, M.E. Selkirk, A.E. Bianco, N.F. Totty, C. Engwerda, C.A. Gray and D.J. Meyer
- Site-directed removal of N-glycosylation sites in BST-1/CD157: effects on molecular and functional heterogeneity **385–392**  
S. Yamamoto-Katayama, A. Sato, M. Ariyoshi, M. Suyama, K. Ishihara, T. Hirano, H. Nakamura, K. Morikawa and H. Jingami
- Identification and characterization of molecular interactions between glucose-regulated proteins (GRPs) mortalin/GRP75/peptide-binding protein 74 (PBP74) and GRP94 **393–398**  
S. Takano, R. Wadhwa, Y. Mitsui and S.C. Kaul
- Characterization of active-site residues in diadenosine tetraphosphate hydrolase from *Lupinus angustifolius* **399–405**  
D. Maksel, P.R. Gooley, J.D. Swarbrick, A. Guranowski, C. Gange, G.M. Blackburn and K.R. Gayler
- Temporal relationships between ceramide production, caspase activation and mitochondrial dysfunction in cell lines with varying sensitivity to anti-Fas-induced apoptosis **407–416**  
C. Rodriguez-Lafresse, G. Alphonse, P. Broquet, M.-T. Aloy, P. Louisot and R. Rousson
- Phenotype–genotype relationships in peroxisome biogenesis disorders of *PEX1*-defective complementation group 1 are defined by Pex1p–Pex6p interaction **417–426**  
S. Tamura, N. Matsumoto, A. Imamura, N. Shimozawa, Y. Suzuki, N. Kondo and Y. Fujiki
- Targeting mutants of PTEN reveal distinct subsets of tumour suppressor functions **427–435**  
N.R. Leslie, D. Bennett, A. Gray, I. Pass, K. Hoang-Xuan and C.P. Downes
- Phosphorylation of human plasma  $\alpha_2$ -Heremans–Schmid glycoprotein (human fetuin) *in vivo* **437–445**  
Å.C. Haglund, B. Ek and P. Ek
- Cloning and characterization of the 5'-flanking region of the rat glutamate-cysteine ligase catalytic subunit **447–455**  
H. Yang, J. Wang, Z.-Z. Huang, X. Ou and S.C. Lu
- Platelet-activating factor acetylhydrolase and transacetylase activities in human plasma low-density lipoprotein **457–464**  
D.C. Tsoukatos, T.A. Liapikos, A.D. Tselepis, M.J. Chapman and E. Ninio
- FK506-binding protein of the hyperthermophilic archaeum, *Thermococcus* sp. KS-1, a cold-shock-inducible peptidyl-prolyl *cis*–*trans* isomerase with activities to trap and refold denatured proteins **465–471**  
A. Ideno, T. Yoshida, T. Iida, M. Furutani and T. Maruyama

Amphibian DNases I are characterized by a C-terminal end with a unique, cysteine-rich stretch and by the insertion of a serine residue into the Ca <sup>2+</sup> -binding site	H. Takeshita, T. Yasuda, R. Iida, T. Nakajima, S. Mori, K. Mogi, Y. Kaneko and K. Kishi	<b>473–480</b>
Comparison of the effects of dietary <i>n</i> –3 and <i>n</i> –6 polyunsaturated fatty acids on very-low-density lipoprotein secretion when delivered to hepatocytes in chylomicron remnants	X. Zheng, M. Avella and K.M. Botham	<b>481–487</b>
Polyamines regulate gap junction communication in connexin 43-expressing cells	L. Shore, P. McLean, S.K. Gilmour, M.B. Hodgins and M.E. Finbow	<b>489–495</b>
Wortmannin-sensitive trafficking steps in the endocytic pathway in rat liver endothelial cells	R. Kjekén, S.A. Mousavi, A. Brech, G. Griffiths and T. Berg	<b>497–503</b>
Determination of the chemical structure of the capsular polysaccharide of strain B33, a fast-growing soya bean-nodulating bacterium isolated from an arid region of China	M.A. Rodríguez-Carvajal, P. Tejero-Mateo, J.L. Espartero, J.E. Ruiz-Sainz, A.M. Buendía-Clavería, F.J. Ollero, S.S. Yang and A.M. Gil-Serrano	<b>505–511</b>
Involvement of the chicken liver 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase sequence His <sup>444</sup> -Arg-Glu-Arg in modulation of the bisphosphatase activity by its kinase domain	Z. Zhu, S. Ling, Q.-H. Yang and L. Li	<b>513–520</b>
Identification of a peroxisome-proliferator-activated-receptor response element in the apolipoprotein E gene control region	R. Galetto, M. Albajar, J.I. Polanco, M.M. Zakin and J.C. Rodríguez-Rey	<b>521–527</b>
Green fluorescent protein (GFP) tagged to the cytoplasmic tail of $\alpha$ IIb or $\beta$ 3 allows the expression of a fully functional integrin $\alpha$ IIb $\beta$ 3: effect of $\beta$ 3 <sub>GFP</sub> on $\alpha$ IIb $\beta$ 3 ligand binding	S. Plançon, M.-C. Morel-Kopp, E. Schaffner-Reckinger, P. Chen and N. Kieffer	<b>529–536</b>
Protein-binding elements in the proximal parotid secretory protein gene enhancer essential for salivary-gland-specific expression	P. Svendsen, K. Kristiansen and J.P. Hjorth	<b>537–544</b>
Functional and conformational properties of the exclusive C-domain from the <i>Arabidopsis</i> copper chaperone (CCH)	H. Mira, M. Vilar, E. Pérez-Payá and L. Peñarribia	<b>545–549</b>
Human cyritestin genes ( <i>CYRN1</i> and <i>CYRN2</i> ) are non-functional	P. Grzmil, Y. Kim, R. Shamsadin, J. Neesen, I.M. Adham, U.A.O. Heinlein, U.J. Schwarzer and W. Engel	<b>551–556</b>
Glucose 6-phosphate transport in fibroblast microsomes from glycogen storage disease type 1b patients: evidence for multiple glucose 6-phosphate transport systems	R. Leuzzi, R. Fulceri, P. Marcolongo, G. Bánhegyi, E. Zammarchi, K. Stafford, A. Burchell and A. Benedetti	<b>557–562</b>
Autocatalytic tyrosine-phosphorylation of protein kinase CK2 $\alpha$ and $\alpha'$ subunits: implication of Tyr <sup>182</sup>	A. Donella-Deana, L. Cesaro, S. Sarno, A.M. Brunati, M. Ruzzene and L.A. Pinna	<b>563–567</b>
Extracellular processing of amphoterin generates a peptide active on erythroleukaemia cell differentiation	B. Sparatore, M. Patrone, M. Passalacqua, M. Pedrazzi, D. Gaggero, S. Pontremoli and E. Melloni	<b>569–574</b>
Inhibition of distant caspase homologues by natural caspase inhibitors	S.J. Snipas, H.R. Stennicke, S. Riedl, J. Potempa, J. Travis, A.J. Barrett and G.S. Salvesen	<b>575–580</b>
Sulphoacetaldehyde sulphydrylase (EC 4.4.1.12) from <i>Desulfonispora thiosulfatigenes</i> : purification, properties and primary sequence	K. Denger, J. Ruff, U. Rein and A.M. Cook	<b>581–586</b>
G-protein-coupled-receptor activation of the smooth muscle calponin gene	N.O. Dulin, S.N. Orlov, C.M. Kitchen, T.A. Voyno-Yasenetskaya and J.M. Miano	<b>587–592</b>

