

Biochemical Journal**Volume 358, part 3****15 September 2001**

First published on the Internet
10 September 2001
<http://www.BiochemJ.org>

Subscribing organizations are encouraged to copy and distribute this table of contents for non-commercial purposes

Research Papers

Carbohydrate specificity of a galectin from chicken liver (CG-16)	A.M. Wu, J.H. Wu, M.-S. Tsai, H. Kaltner and H.-J. Gabius	529–538
Human glycosaminoglycan glucuronyltransferase I gene and a related processed pseudogene: genomic structure, chromosomal mapping and characterization	H. Kitagawa, M. Taoka, Y. Tone and K. Sugahara	539–546
Flavonoids protect neurons from oxidized low-density-lipoprotein-induced apoptosis involving c-Jun N-terminal kinase (JNK), c-Jun and caspase-3	H. Schroeter, J.P.E. Spencer, C. Rice-Evans and R.J. Williams	547–557
All three splice variants of the human sarco/endoplasmic reticulum Ca^{2+} -ATPase 3 gene are translated to proteins: a study of their co-expression in platelets and lymphoid cells	T. Kovács, F. Felföldi, B. Papp, K. Pászty, R. Bredoux, Á. Enyedi and J. Enouf	559–568
Lactate supply as a determinant of the distribution of intracellular pH within the hepatic lobule	S.P. Burns, H.C. Murphy, R.A. Iles and R.D. Cohen	569–571
A normalized plot as a novel and time-saving tool in complex enzyme kinetic analysis	I.G. Bravo, F. Bustos, D. De Arriaga, M.A. Ferrero, L.B. Rodríguez-Aparicio, H. Martínez-Blanco and Á. Reglero	573–583
Kinetic properties of the acylneuraminate cytidylyltransferase from <i>Pasteurella haemolytica</i> A2 ¹	I.G. Bravo, S. Barrallo, M.A. Ferrero, L.B. Rodríguez-Aparicio, H. Martínez-Blanco and Á. Reglero	585–598
<i>Pseudomonas cellulosa</i> expresses a single membrane-bound glycoside hydrolase family 51 arabinofuranosidase	M.-H. Beylot, K. Emami, V.A. McKie, H.J. Gilbert and G. Pell	599–605
The <i>Pseudomonas cellulosa</i> glycoside hydrolase family 51 arabinofuranosidase exhibits wide substrate specificity	M.-H. Beylot, V.A. McKie, A.G.J. Voragen, C.H.L. Doeswijk-Voragen and H.J. Gilbert	607–614
Analysis of aggrecan in human knee cartilage and synovial fluid indicates that aggrecanase (ADAMTS) activity is responsible for the catabolic turnover and loss of whole aggrecan whereas other protease activity is required for C-terminal processing <i>in vivo</i>	J.D. Sandy and C. Verscharen	615–626
Dimethyl sulphoxide enhances the effects of P_i in myofibrils and inhibits the activity of rabbit skeletal muscle contractile proteins	A.C. Mariano, G.M.C. Alexandre, L.C. Silva, A. Romeiro, L.C. Cameron, Y. Chen, P.B. Chase and M.M. Sorenson	627–636
Pulmonary lipid phosphate phosphohydrolase in plasma membrane signalling platforms	M. Nanjundan and F. Possmayer	637–646
Cloning and characterization of the pseudonajatoxin b precursor	N. Gong, A. Armugam, P. Mirtschin and K. Jeyaseelan	647–656
Sd ^a -antigen-like structures carried on core 3 are prominent features of glycans from the mucin of normal human descending colon	C. Capon, E. Maes, J.-C. Michalski, H. Leffler and Y.S. Kim	657–664
Fatty acid and amino acid modulation of glucose cycling in isolated rat hepatocytes	L.A. Gustafson, M. Neeft, D.-J. Reijngoud, F. Kuipers, H.P. Sauerwein, J.A. Romijn, A.W. Herling, H.-J. Burger and A.J. Meijer	665–671
Urokinase-catalysed cleavage of the urokinase receptor requires an intact glycolipid anchor	G. Hoyer-Hansen, U. Pessara, A. Holm, J. Pass, U. Weidle, K. Danø and N. Behrendt	673–679

Parameters affecting <i>in vitro</i> oxidation/folding of maurotoxin, a four-disulphide-bridged scorpion toxin	E. Di Luccio, D.-O. Azulay, I. Regaya, Z. Fajloun, G. Sandoz, P. Mansuelle, R. Kharrat, M. Fathallah, L. Carrega, E. Estève, H. Rochat, M. De Waard and J.-M. Sabatier	681–692
Molecular events involved in up-regulation human Na ⁺ -independent neutral amino acid transporter LAT1 during T-cell activation	T. Nii, H. Segawa, Y. Taketani, Y. Tani, M. Ohkido, S. Kishida, M. Ito, H. Endou, Y. Kanai, E. Takeda and K.-i. Miyamoto	693–704
Targeting of the transcription factor Max during apoptosis: phosphorylation-regulated cleavage by caspase-5 at an unusual glutamic acid residue in position P1	A. Krippner-Heidenreich, R.V. Talanian, R. Sekul, R. Kraft, H. Thole, H. Ottleben and B. Lüscher	705–715
A diacylglycerol-activated Ca ²⁺ channel in PC12 cells (an adrenal chromaffin cell line) correlates with expression of the TRP-6 (transient receptor potential) protein	Y. Tesfai, H.M. Brereton and G.J. Barritt	717–726
Binding of the merlin-I product of the neurofibromatosis type 2 tumour suppressor gene to a novel site in β -fodrin is regulated by association between merlin domains	G.W. Neill and M.R. Crompton	727–735
Multimerization of monocyte chemoattractant protein-1 is not required for glycosaminoglycan-dependent transendothelial chemotaxis	S. Ali, A.C.V. Palmer, S.J. Fritchley, Y. Maley and J.A. Kirby	737–745
Molecular cloning, gene organization and expression of rainbow trout (<i>Oncorhynchus mykiss</i>) inducible nitric oxide synthase (iNOS) gene	T. Wang, M. Ward, P. Grabowski and C.J. Secombes	747–755
Four <i>Trypanosoma brucei</i> fatty acyl-CoA synthetases: fatty acid specificity of the recombinant proteins	D.W. Jiang and P.T. Englund	757–761
Human mucin gene <i>MUC5AC</i> : organization of its 5'-region and central repetitive region	F. Escande, J.-P. Aubert, N. Porchet and M.-P. Buisine	763–772
Lipolytic activity of ricin from <i>Ricinus sanguineus</i> and <i>Ricinus communis</i> on neutral lipids	S. Lombard, M.E. Helmy and G. Piéroni	773–781
Phosphorylation of the leucocyte NADPH oxidase subunit p47 ^{phox} by casein kinase 2: conformation-dependent phosphorylation and modulation of oxidase activity	H.-S. Park, S.M. Lee, J.H. Lee, Y.-S. Kim, Y.-S. Bae and J.-W. Park	783–790
Corrections		
Macrophage migration inhibitory factor of the parasitic nematode <i>Trichinella spiralis</i> by T.H.P. Tan, S.A.V. Edgerton, R. Kumari, M.S.B. McAlister, S.M. Rowe, S. Nagl, L.H. Pearl, M.E. Selkirk, A.E. Bianco, N.F. Totty, C. Engwerda, C.A. Gray and D.J. Meyer (volume 356, pages 373–383, 2001)		791
Activity and genomic organization of human glucose transporter 9 (GLUT9), a novel member of the family of sugar-transport facilitators predominantly expressed in brain and leucocytes by H. Doege, A. Bocianski, H.-G. Joost and A. Schürmann (volume 350, pages 771–776, 2001)		791–792
Reactive oxygen species involved in trichosanthin-induced apoptosis of human choriocarcinoma cells by C.-y. Zhang, Y.-x. Gong, H. Ma, C.-c. An and D.-y. Chen (volume 355, pages 653–661, 2001)		792