

**Research Communication**

- Congenital disorders of glycosylation type I leads to altered processing of N-linked glycans, as well as underglycosylation  
P. Mills, K. Mills, P. Clayton, A. Johnson, D. Whitehouse and B. Winchester **249–254**

**Research Papers**

- Coactosin-like protein, a human F-actin-binding protein: critical role of lysine-75  
P. Provost, J. Doucet, A. Stock, G. Gerisch, B. Samuelsson and O. Rådmark **255–263**
- Conformation of heparin pentasaccharide bound to antithrombin III  
M. Hricovíni, M. Guerrini, A. Bisio, G. Torri, M. Petitou and B. Casu **265–272**
- Promoter I of the ovine acetyl-CoA carboxylase- $\alpha$  gene: an E-box motif at -114 in the proximal promoter binds upstream stimulatory factor (USF)-1 and USF-2 and acts as an insulin-response sequence in differentiating adipocytes  
M.T. Travers, A.J. Vallance, H.T. Gourlay, C.A. Gill, I. Klein, C.B.K. Bottema and M.C. Barber **273–284**
- RhoGDI-binding-defective mutant of Cdc42Hs targets to membranes and activates filopodia formation but does not cycle with the cytosol of mammalian cells  
R.M. Gibson and A.L. Wilson-Delfosse **285–294**
- Identification of a novel class of insect glutathione S-transferases involved in resistance to DDT in the malaria vector *Anopheles gambiae*  
H. Ranson, L. Rossiter, F. Ortelli, B. Jensen, X. Wang, C.W. Roth, F.H. Collins and J. Hemingway **295–304**
- Metabolism of exogenous *sn*-1-alkyl-*sn*-2-lyso-glucosaminyl-phosphatidylinositol in HeLa D cells: accumulation of glucosaminyl(acyl)phosphatidylinositol in a metabolically inert compartment  
A. Wongkajornsilp, D. Sevliver and T.L. Rosenberry **305–313**
- Effects of environment on flavin reactivity in morphinone reductase: analysis of enzymes displaying differential charge near the N-1 atom and C-2 carbonyl region of the active-site flavin  
D.H. Craig, T. Barna, P.C.E. Moody, N.C. Bruce, S.K. Chapman, A.W. Munro and N.S. Scrutton **315–323**
- Calmodulin inhibitors trigger the proteolytic processing of membrane type-1 matrix metalloproteinase, but not its shedding in glioblastoma cells  
B. Annabi, A. Pilorget, N. Bousquet-Gagnon, D. Gingras and R. Béliveau **325–333**
- Sphingosine-induced apoptosis is dependent on lysosomal proteases  
K. Kågedal, M. Zhad, I. Svensson and U.T. Brunk **335–343**
- Survival-factor-induced phosphorylation of Bad results in its dissociation from Bcl-x<sub>L</sub> but not Bcl-2  
I. Hirai and H.-G. Wang **345–352**
- Oxidative dimerization in metallothionein is a result of intermolecular disulphide bonds between cysteines in the  $\alpha$ -domain  
K. Zangger, G. Shen, G. Öz, J.D. Otvos and I.M. Armitage **353–360**
- S-phase-specific expression of the *Mad3* gene in proliferating and differentiating cells  
E.J. Fox and S.C. Wright **361–367**
- Cloning and expression of a single-chain catalytic antibody that acts as a glutathione peroxidase mimic with high catalytic efficiency  
X. Ren, S. Gao, D. You, H. Huang, Z. Liu, Y. Mu, J. Liu, Y. Zhang, G. Yan, G. Luo, T. Yang and J. Shen **369–374**
- Endothelin-1 induces phosphorylation of GATA-4 transcription factor in the HL-1 atrial-muscle cell line  
K. Kitta, S.A. Clément, J. Remeika, J.B. Blumberg and Y.J. Suzuki **375–380**

Identification of the catalytic nucleophile of the Family 31 $\alpha$ -glucosidase from <i>Aspergillus niger</i> via trapping of a 5-fluoroglycosyl-enzyme intermediate	S.S. Lee, S. He and S.G. Withers	<b>381–386</b>
Cloning and characterization of the mouse polyamine-modulated factor-1 (mPMF-1) gene: an alternatively spliced homologue of the human transcription factor	Y. Wang, W. Devereux, P.M. Woster and R.A. Casero, Jr	<b>387–392</b>
Identification of a novel integral plasma membrane protein induced during adipocyte differentiation	T. Albrektsen, H.E. Richter, J.T. Clausen and J. Fleckner	<b>393–402</b>
Functional analysis of the interleukin-1-receptor-associated kinase (IRAK-1) in interleukin-1 $\beta$ -stimulated nuclear factor $\kappa$ B (NF- $\kappa$ B) pathway activation: IRAK-1 associates with the NF- $\kappa$ B essential modulator (NEMO) upon receptor stimulation	E.-L. Cooke, I.J. Uings, C.L. Xia, P. Woo and K.P. Ray	<b>403–410</b>
The result of equilibrium-constant calculations strongly depends on the evaluation method used and on the type of experimental errors	H. Fuchs and R. Gessner	<b>411–418</b>
Different combinations of the heat-shock cognate protein 70 (hsc70) C-terminal functional groups are utilized to interact with distinct tetratricopeptide repeat-containing proteins	S.-J. Wu, F.-H. Liu, S.-M. Hu and C. Wang	<b>419–426</b>
Study of the interaction of the medium chain $\mu$ 2 subunit of the clathrin-associated adapter protein complex 2 with cytotoxic T-lymphocyte antigen 4 and CD28	E.R. Follows, J.C. McPheat, C. Minshull, N.C. Moore, R.A. Pauptit, S. Rowsell, C.L. Stacey, J.J. Stanway, I.W.F. Taylor and W.M. Abbott	<b>427–434</b>
Functional properties of <i>Drosophila</i> inositol trisphosphate receptors	J.E. Swatton, S.A. Morris, F. Wissing and C.W. Taylor	<b>435–441</b>
Characterization of human glucose transporter (GLUT) 11 (encoded by <i>SLC2A11</i> ), a novel sugar-transport facilitator specifically expressed in heart and skeletal muscle	H. Doege, A. Bocianski, A. Scheepers, H. Axer, J. Eckel, H.-G. Joost and A. Schürmann	<b>443–449</b>
Pharmacological characterization of the putative cADP-ribose receptor	J.M. Thomas, R. Masgrau, G.C. Churchill and A. Galione	<b>451–457</b>
Critical roles for the serine 20, but not the serine 15, phosphorylation site and for the polyproline domain in regulating p53 turnover	N. Dumaz, D.M. Milne, L.J. Jardine and D.W. Meek	<b>459–464</b>