

**Research Communication**

- Isoform-specific knockdown and expression of adaptor protein ShcA using small interfering RNA M. Kisielow, S. Kleiner, M. Nagasawa, A. Faisal and Y. Nagamine 1–5

**Research Papers**

- Structure determination of the exopolysaccharide produced by *Lactobacillus rhamnosus* strains RW-9595M and R M.-R. Van Calsteren, C. Pau-Roblot, A. Bégin and D. Roy 7–17
- Platelet-derived growth factor-BB-mediated glycosaminoglycan synthesis is transduced through Akt N.J. Cartel, J. Wang and M. Post 19–28
- Cyclophilin-A is involved in excitotoxin-induced caspase activation in rat neuronal B50 cells M. Capano, S. Virji and M. Crompton 29–36
- Human insulin-like growth factor II leader 2 mediates internal initiation of translation S.K. Pedersen, J. Christiansen, T.v.O. Hansen, M.R. Larsen, and F.C. Nielsen 37–44
- Ceramides increase the activity of the secretory phospholipase A<sub>2</sub> and alter its fatty acid specificity K.S. Koumanov, A.B. Momchilova, P.J. Quinn and C. Wolf 45–51
- An alternative model of H ferritin promoter transactivation by c-Jun M.C. Faniello, G. Chirico, B. Quaresima, G. Cuda, G. Allevalo, M.A. Bevilacqua, F. Baudi, V. Colantuoni, F. Cimino, S. Venuta, V.E. Avvedimento and F. Costanzo 53–58
- Evidence that zymogen granules do not function as an intracellular Ca<sup>2+</sup> store for the generation of the Ca<sup>2+</sup> signal in rat parotid acinar cells A. Nezu, A. Tanimura, T. Morita, K. Irie, T. Yajima and Y. Tojyo 59–66
- Investigation of the role of Endo180/urokinase-type plasminogen activator receptor-associated protein as a collagenase 3 (matrix metalloproteinase 13) receptor L. Bailey, D. Wienke, M. Howard, V. Knäuper, C.M. Isacke and G. Murphy 67–72
- Monitoring of exocytosis and endocytosis of insulin secretory granules in the pancreatic  $\beta$ -cell line MIN6 using pH-sensitive green fluorescent protein (pHluorin) and confocal laser microscopy M. Ohara-Imaizumi, Y. Nakamichi, T. Tanaka, H. Katsuta, H. Ishida and S. Nagamatsu 73–80
- Expression, purification and characterization of human glutamate dehydrogenase (GDH) allosteric regulatory mutations J. Fang, B.Y.L. Hsu, C.M. MacMullen, M. Poncz, T.J. Smith and C.A. Stanley 81–87
- Solute carrier 11a1 (*Slc11a1*; formerly *Nramp1*) regulates metabolism and release of iron acquired by phagocytic, but not transferrin-receptor-mediated, iron uptake V. Mulero, S. Searle, J.M. Blackwell and J.H. Brock 89–94
- Involvement of protein kinase D in Fc $\gamma$ -receptor activation of the NADPH oxidase in neutrophils J.K. Davidson-Moncada, G. Lopez-Lluch, A.W. Segal and L.V. Dekker 95–103
- Major outer membrane proteins and proteolytic processing of RgpA and Kgp of *Porphyromonas gingivalis* W50 P.D. Veith, G.H. Talbo, N. Slakeski, S.G. Dashper, C. Moore, R.A. Paolini and E.C. Reynolds 105–115
- Maintenance of the filamentous actin cytoskeleton is necessary for the activation of store-operated Ca<sup>2+</sup> channels, but not other types of plasma-membrane Ca<sup>2+</sup> channels, in rat hepatocytes Y.-J. Wang, R.B. Gregory and G.J. Barritt 117–126
- Protein kinase A enhances, whereas glycogen synthase kinase-3 $\beta$  inhibits, the activity of the exon 2-encoded transactivator domain of heterogeneous nuclear ribonucleoprotein D in a hierarchical fashion M. Tolnay, Y.-T. Juang and G.C. Tsokos 127–136



|  |  |                |
|--|--|----------------|
| Glycosylation and epitope mapping of the 5T4 glycoprotein oncofoetal antigen   | D.M. Shaw, A.M. Woods, K.A. Myers, C. Westwater, V. Rahi-Saund, M.J. Davies, D.V. Renouf, E.F. Hounsell and P.L. Stern | <b>137–145</b> |
| Comparative characterization of two DEAD-box RNA helicases in superfamily II: human translation-initiation factor 4A and hepatitis C virus non-structural protein 3 (NS3) helicase | M.X. Du, R.B. Johnson, X.-L. Sun, K.A. Staschke, J. Colacino and Q.M. Wang   | <b>147–155</b> |
| Nuclear receptor corepressor-dependent repression of peroxisome-proliferator-activated receptor $\delta$ -mediated transactivation   | A.-M. Krogsdam, C.A.F. Nielsen, S. Neve, D. Holst, T. Helledie, B. Thomsen, C. Bendixen, S. Mandrup and K. Kristiansen | <b>157–165</b> |
| Characterization of the role of the AMP-activated protein kinase in the stimulation of glucose transport in skeletal muscle cells  | L.G.D. Fryer, F. Fougelle, K. Barnes, S.A. Baldwin, A. Woods and D. Carling  | <b>167–174</b> |
| Opposing roles of serine/threonine kinases MEKK1 and LOK in regulating the CD28 responsive element in T-cells  | L. Tao, S. Wadsworth, J. Mercer, C. Mueller, K. Lynn, J. Siekierka and A. August                                       | <b>175–182</b> |
| <i>De novo</i> -synthesized ceramide is involved in cannabinoid-induced apoptosis  | T. Gómez del Pulgar, G. Velasco, C. Sánchez, A. Haro and M. Guzmán   | <b>183–188</b> |
| Glutamic acid-65 is an essential residue for catalysis in <i>Proteus mirabilis</i> glutathione S-transferase B1-1  | N. Allocati, M. Masulli, E. Casalone, S. Santucci, B. Favaloro, M.W. Parker and C. Di Ilio                             | <b>189–193</b> |
| Interaction of the transforming acidic coiled-coil 1 (TACC1) protein with ch-TOG and GAS41/NuBI1 suggests multiple TACC1-containing protein complexes in human cells               | B. Lauffart, S.J. Howell, J.E. Tasch, J.K. Cowell and I.H. Still   | <b>195–200</b> |