

Research Papers

| | | |
|--|---|---------|
| Assay of advanced glycation endproducts (AGEs): surveying AGEs by chromatographic assay with derivatization by 6-aminoquinolyl- <i>N</i> -hydroxysuccinimidyl-carbamate and application to <i>N</i> _ε -carboxymethyl-lysine- and <i>N</i> _ε -(1-carboxyethyl)lysine-modified albumin | N. Ahmed, O.K. Argirov, H.S. Minhas, C.A.A. Cordeiro and P.J. Thornalley | 1–14 |
| Chromatographic assay of glycation adducts in human serum albumin glycated <i>in vitro</i> by derivatization with 6-aminoquinolyl- <i>N</i> -hydroxysuccinimidyl-carbamate and intrinsic fluorescence | N. Ahmed and P.J. Thornalley | 15–24 |
| Characteristics of physiological inducers of the ethanol utilization (<i>alc</i>) pathway in <i>Aspergillus nidulans</i> | M. Flipphi, J. Kocalkowska and B. Felenbok | 25–31 |
| Somatostatin-induced paradoxical increase in intracellular Ca ²⁺ concentration and insulin release in the presence of arginine vasopressin in clonal HIT-T15 β-cells | H. Cheng, S. Yibchok-Anun, S.-C. Park and W.H. Hsu | 33–39 |
| Protein kinase CK2 inhibitor 4,5,6,7-tetrabromobenzotriazole (TBB) induces apoptosis and caspase-dependent degradation of haematopoietic lineage cell-specific protein 1 (HS1) in Jurkat cells | M. Ruzzene, D. Penzo and L.A. Pinna | 41–47 |
| The same rat Δ6-desaturase not only acts on 18- but also on 24-carbon fatty acids in very-long-chain polyunsaturated fatty acid biosynthesis | S. D'Andrea, H. Guillou, S. Jan, D. Catheline, J.-N. Thibault, M. Bouriel, V. Rioux and P. Legrand | 49–55 |
| μ-1,2-Peroxybridged di-iron(III) dimer formation in human H-chain ferritin | F. Bou-Abdallah, G.C. Papaefthymiou, D.M. Scheswohl, S.D. Stanga, P. Arosio and N.D. Chasteen | 57–63 |
| Primary arrest of circulating platelets on collagen involves phosphorylation of Syk, cortactin and focal adhesion kinase: studies under flow conditions | G. Arderiu, M. Díaz-Ricart, B. Buckley, G. Escolar and A. Ordinas | 65–71 |
| A novel pathway for adrenergic stimulation of cAMP-response-element-binding protein (CREB) phosphorylation: mediation via α ₁ -adrenoceptors and protein kinase C activation | H. Thonberg, J.M. Fredriksson, J. Nedergaard and B. Cannon | 73–79 |
| Isolation and characterization of a polymerized prion protein | B.-Y. Lu and J.-Y. Chang | 81–87 |
| Identification of an initiator-like element essential for the expression of the tissue inhibitor of metalloproteinases-4 (<i>Timp-4</i>) gene | D.A. Young, B.W. Phillips, C. Lundy, R.K. Nuttall, A. Hogan, G.A. Schultz, K.J. Leco, I.M. Clark and D.R. Edwards | 89–99 |
| L-Lactate transport into rat heart mitochondria and reconstruction of the L-lactate/pyruvate shuttle | D. Valenti, L. de Bari, A. Atlante and S. Passarella | 101–104 |
| Constitutive expression of hZnT4 zinc transporter in human breast epithelial cells | A.A. Michalczyk, J. Allen, R.C. Blomeley and M.L. Ackland | 105–113 |
| Elevation of hepatic sulphotransferase activities in mice with resistance to cystic fibrosis | J.L. Falany, H. Greer, T. Kovacs, E.J. Sorscher and C.N. Falany | 115–120 |
| Plant-derived phenolic compounds prevent the DNA single-strand breakage and cytotoxicity induced by <i>tert</i> -butylhydroperoxide via an iron-chelating mechanism | P. Sestili, G. Diamantini, A. Bedini, L. Cerioni, I. Tommasini, G. Tarzia and O. Cantoni | 121–128 |

| | | |
|--|--|----------------|
| Isoaspartyl dipeptidase activity of plant-type asparaginases | M. Hejazi, K. Piotukh, J. Mattow, R. Deutzmann, R. Volkmer-Engert and W. Lockau | 129–136 |
| Only the large soluble form of preadipocyte factor-1 (Pref-1), but not the small soluble and membrane forms, inhibits adipocyte differentiation: role of alternative splicing | B. Mei, L. Zhao, L. Chen and H.S. Sul | 137–144 |
| Characterization of a murine gene encoding a developmentally regulated cytoplasmic dual-specificity mitogen-activated protein kinase phosphatase | R.J. Dickinson, D.J. Williams, D.N. Slack, J. Williamson, O.-M. Seternes and S.M. Keyse | 145–155 |
| Novel disease-causing mutations in the dihydropyrimidine dehydrogenase gene interpreted by analysis of the three-dimensional protein structure | A.B.P. van Kuilenburg, D. Dobritzsch, R. Meinsma, J. Haasjes, H.R. Waterham, M.J.M. Nowaczyk, G.D. Maropoulos, G. Hein, H. Kalhoff, J.M. Kirk, H. Baaske, A. Aukett, J.A. Duley, K.P. Ward, Y. Lindqvist and A.H. van Gennip | 157–163 |
| Crystal structure of human dehydroepiandrosterone sulphotransferase in complex with substrate | P.H. Rehse, M. Zhou and S.-X. Lin | 165–171 |
| Structural basis for the unusual carbohydrate-binding specificity of jacalin towards galactose and mannose | Y. Bourne, C.H. Astoul, V. Zamboni, W.J. Peumans, L. Menu-Bouaouiche, E.J.M. Van Damme, A. Barre and P. Rougé | 173–180 |
| The 45 kDa collagen-binding fragment of fibronectin induces matrix metalloproteinase-13 synthesis by chondrocytes and aggrecan degradation by aggrecanases | H. Stanton, L. Ung and A.J. Fosang | 181–190 |
| Gastric MUC5AC and MUC6 are large oligomeric mucins that differ in size, glycosylation and tissue distribution | H. Nordman, J.R. Davies, G. Lindell, C. de Bolós, F. Real and I. Carlstedt | 191–200 |
| Karyopherin $\alpha 2$: a control step of glucose-sensitive gene expression in hepatic cells | G. Guillemain, M.J. Muñoz-Alonso, A. Cassany, M. Loizeau, A.-M. Fausat, A.-F. Burnol and A. Leturque | 201–209 |
| Nitric oxide inhibits the shedding of the glycosylphosphatidylinositol-anchored dipeptidase from porcine renal proximal tubules | S.W. Park, H.J. Yoon, H.B. Lee, N.M. Hooper and H.S. Park | 211–218 |
| Adenovirus-mediated overexpression of liver carnitine palmitoyltransferase I in INS1E cells: effects on cell metabolism and insulin secretion | B. Rubí, P.A. Antinozzi, L. Herrero, H. Ishihara, G. Asins, D. Serra, C.B. Wollheim, P. Maechler and F.G. Hegardt | 219–226 |
| Engineering N-terminal domain of tissue inhibitor of metalloproteinase (TIMP)-3 to be a better inhibitor against tumour necrosis factor- α -converting enzyme | M.-H. Lee, V. Verma, K. Maskos, D. Nath, V. Knäuper, P. Dodds, A. Amour and G. Murphy | 227–234 |
| Ca ²⁺ uptake by the endoplasmic reticulum Ca ²⁺ -ATPase in rat microvascular endothelial cells | F. Moccia, R. Berra-Romani, S. Baruffi, S. Spaggiari, S. Signorelli, L. Castelli, J. Magistretti, V. Taglietti and F. Tanzi | 235–244 |
| Diacylglycerol activates the influx of extracellular cations in T-lymphocytes independently of intracellular calcium-store depletion and possibly involving endogenous <i>TRP6</i> gene products | A. Gamberucci, E. Giurisato, P. Pizzo, M. Tassi, R. Giunti, D.P. McIntosh and A. Benedetti | 245–254 |
| Cloning and characterization of hELD/OSA1, a novel BRG1 interacting protein | A.F.L. Hurlstone, I.A. Olave, N. Barker, M. van Noort and H. Clevers | 255–264 |
| Characterization of a novel intracellular heparanase that has a FERM domain | K.J. Bame, I. Venkatesan, J. Dehdashti, J. McFarlane and R. Burfeind | 265–274 |
| Turnover of matrix proteins in mammalian mitochondria | W. Huth, S. Rolle and I. Wunderlich | 275–284 |
| Use of a purified and functional recombinant calcium-channel β_4 subunit in surface-plasmon resonance studies | S. Geib, G. Sandoz, K. Mabrouk, A. Matavel, P. Marchot, T. Hoshi, M. Villaz, M. Ronjat, R. Miquelis, C. Lévêque and M. de Waard | 285–292 |
| Role of the self-association of β subunits in the oligomeric structure of Na ⁺ /K ⁺ -ATPase | A.V. Ivanov, N.N. Modyanov and A. Askari | 293–299 |
| High cellular accumulation of sulphoraphane, a dietary anticarcinogen, is followed by rapid transporter-mediated export as a glutathione conjugate | Y. Zhang and E.C. Callaway | 301–307 |

| | | |
|---|--|----------------|
| Contribution by different fuels and metabolic pathways to the total ATP turnover of proliferating MCF-7 breast cancer cells | M. Guppy, P. Leedman, X. Zu and V. Russell | 309–315 |
| Cardiolipin prevents rate-dependent uncoupling and provides osmotic stability in yeast mitochondria | V. Koshkin and M.L. Greenberg | 317–322 |
| Characteristics and composition of the vitamin K-dependent γ -glutamyl carboxylase-binding domain on osteocalcin | R.J.T.J. Houben, D.T.S. Rijkers, T.B. Stanley, F. Acher, R. Azerad, S.-M. Käkönen, C. Vermeer and B.A.M. Soute | 323–328 |