

### Online Commentary

New biochemical approaches towards understanding the Parkinson's disease-associated kinase, LRRK2

*Published on the Internet 23 October 2009,*

*doi:10.1042/BJ20091540*

(see accompanying paper, pp. 47–60)

G.-Y. Liou and K. A. Gallo

e1–e3

### Accelerated Publication

Ein2, the central regulator of ethylene signalling, is localized at the ER membrane where it interacts with the ethylene receptor ETR1

*Published as BJ Immediate Publication 21 September 2009,*

*doi:10.1042/BJ20091102*

M.M.A. Bisson, A. Bleckmann, S. Allekotte and G. Groth

1–6



### Research Papers

Existence of exocytotic hemifusion intermediates with a lifetime of up to seconds in type II pneumocytes

*Published as BJ Immediate Publication 27 August 2009,*

*doi:10.1042/BJ20091094*

P. Miklavc, S. Albrecht, O.H. Wittekindt, P. Schullian, T. Haller and P. Dietl

7–14



A survey of proteins encoded by non-synonymous single nucleotide polymorphisms reveals a significant fraction with altered stability and activity  
*Published as BJ Immediate Publication 25 August 2009,*  
*doi:10.1042/BJ20090723*

A. Allali-Hassani, G.A. Wasney, I. Chau, B.S. Hong, G. Senisterra, P. Loppnau, Z. Shi, J. Moult, A.M. Edwards, C.H. Arrowsmith, H.W. Park, M. Schapira and M. Vedadi

15–26



Regulation of histidine-rich glycoprotein (HRG) function via plasmin-mediated proteolytic cleavage  
*Published as BJ Immediate Publication 27 August 2009,*  
*doi:10.1042/BJ20090794*

I.K.H. Poon, A.-K. Olsson, M.D. Hulett and C.R. Parish

27–37



Human and mouse trace amine-associated receptor 1 have distinct pharmacology towards endogenous monoamines and imidazoline receptor ligands  
*Published as BJ Immediate Publication 2 September 2009,*  
*doi:10.1042/BJ20090998*

L.A. Hu, T. Zhou, J. Ahn, S. Wang, J. Zhou, Y. Hu and Q. Liu

39–45

Substrate specificity and inhibitors of LRRK2, a protein kinase mutated in Parkinson's disease  
*Published as BJ Immediate Publication 9 September 2009,*  
*doi:10.1042/BJ20091035*

R.J. Nichols, N. Dzamko, J.E. Hutton, L.C. Cantley, M. Deak, J. Moran, P. Bamforth, A.D. Reith and D.R. Alessi

47–60



Conditions of endoplasmic reticulum stress stimulate lipid droplet formation in *Saccharomyces cerevisiae*  
*Published as BJ Immediate Publication 26 August 2009,*  
*doi:10.1042/BJ20090785*

W. Fei, H. Wang, X. Fu, C. Bielby and H. Yang

61–67



Design of proteolytically stable RI-anchoring disruptor peptidomimetics for *in vivo* studies of anchored type I protein kinase A-mediated signalling  
*Published as BJ Immediate Publication 28 August 2009,*  
*doi:10.1042/BJ20090933*

E.A. Torheim, E. Jarnæss, B. Lygren and K. Taskén

69–78

Active-site determinants of substrate recognition by the metalloproteinases TACE and ADAM10 <i>Published as BJ Immediate Publication 28 August 2009, doi:10.1042/BJ20090549</i>	C.I. Caescu, G.R. Jeschke and B.E. Turk	 online data	79–88
Analysis of the dual function of the ESCRT-III protein Snf7 in endocytic trafficking and in gene expression <i>Published as BJ Immediate Publication 2 September 2009, doi:10.1042/BJ20090957</i>	P. Weiss, S. Huppert and R. Kölling		89–97
Importance of the bioenergetic reserve capacity in response to cardiomyocyte stress induced by 4-hydroxynonenal <i>Published as BJ Immediate Publication 10 September 2009, doi:10.1042/BJ20090934</i>	B.G. Hill, B.P. Dranka, L. Zou, J.C. Chatham and V.M. Darley-Usmar		99–107
IRAK1-independent pathways required for the interleukin-1-stimulated activation of the Tpl2 catalytic subunit and its dissociation from ABIN2 <i>Published as BJ Immediate Publication 15 September 2009, doi:10.1042/BJ20091271</i>	H. Handoyo, M.J. Stafford, E. McManus, D. Baltzis, M. Peggie and P. Cohen	 online data	109–118
The human <i>urocortin 2</i> gene is regulated by hypoxia: identification of a hypoxia-responsive element in the 3'-flanking region <i>Published as BJ Immediate Publication 26 August 2009, doi:10.1042/BJ20090311</i>	K. Bühler, I. Plaisance, T. Dieterle and M. Brink	 online data	119–127
<i>Escherichia coli</i> glycogen metabolism is controlled by the PhoP-PhoQ regulatory system at submillimolar environmental Mg <sup>2+</sup> concentrations, and is highly interconnected with a wide variety of cellular processes <i>Published as BJ Immediate Publication 25 August 2009, doi:10.1042/BJ20090980</i>	M. Montero, G. Eydallin, A.M. Viale, G. Almagro, F.J. Muñoz, M. Rahimpour, M.T. Sesma, E. Baroja-Fernández and J. Pozueta-Romero	 online data	129–141
The human <i>HIF</i> (hypoxia-inducible factor)-3 $\alpha$ gene is a HIF-1 target gene and may modulate hypoxic gene induction <i>Published as BJ Immediate Publication 21 August 2009, doi:10.1042/BJ20090120</i>	T. Tanaka, M. Wiesener, W. Bernhardt, K.-U. Eckardt and C. Warnecke	 online data	143–151
Regulation of Rnd3 localization and function by protein kinase C $\alpha$ -mediated phosphorylation <i>Published as BJ Immediate Publication 1 September 2009, doi:10.1042/BJ20082377</i>	J.P. Madigan, B.O. Bodemann, D.C. Brady, B.J. Dewar, P.J. Keller, M. Leitges, M.R. Philips, A.J. Ridley, C.J. Der and A.D. Cox		153–161