

ADVERTISING RATES

Cover and Special Positions	£90
Full Page	£70
Half Page	£40
Quarter Page	£25

Agency Commission	10%
Publisher's Discount	10%

All communications regarding advertising matters should be addressed to:

Advertisement Section,
Biochemical Society,
7 Warwick Court,
London WC1R 5DP

Telephone: 01-242 1076
(4 lines)

THE BIOCHEMICAL JOURNAL

The Biochemical Journal Cumulative Indices

Save Research Time

Busy researchers and authors find that these multi-volume indices are invaluable. Where time is money the cost is recovered quickly. Produced as a service to subscribers and priced accordingly.

Volumes	£ p	U.S.\$
21-30	0.75	2.50
31-40	2.00	5.50
41-65	2.75	7.50
66-90	3.25	9.00
91-115	5.25	13.00
116-130	5.50	15.00

All prices include postage (by surface mail to overseas)

Available from:


The Biochemical Society
(Publications)

PO BOX 32 Commerce Way
Whitehall Industrial Estate
Colchester CO2 8HP Essex

The **amicon** DC2/CH3
 Hollow Fibre Systems
 FOR THE RAPID
Concentration/Dialysis
of Macromolecular Solutions



DC2



CH3

SYSTEM FEATURES

- * High Ultrafiltration Rates
- * Process from 200 ml to 2000 ml
- * Choice of membrane fibres 10,000 and 50,000 MW cuts offs

- * Concentration or dialysis selected by switch and tube changes
- * Fast operation minimises possible denaturation
- * Sample volume maintained automatically during desalting
- * Ultrafiltration rate is indicated on built-in flowmeter

TYPICAL APPLICATION DC2

Problem: Salt exchange in 750 ml of 3% mixed Globulin.

Solution: DC2 Dialyzer/Concentrator with 50,000 MW Diafiber Cartridge

Reasoning: The high surface area of this integrated system makes for optimal exchange time.

- * Can be used for concentration or fractionation
- * Variable back pressure valve prevents fibre damage

TYPICAL APPLICATION CH3

99% of most common salts in 1 litre of water can be exchanged in less than 4 hours.

DIAFIBER MEMBRANE CARTRIDGES ARE CENTRAL TO BOTH SYSTEMS

Both the DC2 and the CH3 systems each include DIAFIBER membrane cartridges. These contain 1000 fibres with an effective area of 900 cm². Each fibre is a uniform anisotropic (skinned) membrane cylinder whose inner retentive-side is exposed to the process stream. As the liquid flows along the thin channel inside each fibre, shear forces minimise high solute concentration at the membrane surface (concentration polarization) and support transmembrane flow.

Diafiber membrane cartridges are made from environmentally resistant polymers and are completely non-cellulosic. This gives them far higher ultrafiltration rates than homogeneous cellulosic fibres of equal area. Cartridges are available with nominal retentions of 10,000 and 50,000 MW. Unaffected by enzymes, they offer months of useful life without failure and are easily cleansed for reuse by circulating water and detergent through the system.

amicon

57 QUEENS ROAD, HIGH WYCOMBE, BUCKS
 Telephone: HIGH WYCOMBE 21898

ANNOUNCING

**BIOCHEMICAL SOCIETY SYMPOSIA
No. 37**

**THE STRUCTURE AND FUNCTIONS
OF EUKARYOTIC RIBOSOMES**

(January 1974)

130pp. £3.50 (U.S. \$10.50)

This symposium was held in the University of Glasgow in June 1972 as part of the Biochemical Society Meetings Programme. The speakers who were invited to take part in this symposium were chosen because of their intimate and expert knowledge of the latest developments in our understanding of the structure and formation of ribosomes and particular attention is paid to ribosomes from eukaryotic cells. This publication contains the full papers given at the symposium, together, with plates, tables and figures.

List of contents and authors:

Chairman's Opening Remarks by **J. N. Davidson**

Ribosomal Proteins and the Assembly of Ribosomes in Eukaryotes by **J. R. Warner, A. Kumar, S. A. Udem & R. S. Wu**

Ribosomal Ribonucleic Acid and Ribosome Formation in Hela Cells by **B. E. H. Maden, M. Salim & J. Shepherd**

Structure and Function of Ribosomal Proteins of Escherichia coli by **H. G. Wittman, R. R. Crichton & G. Stoffler**

80S-Ribosomal Proteins by **H. J. Gould, O. H. W. Martini & H. S. King**

The Genes for Ribosomal Ribonucleic Acid by **P. J. Ford**

Ribosome Formation in Yeast by **T. H. Rozijn**

Ribosomal Ribonucleic Acid in Evolution by **U. E. Loening**

Regulation of Ribosome Synthesis by **R. P. Perry**

Chairman's Concluding Remarks by **H. R. V. Arnstein**

Send orders to:

The Biochemical Society (Publications)

P.O. Box 32, Commerce Way,

Colchester CO2 8HP, Essex