

# Clinical SCIENCE

## C O N T E N T S

### COMMENT

**Familial combined hyperlipidaemia: how can genetic disorders be common, complex and comprehensible?**

A. Sniderman, S. D. Bailey and J. C. Engert (see accompanying paper pp. 375–381)

365–367

Published as Immediate Publication 4 July 2007, doi:10.1042/CS20070220

### PAPERS

**Assessment of orthostatic fluid shifts with strain gauge plethysmography**

R. D. Thijs, M. Bruijnzeels, A. M. Kamper, A. D. van Dijk and J. G. van Dijk

369–374

Published as Immediate Publication 6 June 2007, doi:10.1042/CS20070060

**Five-year follow-up of waist circumference, insulin and ALT levels in familial combined hyperlipidaemia**

M. C. G. J. Brouwers, M. M. J. van Greevenbroek, V. M. M.-J. Vermeulen, J. M. J. P. van Lin,  
C. J. H. van der Kallen and T. W. A. de Bruin



375–381

Published as Immediate Publication 12 June 2007, doi:10.1042/CS20070101

**Relationship of waist and hip circumference with coagulation and fibrinolysis in postmenopausal women**

R. E. Peverill, H. J. Teede, E. Malan, D. Kotsopoulos, J. J. Smolich and B. P. McGrath

383–391

Published as Immediate Publication 4 July 2007, doi:10.1042/CS20070084

### CORRESPONDENCE

**Global longitudinal strain by two-dimensional speckle tracking echocardiography is closely related to myocardial infarct size in chronic ischaemic heart disease**

Z.-Z. Song

393

Published as Immediate Publication 4 June 2007, doi:10.1042/CS20070171

**Global longitudinal strain by two-dimensional speckle tracking echocardiography is closely related to myocardial infarct size in chronic ischaemic heart disease: authors' reply**

O. Gjesdal and T. Edvardsen

394–395

Published as Immediate Publication 4 June 2007, doi:10.1042/CS20070177