PUBLISHED BY THE MEDICAL RESEARCH SOCIETY AND THE BIOCHEMICAL SOCIETY

© The Medical Research Society and the Biochemical Society 1983 ISSN 0143-5221

Printed in Great Britain by Spottiswoode Ballantyne Ltd.
Colchester and London

CLINICAL SCIENCE

1983

Guidance for Authors

CONTENTS

1. POLICY OF THE JOURNAL

ultimate responsibility of acceptance for publication

1	D-11-					1.1. Scope
٠.	Poucy	of the Journal				Clinical Science publishes papers in the field of
	1.1.	Scope			i	clinical investigation, provided they are of a suit-
	1.2.	The Editorial Board			i	able standard and contribute to the advancement of
	1.3.	The editorial process			i	knowledge in this field. The term 'clinical in-
		Ethics of investigations o	n			vestigation' is used in its broad sense to include
		human subjects .			ii	studies in animals and the whole range of bio-
	1.5.	Originality of papers			ii	chemical, physiological, immunological and other
2.		ission of Manuscripts: Ge	eneral			approaches that may have relevance to disease in
		nation and Format				man. Studies which are confined to normal sub-
	2.1.	General			ii	jects, or animals, or are purely methodological in
	2.2.	Full papers			ii	nature may be acceptable. The material presented
	2.3.	Short Communications			iii	should permit conclusions to be drawn and should
	2.4.	Correspondence .			iii	not be only of a preliminary nature. The journal
	2.5.	Arrangements for large		ts		publishes four types of manuscript, namely invited
		of information .			iii	Editorial Reviews, Full Papers, Short Communi-
	2.6.	Proof corrections .			iii	cations and Correspondence. In addition, Clinical
	2.7.	Offprints			iii	Science publishes abstracts of the proceedings of
	2.8.	Availability on MEDLINE			iii	the Medical Research Society and also that
3.	Misc	ellaneous Notes				Society's Annual Guest Lecture.
	3.1.	Abbreviations .			iv	•
	3.2.	Anatomical nomenclatur			iv	1.2. The Editorial Board
	3.3.	Animals, plants and mic	ro-			The Board comprises Editors for the Medical
		organisms			iv	Research Society and the Biochemical Society and
	3.4.	Buffers and salts .			iv	a Chairman and two Deputy Chairmen who are
	3.5.	Computer modelling			vi	drawn alternately from the two Societies. Members
	3.6.	Doses			iv	of the Board retire after a maximum of 5 years;
		Enzymes			iv	the Chairman serves in his capacity for 2 years.
	3.8.	Evaluation of measurem		-		The membership of the Board is designed to cover
	• • • •	procedures			iv	as wide a range of interests as possible.
	3.9.	Figures and Tables .			iv	The main function of the Board is to decide on
		Footnotes			v	the acceptability of submitted papers, but it also
		Isotope measurements			v	deals with general matters of editorial policy.
		Radionuclide application			v	Financial policy is dealt with separately by the
		Methods		_	v	Committee of Management.
	3.14.	Nomenclature of disease		_	v	1.3. The editorial process
		Powers in Tables and Fi			v	A submitted paper is first read by the Chairman
		References			v	or one of the Deputy Chairmen of the Editorial
		Solutions			vi	Board who then sends it to an Editor. The latter
		Spectrophotometric data			vi	considers the paper in detail and sends it to one or
		Spelling			vi	more referees (who remain anonymous) from out-
	3.20.	Statistics			vi	side the membership of the Board. The Editor
		Trade names		•	vi	returns it with his recommendation to the Chair-
4		s: The SI System .			vi	man who then writes formally to the authors. The

vii

4. Units: The SI System

5. Abbreviations, Conventions etc.

lies with the Chairman. If the Chairman is for any reason unavailable, a Deputy Chairman assumes this function.

1.4. Ethics of investigations on human subjects

Authors must state in the text of their paper the manner in which they have complied, where necessary, with the recommendations on human investigations published in the Medical Research Council report of 1962/63 [British Medical Journal (1964) ii, 178–180]. Consent must be obtained from each patient or subject after full explanation of the purpose, nature and risks of all procedures used and the fact that such consent has been given should be recorded in the paper. Papers should also state that the Ethical Committee of the Institution in which the work was performed has given approval to the protocol. The Editorial Board will not accept papers the ethical aspects of which are, in the Board's opinion, open to doubt.

1.5. Originality of papers

Submission of a paper to the Editorial Board is taken to imply that it reports work that has not been published in either the same or a substantially similar form, that it is not under consideration for publication elsewhere and that, if accepted for publication by Clinical Science, it will not be published elsewhere in the same form, either in English or in any other language, without the consent of the Editorial Board. This does not usually apply to previous publication of oral communications in brief abstract form. In such cases authors should enclose copies of the abstracts or previous publications. The author, or in the case of multiple authorship the authors, will be asked to sign a statement vesting the copyright in the publishers. Requests for consent for reproduction of material published in Clinical Science should be addressed to the Editorial Manager.

2. SUBMISSION OF MANUSCRIPTS: GENERAL INFORMATION AND FORMAT

2.1. General

Papers submitted for publication should be sent to the Editorial Manager, Clinical Science, 7 Warwick Court, London WC1R 5DP.

The submission should contain four copies (of which three may be photocopies) of the typescript, Tables, Figures etc. The authors should retain one copy of the paper. The Editorial Board does not accept responsibility for damage or loss of papers

submitted, although great care is taken to ensure safety and confidentiality of the typescript during the editorial process. In the case of multiple authorship, the covering letter should indicate that the approval of all co-authors has been obtained.

Papers should be presented so that they are intelligible to the non-specialist reader of the journal. This is particularly important in highly specialized fields and a very brief resume of the current state of knowledge is usually helpful. Certain types of material, e.g. mathematical formulations requiring more than trivial derivations, should be given in a separate Appendix.

Where the reader is referred to previous works by the same author(s) for important details relevant to the present work, copies or reprints of the publication should be sent with the typescript. This is of particular importance in relation to methodology.

The dates of receipt and acceptance of the paper will be published. If the paper has to be returned to the authors for revision and is not resubmitted within 1 month, the date of receipt will be revised accordingly. Papers returned by authors later than 12 months after the original submission date will be treated as new papers. For Short Communications the published date will always be that of receipt of the final version. It is emphasized that badly presented or unduly long papers will be returned for revision and delays in publication will be inevitable. Similar delays will be incurred if the typescript is not prepared strictly in accordance with the instructions detailed below.

2.2. Full papers

The authors should refer to a current issue of Clinical Science to make themselves familiar with the general layout. Concise presentation is very important for rising costs are a severe constraint on space. The length of manuscript and the number of Figures and Tables must be kept to a minimum. Extensive Tables of data can be deposited with the Royal Society of Medicine (see 2.5). Guidance for Authors is usually published in the January issue of the journal, and revised periodically.

Typescripts should be, in general, arranged as follows:

(a) Title page. Title: this should be as informative as possible, since titles of papers are being increasingly used in indexing and coding for information storage and retrieval. The title should indicate the species in which the observations reported have been made. The numbering of parts in a series of papers is not permitted.

List of authors' names (degrees and appointments are not required).

Laboratory or Institute of origin.

Key words: for indexing the subject of the paper; they should, if possible, be selected from the current issues of 'Medical Subject Headings' (MeSH), produced by the *Index Medicus*.

Short title: for use as a running heading in the printed text; it should not exceed forty-five characters and spaces.

Author for correspondence: the name and address of the author to whom queries and requests for reprints should be sent.

- (b) Summary. This should be a brief statement arranged in numbered paragraphs of what was done, what was found and what was concluded and should rarely exceed 250 words. Abbreviations should be avoided as far as possible and must be defined. Statistical and methodological details including exact doses should also be avoided unless they are essential to the understanding of the summary.
- (c) Introduction. This should contain a clear statement of the reason for doing the work, but should not include either the findings or the conclusions.
- (d) Methods. The aim should be to give sufficient information in the text or by reference to permit the work to be repeated without the need to communicate with the author.
- (e) Results. This section should not include material appropriate to the Discussion section.
- (f) Discussion. This should not contain results and should be pertinent to the data presented.
- (g) Acknowledgments. These should be as brief as possible.
 - (h) References. See p. v for the correct format.
 - (i) Figures and Tables. See p. iv.

2.3. Short Communications

The Short Communication should describe completed work, and should not be merely a preliminary communication. The format of Short Communications should be similar to that of Full Papers, but should not exceed 1200 words of text. One Figure or Table is allowed, but if neither is included the text may be expanded to 1400 words. The passage of Short Communications through the editorial process can frequently be expedited and contributors are encouraged to take advantage of these facilities when rapid publication is of importance and the material can be presented concisely. The paper should appear in print within 3 months of acceptance. When submitting Short Communi-

cations, authors should make it quite clear that the work is intended to be treated as a Short Communication.

2.4. Correspondence

Letters containing original observations or critical assessments of material published in Clinical Science, including Editorial Reviews, will be considered for the Correspondence section of the journal. Letters should be no longer than 750 words, with one Figure or Table and up to six references, or 1000 words maximum without a Figure or Table. Letters relating to material previously published in Clinical Science should be submitted within 6 months of the appearance of the article concerned. They will be sent to the authors for comment and both the letter and any reply by the author will be published together. Further correspondence arising therefrom will also be considered for publication. Consideration will also be given to publication of letters on ethical matters.

2.5. Arrangements for large amounts of information

It is impracticable to publish very large sets of individual values or very large numbers of diagrams, and under these circumstances a summary of the information only should be included in the paper. The information from which the summary was derived should be submitted with the typescript and, if the latter is accepted, the Editors may ask for a copy of the full information and diagrams to be deposited with the Librarian, the Royal Society of Medicine, 1 Wimpole Street, London W1M 8AE, who will issue copies on request. Experience has shown that such requests are frequently received.

2.6. Proof corrections

These are expensive and corrections of other than printers' errors may have to be charged to the author.

2.7. Offprints

Fifty offprints are supplied free and additional copies may be obtained at terms, based upon the cost of production, that will be given with the proofs. All offprints should be ordered when the proofs are returned.

2.8. Availability on MEDLINE

Summaries of papers in Clinical Science are available on-line on teleprinters participating in the

MEDLINE system run by the National Library of Medicine, National Institutes of Health, Bethesda, Maryland, U.S.A.

3. MISCELLANEOUS NOTES

3.1. Abbreviations

Abbreviations should be avoided; if used they must be defined at the first mention; new abbreviations should be coined only for unwieldy names which occur frequently. Abbreviations should not appear in the title nor, if possible, in the Summary. A list of accepted abbreviations is on p. vi. Numbers, not initials, should be used for patients and subjects.

3.2. Anatomical nomenclature

This should follow the recommendations of the International Anatomical Nomenclature Committee (1966) *Nomina Anatomica*, 3rd edn, Excerpta Medica Foundation, Amsterdam.

3.3. Animals, plants and micro-organisms

The full binomial specific names should be given at first mention for all experimental animals other than common laboratory animals. The strain and, if possible, the source of laboratory animals should be stated. Thereafter in the text, single letter abbreviations may be given for the genus; if two genera with the same initial letter are studied, abbreviations such as *Staph*. and *Strep*. should be used.

3.4. Buffers and salts

The acidic and basic components should be given, together with the pH. Alternatively, a reference to the composition of the buffer should be given. Further details are provided in the *Biochemical Journal* (1978) 169, 9.

When describing solutions containing organic anions and their parent acids, the salt designator (e.g. lactate, urate, oxalate) should be used in preference to the name of the acid (lactic, uric, oxalic) unless it is certain that virtually all of the acid is in the undissociated form.

The composition of incubation media should be described, or a reference to the composition should be given.

3.5. Computer modelling

Papers concerned primarily with computer modelling techniques are acceptable provided that use of such techniques leads to a clear choice between two or more alternative hypotheses, or to the formulation of a new hypothesis amenable to experimental challenge or verification, or provides some new insight into the behaviour of a particular

physiological system. Extensive technical details of hardware and software should not be given.

3.6. Doses

Doses of drugs should be expressed in mass terms, e.g. milligrams (mg) or grams (g), and also (in parentheses) in molar terms, e.g. mmol, mol, where this appears to be relevant. Molecular weights of many drugs may be found in *The Merck Index*, 8th edn, Merck and Co. Inc., N.J., U.S.A.

3.7. Enzymes

Nomenclature should follow that given in Enzyme Nomenclature (1978), Academic Press, London and New York, and the Enzyme Commission (EC) number should be quoted at the first mention. Where an enzyme has a commonly used informal name, this may be employed after the first formal identification. A unit of enzyme activity should preferably be expressed as that amount of material which will catalyse transformation of 1 µmol of the substrate/min; under defined conditions, including temperature and pH. Alternatively, or when the natural substrate has not been fully defined, activity should be expressed in terms of units of activity relative to that of a recognized reference preparation, assayed under identical conditions. Activities of enzymes should normally be expressed as units/ml or units/mg of protein.

3.8. Evaluation of measurement procedures

When a new measuring procedure has been used, or when an established procedure has been applied in a novel fashion, an estimate of the precision of the procedure should be given. This should, as far as possible, indicate what sources of variation have been included in this estimate, e.g. variation of immediate replication, variation within different times of day, or from day to day etc.

If the precision of measurement varies in proportion to the magnitude of the values obtained, it can best be expressed as the coefficient of variation; otherwise it should be expressed by an estimate of the (constant) standard error of a single observation, or by estimates at several points within the range of observed values.

When recovery experiments are described the approximate ratio of the amount added to the amount already present and the stage of the procedure at which the addition was made should be stated.

3.9. Figures and Tables

These are expensive to print and their number should be kept to a minimum. Their appropriate position in the paper should be indicated in the margin of the text. References to Figures and Tables should be in Arabic numerals, e.g. Fig. 3, and they should be numbered in order of appearance. In general, the same data should not be presented in both a Figure and a Table.

Figures, with captions attached, should be supplied as original drawings or matt photographs together with photocopies. All Figures should have their number and the authors' names written in pencil on the back; the top of the Figure should be indicated with a pencilled arrow. Acceptable symbols for experimental points are \bigcirc , \triangle , \square , O, \triangle , \square . The symbols \times or + must be avoided. The same symbols must not be used for two curves where the points might be confused. For scatter diagrams, solid symbols are preferred. When a particular variable appears in more than one Figure, the same symbol should be used for it throughout, if possible.

Curves should not be drawn beyond the experimental points, nor should axes extend appreciably beyond the data. Only essential information that cannot readily be included in the legend should be written within the Figure.

Figures for half-tone reproduction should be submitted as glossy prints. They are particularly expensive to print and their use should be avoided as far as possible.

Tables should be typed separately from the text. They should have an underlined title followed by any legend.

Captions for the Figures, and titles and legends for the Tables, should make them readily understandable without reference to the text. Adequate statistical information, including that on regression lines, should be included in Figure captions where appropriate.

3.10. Footnotes

These should be avoided as far as possible but where they are used in Tables they should be identified by the symbols $^{\bullet}$ † † § § § § , in that order.

3.11. Isotope measurements

The information given should include (a) conditions of radioactivity counting, e.g. infinitely thick, infinitely thin; (b) the nature of the phosphor used in liquid-scintillation counting; (c) details of corrections made to the observed count rate, e.g. for 'quenching' or 'cross-over'; (d) standard deviation of the results or a statement of the minimum total counts above background collected and the background value.

In general the specific radioactivity of the starting materials should be given, preferably in terms of radioactivity per unit weight or, for stable isotopes, as atoms % excess.

Pending the general introduction of SI units radioactivity should continue to be expressed in terms of the curie (Ci) followed by the corresponding figure in terms of the becquerel (Bq: disintegrations/s), in parentheses, and suitably rounded.

3.12. Radionuclide applications in man

If new or modified radionuclide applications in man are described, an estimate of the maximal possible radiation dose to the body and critical organs should be given.

For the time being this can continue to be expressed in rems, but with the corresponding figure in sieverts (Sv) given in parentheses after it.

3.13. Methods

In describing certain techniques, namely centrifugation (when the conditions are critical), chromatography and electrophoresis, authors should follow the recommendations published by the Biochemical Society (currently, *Biochemical Journal* (1981) 193, 1–21).

3.14. Nomenclature of disease

This should follow the *International Classification of Disease* (8th revision, World Health Organization, Geneva, 1969) as far as possible.

3.15. Powers in Tables and Figures

Care is needed where powers are used in Table headings and in Figures to avoid numbers with an inconvenient number of digits. For example: (i) an entry '2' under the heading 10^3k means that the value of k is 0.002; an entry '2' under the heading $10^{-3}k$ means that the value of k is 2000. (ii) A concentration 0.00015 mol/l may be expressed as 0.15 under the heading 'concn. (mmol/l)' or as 150 under the heading 'concn. (μ mol/l)' or as 15 under the heading '10⁵ × concn. (mol/l)', but not as 15 under the heading 'concn. (mol/l) × 10^{-5})'.

3.16. References

The numerical citation system is now used: references in the text are numbered consecutively in the order in which they are first mentioned, the numerals being given in brackets, e.g. [22]. References cited in Figure legends or Tables only should be numbered in a sequence determined by the position of the first mention in the text of the Figure or Table. References should be listed in

numerical order and the names of all authors of a paper should be given, with the full title of the paper and the source details in full including the first and last page numbers, e.g.

[2] CLARK, T.J.H., FREEDMAN, S., CAMPBELL, E.J.M. & WINN, B.R. (1969) The ventilatory capacity of patients with chronic airways obstruction. Clinical Science, 36, 307-316.

When the quotation is from a book, the following format should be used, giving the relevant page or chapter number:

- [20] MOLLISON, P.L. (1967) Blood Transfusion in Clinical Medicine, 4th edn, p. 50. Blackwell Scientific Publications, Oxford.
- [22] Reid, L. (1968) In: The Lung, p. 87. Ed. Liebow, A.A. & Smith, D.E. Williams and Wilkins, Baltimore.

References to 'personal communications' and 'unpublished work' should appear in the text only and not in the list of references. The name and initials of the source of information should be given. When the reference is to material that has been accepted for publication but has not yet been published, this should be indicated in the list of references by 'In press' together with the name of the relevant journal and, if possible, the expected date of publication. If such a citation is of major relevance to the manuscript submitted for publication authors are advised that the editorial process might be expedited by the inclusion of a copy of such work. In the case of quotations from personal communications the authors should state in the covering letter that permission for quotation has been obtained.

3.17. Solutions

Concentration of solutions should be described where possible in molar terms (mol/l and subunits thereof), stating the molecular particle weight if necessary. Values should not be expressed in terms of normality or equivalents. Mass concentration should be expressed as g/l or subunits thereof, for example mg/l or μ g/l. For solutions of salts, molar concentration is always preferred to avoid ambiguity as to whether anhydrous or hydrated compounds are used. Concentrations of aqueous solutions should be given as mol/l or mol/kg (g/l or g/kg if not expressed in molar terms) rather than % (w/v) or % (w/w). It should always be made clear whether concentrations of components in a reaction mixture are final concentrations or the concentrations in solutions added.

3.18. Spectrophotometric data

The term 'absorbance' $[\log (I_0/I)]$ should be used rather than 'optical density' or 'extinction'. The

solvent, if other than water, should be specified. Symbols used are: A, absorbance; a, specific absorption coefficient (litre g^{-1} cm⁻¹) (alternatively use $A_{1cm}^{1\%}$); ε , molar absorption coefficient (the absorbance of a molar solution in a 1 cm lightpath) (litre mol⁻¹ cm⁻¹, not cm² mol⁻¹).

3.19. Spelling

Clinical Science uses as standards for spelling the Concise or Shorter Oxford Dictionary of Current English (Clarendon Press, Oxford) and Butterworth's Medical Dictionary (Butterworths, London).

3.20. Statistics

Papers are frequently returned for revision (and their publication consequently delayed) because the authors use inappropriate statistical methods. Two common errors are the use of means, standard deviations and standard errors in the description and interpretation of grossly non-normally distributed data and the application of t-tests for the significance of difference between means in similar circumstances, or when the variances of the two groups are non-homogeneous. In some circumstances it may be more appropriate to provide a 'scattergram' than a statistical summary.

A reference should be given for all methods used to assess the probability of a result being due to chance. The format for expressing mean values and standard deviations or standard errors of the mean is, for example: mean cardiac output 10.4 l/min (sD 1.2; n=11). Degrees of freedom should be indicated where appropriate. Levels of significance are expressed in the form P < 0.01.

3.21. Trade names

The name and address of the supplier of special apparatus and of biochemicals should be given. In the case of drugs, approved names should always be given with trade names and manufacturers in parentheses.

4. UNITS: THE SI SYSTEM

The recommended Système International (SI) units [see Quantities, Units and Symbols, 2nd edn (1975) The Royal Society, London] are used by Clinical Science. All papers submitted should use these units except for blood pressure values, which should be expressed in mmHg, or gas tensions, where values at the author's discretion may be given as mmHg (with kPa in parentheses) or as kPa (with mmHg in parentheses) in the text and either as mmHg or as kPa in Figures, which (if practicable) should have scales in both units. Airways pressure should be expressed in kPa. Where molecular weight is known, the amount of a chemical or drug should be expressed in mol or in

an appropriate subunit, e.g. mmol. Energy should be expressed in joules (J).

The basic SI units and their symbols are as follows:

Physical quantity	Name	Symbol
length	metre	m
mass	kilogram	kg
time	second	S
electric current	ampere	Α
thermodynamic temperature	kelvin	K
luminous intensity	candela	cd
amounts of substance	mole	mol

The following are examples of derived SI units:

Physical quantity	Name	Symbol	Definition
energy	joule	J	kg m ² s ⁻²
force	newton	N	$kg \ m \ s^{-2} = J \ m^{-1}$
power	watt	W	$kg m^2 s^{-3} =$
pressure	pascal	Pa	$kg m^{-1} s^{-2} = N m^{-2}$
electric charge	coulomb	С	As
electric potential	volt	v	$kg m^2 s^{-2} A^{-1}$ = $J A^{-1} s^{-1}$
electric resistance	ohm	Ω	$kg m^2 s^{-3} A^{-2}$ = $V A^{-1}$
electric conductance	siemens	S	$kg^{-1} m^{-2} s^3 A^2$ = Ω^{-1}
electric capacitance	farad	F	$A^2 s^3 kg^{-1} m^{-2}$ = $A s V^{-1}$
frequency	hertz	Hz	s ⁻¹
volume	litre	1	10 ⁻³ m ³

The word 'litre' has been accepted as a special name for cubic decimetre (1 litre = 1 dm^3).

Both the basic and derived SI units, including the symbols of derived units that have special names, may be preceded by prefixes to indicate multiples and submultiples. The prefixes should be as follows:

	Prefix	Symbol	Multiple	Prefix	Symbol
10 ⁶	mega	M	10-3	milli	m
10³	kilo	k	10-6	micro	μ
10 ²	hecto	h*	10 ⁻⁹	nano	n
10	deka	da	10-12	pico	р
10 ⁻¹	deci	d*	10-15	femto	f
10-2	centi	*ء			

To be avoided where possible (except for cm).

Compound prefixes should not be used, e.g. 10^{-9} m should be represented by 1 nm, not 1 m μ m.

Notes:

- (i) Full stops are not used after symbols.
- (ii) Minutes (min), hours (h), days and years will continue to be used in addition to the SI unit of time [the second (s)].
- (iii) The solidus may be used in a unit as long as it does not have to be employed more than once,

e.g. mmol/l is acceptable, but ml/min/kg is not, and should be replaced by ml min⁻¹ kg⁻¹.

5. ABBREVIATIONS, CONVENTIONS, DEFINITIONS, SYMBOLS AND SPECIAL COMMENTS

As well as standard symbols and abbreviations that have been accepted by international bodies, and which can be used without definition, this list shows selected abbreviations in the form of groups of capital letters (e.g. ALA, ECF, MCHC) which when used must be defined in the text as indicated on p. iv. The standard abbreviations for amino acids are only for use in Figures and Tables or for peptide sequences.

absorbance	A
acceleration due to gravity	g
adenosine 3': 5'-cyclic mono-	cyclic AMP
phosphate	0,00 111
	AMP
adenosine 5'-phosphate	
adenosine 5'-pyrophosphate	ADP
adenosine 5'-triphosphate	ATP
adenosine triphosphatase	ATPase
adrenocorticotropic hormone	ACTH
adrenoceptor (see also	
blocking agents)	
alanine	Ala
alternating current	a.c.
alveolar minute ventilation	\dot{V}_{A}
alveolar to arterial oxygen	$(\stackrel{\wedge}{P}_{A}, o_2 - Pa, o_2)$
tension difference	(1 A,O ₂ 1 a,O ₂)
	A
ampere aminolaevulinic acid	ALA
angiotensin	ANG; reference amino
	acid abbreviations are
	used as prefix within
	brackets: e.g.
	[Sar ¹ ,Val ⁵ ,Ala ⁸]ANG
Ångstrom (Å)	not used; express in nm
	$(1 \text{ Ångstrom} = 10^{-1})$
	nm)
antidiuretic hormone	ADH (when referring to
	the physiological
	secretion)
arginine	
arteriovenous	Arg
arteriovenous	a-v: permitted in
•	Figures and Tables
asparagine	Asn
aspartic acid	Asp
atmosphere (unit of pressure)	not used; express in kPa
	(1 atmosphere =
	101·325 kPa)
atomic weight	at. wt.
becquerel	Bq (1 d.p.s.)
blocking agents	e.g. β-adrenoceptor
blocking agents	antagonists preferred
blood messages	
blood pressure	express in mmHg
blood urea nitrogen	not used; recalculate as
	urea, express in
	mmol/l
blood volume	BV
body temperature and pres-	BTPS
sure, saturated	
.,	

****	Guidancejo	714111015	
British Pharmacopoeia	write in full and give	electromotive force	e.m.f.
	edition	electron spin resonance	e.s.r.
calculated	calc. (in Tablesonly)	electronvolt	eV (for radiation
'Calorie' (= 1000 cal)	not used; recalculate as		energies)
	kilojoules (1 'Calorie'	equation	eqn.
	= 4.184 kJ)	equivalents (amount of a	not used; recalculate in
carbon dioxide output (in res-	VCO ₂ ; express in ml	chemical)	molar terms
piratory physiology)	STP/min	erythrocyte count	express as 1012 cells/l
cardiac frequency	f_c ; in beats/min	erythrocyte sedimentation	ESR
cardiac output	express in I/min	rate	
centimetre	cm	ethanol, ethanolic	not ethyl alcohol or al-
clearance of x	Cx A and and CoA		coholic
coenzyme A and its acyl derivatives	CoA and acyl-CoA	ethylenediaminetetra-acetate	EDTA
	cf.	exchangeable	Na, K, etc., for total
compare complement fractions	C1-C9	•	exchangeable sodium,
compliance (respiratory	C; express in 1 kPa ⁻¹		potassium etc.
physiology)	C, express in 1 kFa	Experiment (with reference	Expt.; plural, Expts.
concentrated	conc.	numeral)	zapta, piarai, zapto.
concentration	concn.; may be denoted	expired minute ventilation	$\dot{V}_{_{\mathbf{E}}}$
concenti attori	[]; e.g. plasma	extinction	use absorbance
	[HCO ₃]	extracellular fluid	ECF
conductance (respiratory	G; express in 1 s ⁻¹ kPa ⁻¹	extracellular fluid volume	ECFV
physiology)	0, onpress in 10	extraction ratio of x (renal)	E _x
correlation coefficient	r: may be used without	Figure (with reference	Fig.; plural, Figs.
	definition	numeral)	g, p,g
counts/min, counts/s	c.p.m., c.p.s.	filtered load of x (renal)	F
cubic centimetres	use ml	follicle-stimulating hormone	<i>F</i> x FSH
curie	Ci (1 Ci = 3.7×10^{10}	forced expiratory volume in	FEV _{1.0}
	d.p.s.)	1.0 s	- ~ · 1.0
cycle/s	Hz	fractional concentration in	F
cysteine	Cys	dry gas	1
dates	e.g. 11 August 1970	fractional disappearance rate	k (as in $A = A_0 e^{-\kappa t}$)
dead-space minute ventilation	ν _D ν _D °C	frequency of respiration	$f_{\rm R}$; in breaths/min
dead-space volume	V_{D}	functional residual capacity	FRC
degrees, Celsius or centigrade	°C	gas-liquid chromatography	g.l.c.
deoxy (prefix)	not desoxy	gas transfer factor	T; in mmol min ⁻¹ kPa ⁻¹
deoxycorticosterone	DOC	glomerular filtration rate	GFR
deoxycorticosterone acetate	DOCA	glutamic acid	Glu
deoxyribonucleic acid	DNA	glutamine	Gln
dialysate	diffusate preferred;	glutathione	GSH (reduced); GSSG
	'dialysate' should be	3	(oxidized)
distinct and a sthert called an	clearly defined	glycine	Gly
diethylaminoethylcellulose differential of x with respect to	DEAE-cellulose	gram(me)	g
time	$x (= \alpha x/\alpha t)$	gravitational field, unit of	g
1,25-dihydroxycholecalciferol	1,25-(OH),D,	(9·81 m s ⁻¹)	
dilute	dil.	growth hormone	GH; if human, HGH
2,3-diphosphoglycerate	2,3-DPG	guery	Gy (100 rads)
direct current	• ·	haematocrit	not allowed; use packed
disintegrations/min	d.c.		cell volume (PCV)
disintegrations/s	d.p.m	haemoglobin	Hb; express in g/dl
•	d.p.s.	half-life	<i>t</i> ₄
dissociation constant acidic	ν	hertz (s ⁻¹)	$t_{ m i} \ { m Hz}$
basic	K_a	histidine	His
	K_b	hour	h
apparent minus log of	e.g. K'_a	human chorionic gon-	HCG
_	pK	adotropin	
doses	avoid Latin designa-	human placental lactogen	HPL
	tions such as b.d. and t.i.d.	hydrocortisone	use cortisol
duna		hydrogen ion activity	aH; express in nmol/l
dyne	dyn; used for vascular	minus log of	pH
alastanas	resistance	25-hydroxycholecalciferol	-
elastance	E; express in Pa m ⁻³		25-(OH)D ₃
electrocardiogram	ECG	hydroxyproline	Нур
electroencephalogram	EEG	immunoglobulins	IgA, IgD, IgE, IgG, IgM

	•	•	
injection routes:	use abbreviations only in Figures	millimetre of mercury	mmHg; for blood pres- sure and, at authors'
intra-arterial	i.a.		discretion, for gas ten-
intramuscular	i.m.		sions: see p. vi (1
intraperitoneal	i.p.		mmHg = 0.133 kPa
intravenous	_ -	:!!!()	
	i.y.	millimolar (concentration)	mmol/l; not mm
subcutaneous	s.c.	millimole	mmol
international unit	i.u. (definition and	minimum	min.
	reference should	minute (60 s)	min
	be given for uncom-	molal	mol/kg
	mon or ambiguous	molar (concentration)	mol/l; not M
	applications, e.g. en-	molar absorption coefficient	ε (the absorbance of a
	zymes)	moiar abborpaon coemercine	molar solution in a
intracellular fluid	ICF		1 cm light-path)
intracellular fluid volume	ICFV	1	
ionic strength	I	mole	mol
_		molecular weight	mol. wt.
isoleucine	Ile	nicotinamide-adenine	NAD if oxidation state
isotonic	not used; specify com-	dinucleotide	not indicated
	position of fluid, e.g.		NAD+ if oxidized
	NaCl, 150 mmol/l		NADH if reduced
isotopically labelled com-	e.g. [U-14C]glucose,	nicotinamide-adenine	NADP if oxidation
pounds	[1-14C]glucose,	dinucleotide phosphate	state not indicated
•	sodium [1-14C]-	umucicoude phospitate	NADP+ if oxidized
	acetate; use 131I-		
	labelled albumin, not		NADPH if reduced
	[131I]albumin	normal	should not be used to
	for simple molecules:		denote the concentra-
			tion or osmolarity of
	¹⁴ CO ₂ , ³ H ₂ O		a solution
joule	j	normal temperature and	use standard temp-
kilogram(me)	kg	pressure	erature and pressure
kilopond	not used; 1 kilopond =	•	(STP)
	9-8067 N	nuclear magnetic resonance	n.m.r.
lactate dehydrogenase	LDH	number (in enumerations)	no. (in Tables only)
leucine	Leu	observed	obs. (in Tables only)
leucocyte count	express as 10° cells/l	ohm	Ω
lipoproteins (serum)	-	ornithine	Orn
high density	HDL		
low density	LDL	ortho-	<i>0-</i>
very low density	VLDL	orthophosphate (inorganic)	P_{l}
litre	1 (write in full if con-	osmolarity	express in osmol (or
	fusion with the		_ mosmol)/l
	numeral 1 is possible)	oxygen uptake per minute	Vo_2 ; express in ml
lana-ish (hana 10)		(in respiratory physiology)	STP/min
logarithm (base 10)	log	packed cell volume	PCV
logarithm (base e)	ln	page, pages	p., pp.
luteinizing hormone	LH	para-	p-
lysine	Lys	para-aminohippurate	PAH
maximum	max.	partial pressure	P; express in either kPa
mean corpuscular	MCH; express in pg		or mmHg (see p. vi)
haemoglobin		e.g. alveolar, of O ₂	PAO ₂
mean corpuscular	MCHC; express in g/dl	arterial, of CO ₂	Paco ₂
haemoglobin concentration		capillary, of O ₂	Pcapo,
mean corpuscular volume	MCV; express in fl (1	mixed venous, of CO,	$P\bar{v}co_{2}$
	$\mu \mathrm{m}^3 = 1 \mathrm{fl})$	pascal	Pa
median lethal dose	LD ₅₀	p e r	/
meta-	m-	•	/ %
melting point	m.p.	per cent	
methanol, methanolic	not methyl alcohol	petroleum ether	not used; use light
methionine	Met		petroleum and give
			boiling range
metre	m v	phenylalanine	Phe
Michaelis constant	K _m	plasma renin activity	express as pmol of
micromole	μmol		angiotensin I h ⁻¹
micron (10 ⁻⁶ m)	μ m; not μ		ml ⁻¹
milliequivalent	not used; give amount in	plasma volume	PV
	mmol	poise	1 poise = 10^{-1} N s
millilitre	ml		m ⁻²

		o. 1140.00 b	
potential difference power output	p.d. $W (1 W = 0.1635)$	specific conductance of airways	sGaw; express in s ⁻¹ kPa ⁻¹
precipitate	kpm/min) ppt. P; express in kPa	standard deviation standard error of the mean	SD sem may be used without definition
pressure	(except for blood pressures and gas	standard temperature and pressure	STP
	tensions: see p. 6); 1 kPa = 7.5 mm Hg	steroid nomenclature	see Biochemical Journal (1969) 113, 5-28;
probability of an event being due to chance alone	P	aulahudeul	(1972) 127, 613–617 use thiol or SH
proline	Pro	sulphydryl sum	Σ thio of SA
protein-bound iodine	PBI	Svedberg unit	S
(plasma) pulmonary capillary blood flow	Ċс	temperature (absolute) (empirical) temperature, thermodynamic	T ! °K
pyrophosphate (inorganic)	PPi	thin-layer chromatography	t.l.c.
rad (radiation dose; 10 ⁻⁵ J	not abbreviated	threonine	Thr
absorbed/g of material) red blood cell	(100 rads = 1 Gy) use erythrocyte;	thyrotrophic hormone	TSH TRH
700 01000 00H	express counts as 10^{12} cells/l	thyrotrophin-releasing hor- mone tidal volume	V _T
relative band speed (partition	R_F	time (symbol)	t^{T}
chromatography)		time of day	e.g. 18.15 hours
rem renin	100 ergs/g × quality factor	torr	not used; use kPa (1 torr = 0.133 kPa)
reimi	see plasma renin activity	total lung capacity tryptophan	TLC Trp
residual volume	RV	tubular maximal reabsorptive	T _{m,x}
resistance (rheological)	R; express in kPa l ⁻¹ s	capacity for x	•
respiratory exchange ratio (pulmonary)	R	tyrosine ultraviolet	Tyr u.v.
respiratory quotient (metabolic)	RQ	urinary concentration of x valency	U _x e.g. Fe ²⁺ , not Fe ⁺⁺
revolutions	rev.	valine variance ratio	Val F
rev./min	not r.p.m.; use g if possible (see p. viii)	vascular resistance	express in kPa l ⁻¹ s (with value in dyn s cm ⁻⁵
ribonucleic acid röntgen	RNA R		in parentheses);
saline	define at first mention {e.g. NaCl solution (154 mmol/l)]		primary values of dif- ferential vascular pres- sure (mmHg) and
saturation	S, e.g. Sao ₂ for arterial oxygen saturation (see partial pressure		flow (I/min) should always also be given in Tables or text as
	for other analogous	velocity	appropriate v; express as m s ⁻¹
second (time)	abbreviations) s	venous admixture	\dot{Q}_{va}
serine	Ser	veronal	used only for buffer mix-
sievert	Sv (1 J/kg × quality factor)		tures; otherwise use 5,5'-diethylbarbituric
solvent systems	e.g. butanol/acetic acid/ water (4:1:1, by vol.), butanol/ acetic acid (4:1, v/v)	viscosity, dynamic viscosity, kinematic vital capacity	acid η υ VC
species	sp., plural spp.	volt	V O; use O for blood flow
specific activity	sp. act. Confusion must be avoided	volume of blood (in cardio- respiratory physiology) watt	Q; use Q for blood flow rate W
	between e.g. specific radioactivity and the	watt wavelength	λ
	specific activity of an	weight	wt.
	enzyme	white blood cell	use leucocyte: express counts as 10° cells/l

Volume 64

AUTHOR INDEX

Аввотт, R.J. 617-622
ABRAMSON, S.B. 193-205
Afford, S.C. 223-230
Albano, J.D.M. 623-627
ALBERTI, K.G.M.M. 41-47
ARIEFF, A.I. 573-580
ARNER, P. 235-237
Aronson, J.K. 253-258
Aviram, M. 637-642
Ayres, J.G. 555-557
AZAD KHAN, A.K. 349-354

BACON, P.A. 551-553 BAGGER, J.P 33-40 BAHLMANN, J. 141-152 BALLARD, F.J. 315-320 BANDIERA, F. 137-140 BANKS, R.A. 471-474 BARNETT, D.B. 617-622 BARRADAS, M. 239-241 BASSENDINE, M.F. 643-648 BASTIAN, B.C. 593-599 BATTILANA, G. 259-263 BAZELMANS, J. 511-516 BEASTALL, G.H. 117-118 Веск, F. 183-186 BECK, F.W.J. 295-301 Beilin, L.J. 471-474 Bellini, G. 259-263 Bennett, T. 371-376, 377-382 Bergel, D.H. 455-461 Bessent, R.G. 527-535 BIANCHETTI, A. 259-263 BINGHAM, S. 629-635 BISTRIAN, B.R. 321-331 BLACKBURN, G.L. 321–331 BLAKE, D.R. 551-553 BLANN, A. 551-553 BLOMOVIST, C.G. 593-599 BODOLA, F. 481-486 BOER, P. 153-160 BORST, C. 581-586 Brod, J. 141–152 Вкоок, G. 637-642 Brown, I.M.H. 207-212 Brun-Pascaud, M. 497-504

CACHOVAN, M. 141-152

Burston, D. 433-439

BURGOYNE, J.L. 315-320

CAMPANACCI, L. 259-263 CARMEL, R. 193-205 CARRETTA, R. 259-263 CHAMBERLAIN, M.J. 69-78 CHATELAIN, R.E. 355-358 CICLITIRA, P.J. 655-659 CLARK, F. 41-47 CLARK, T.J.H. 555-557 CLAYTON, R.N. 1-6 CLUGSTON, G.A. 231-233 COADE, S.B. 333-340 Cook, D.B. 41-47 CORNISH, K.G. 281-287 CORRALL, R.J.M. 49-53 CRAFT, I.L. 239-241 CROSSLEY, I.R. 247-252 CUMBERBATCH, M. 167-176, 441-447 CUMMINGS, J.H. 629-635

DALTON, M. 307-314 DALTON, N. 475-479 DANDONA, P. 239-241 DANIELSON, B.G. 399-405 DAVIDSON, N.M. 49-53 DE GROOTE, J. 85-90 DESAI, S.P. 321-331 Dessi'-Fulgheri, 137-140 DONOHOE, K. 601-609 DORHOUT MEES, E.J. 153-160 DORNHORST, A.C. 661-662 Doyle, A.E. 463-470 DUFFY, S.G. 463-470 DULFANO, M.J. 449-451 DULL, W.L. 25-31 DUNNING, A.J. 581-586

EDDLESTON, A.L.W.F. 113-116
EDWARDS, R.H.T. 55-62, 547-550
ELANDER, B. 423-431
ELIA, M. 517-526

FABRIS, B. 259-263 FARR, M. 551-553 FARRI, L. 611-616 FAVRE, L. 407-415 FELL, G.S. 527-535 Fellenius, E. 423–431 Fellström, B. 399–405 Fern, E.B. 101–108 Feuer, G. 303–306 Fevery, J. 85–90 Flecknell, P.A. 161–165 Fois, G. 137–140 Forbes, G.B. 601–609 Franks, R.C. 481–486 French, E.B. 49–53 Frier, B.M. 49–53 Fröseth, B. 453–454

GALTON, D.J. 559-563 GARDNER, M.L.G. 433-439 GARDINER, I.T. 487-495 S.M. 371-376, GARDINER. 377-382 GARLICK, P.J. 101-108, 231-233 GEERS, A.B. 153-160 GERTZ, E.W. 573-580 GIBSON, G.J. 487-495 GILMORE, J.P. 281-287 GIRARD, P. 497-504 GLASS, R.E. 101-108 GLASSON, PH. 407-415 GLORIOSO, N. 137-140 GOLDBLATT, D. 601-609 GORDON, D. 273-280 Graham, R.M. 593-599 Grainger, S.L. 207-212 GRIBBIN, H.R. 487-495 GRIGGS, R.C. 601-609 GUTHRIE, G. 349-354 GUTTERIDGE, J.M.C. 551-553

Haglund, U. 423–431 Haines, A.J. 113–116 Hall, G.M. 109–111 Halliday, D. 243–246 Halliwell, B. 551–553, 649–653 Harkness, R.A. 333–340 Hayashi, R.H. 481–486 Heinz III, G.J. 487–495 Helander, H.F. 423–431 Henningsen, P. 33–40 Hervey, G.R. 7–18 Höck, S.A. 505–510 iv Author Index

HOPKINS, W.M. 49-53 HOWELL, J.B.L. 417-421 HUTTON, R. 239-241

INSEL, P.A. 265–272 IMESON, J.D. 273–280 IQBAL, M.J. 307–314

Jäppinen, P. 187–191 John, M. 161–165 Johnson, R.H. 587–591 Johnston, D.G. 41–47 Johnston, H.H. 349–354 Jones, P.B.B. 387–394, 395–398

Karlström, B. 399–405 Kaufman, A.M. 565–572 Keeling, P.W.N. 207–212 Kellen, J.A. 303–306 Khan, T. 565–572 Khokher, M. 239–241 Koomans, H.A. 153–160 Krølner, B. 537–540, 541–546 Krück, F. 505–510

Lambie, D.G. 587–591
Lane, D.J. 213–222
Lazarowitz, V.C. 573–580
Leach, W. 573–580
Lennox, E.S. 655–659
Lewis, C.M. 289–293
Lewis, S.F. 593–599
Lindblad, B.S. 433–439
Lithell, H. 399–405
Livesey, G. 517–526
Ljunghall, S. 399–405
Lofts, F.J. 63–68
Luk, C.K. 449–451
Lundholm, K. 243–246

Mac-Moune, F.L. 565-572
Madeddu, P. 137-140
Marcer, D. 417-421
Marigold, J.H. 207-212
Martin, V.I. 295-301
Matthews, D.M. 433-439
Maury, C.P. J. 453-454
McCulloch, A.J. 41-47
McFarlane, B.M. 113-116
McFarlane, I.G. 113-116, 127-135
Meade, T.W. 273-280
Meloni, F. 137-140
Millar, J.G.B. 623-627

MILLS, K.R. 55-62

MILLS, P.R. 527-535
MILLWARD, D.J. 243-246
MIR, M.A. 79-83
MOLDAWER, L.L. 321-331
MOORE, P.K. 63-68
MORGAN, D.B. 167-176,
177-182, 441-447
MORGAN, W.K.C. 69-78
MORTON, J.J. 359-370
MOTULSKY, H.J. 265-272
MOXHAM, J. 547-550
MOXLEY, R.T. 601-609
MURACA, M. 85-90
MURRAY, D.J. 341-347

NARBED, P.G. 417–421 NELSON, L.M. 527–535 NESTEL, P.J. 511–516 NEWHAM, D.J. 55–62 NEWHAM, D. 547–550 NIELSEN, T.T. 33–40 NOLAN, C. 511–516 NOSADINI, R. 41–47 NOWOTNY, P. 383–386

O'CONNOR, D.T. 265–272 Olbe, L. 423–431 O'MALLEY, B.P. 617–622 ÖSTMAN, J. 235–237

PALMER, K.T. 587-591 PALOMBO, J.D. 321-331 Papageorgiou, A. 611-616 PARK, R. 573-580 PATERSON, J.L. 109-111 PEART, W.S. 273-280 PENCHARZ, P.B. 611-616 PENFOLD, J.L. 315-320 Pernet, A. 41-47 PETERS, T.J. 341-347 PETTINGER, W.A. 593-599 Peveler, R.C. 455-461 PINIEWSKA, M. 41-47 POCIDALO, J.-J. 497-504 Polak, A. 623-627 Polu, J.M. 25-31 Pors Nielsen, S. 541-546 Pretschner, P. 141-152 PRIDE, N.B. 487-495

Quigley, B.M. 55-62

RAPPELLI, A. 137-140 REES, A. 559-563 RENNER, I.G. 193-205 RENNIE, M.J. 243-246 RENTHAL, R.D. 481-486 RIONDEL, A. 407–415
ROBINSON, B.J. 587–591
ROBINSON, J.L. 455–461
ROOS, J.C. 153–160
ROSENTHAL, F.D. 617–622
ROSHANAI, F. 91–99
ROTHWELL, N.J. 19–23
ROWLEY, D.A. 649–653
RUBYTHON, E.J. 177–182, 441–447
RUSSELL, R.G.G. 387–394, 395–398
RUSSELL, R.I. 527–535

SADOUL, P. 25-31 **SAKAMOTO, A.** 321-331 SANDERS, T.A.B. 91-99 SAVOLAINEN, H. 187-191 SAWERS, R.S. 307-314 SCHLEBUSCH, H. 505-510 SEID, J.M. 387–394 SEMPLE, P.D.'A. 117-118 SHERLOCK, S. 643-648 SIMMONDS, R.J. 333-340 SLEIGHT, P. 455-461 SOLTYS, J. 471-474 SORGER, M. 505-510 Sowers, J. 183-186 Sowers, J.R. 295-301 SPENCER, E. 417-421 STERN, N. 183-186 STEWART, R.I. 289-293 STIEL, D. 341-347 STOCK, M.J. 19-23 STOCKLEY, R.A. 119–126, 223-230 STOCKS, J. 559-563 STRADLING, J.R. 213-222

Таката, Ү. 463-470 TAYLOR, W.F. 593-599 TENHUNEN, R. 187-191 Teppo, A.-M. 453-454 THOMAS, H.C. 643-648 THOMASSEN, A.R. 33-40 THOMPSON, R.P.H. 207-212 Timson, L. 617-622 TOBIN, G. 7-18 Тогт, В. 537-540, 541-546 Tomas, T.M. 315-320 Tøndevold, E. 541-546 Tonolo, G. 137-140 TONUTTI, L. 259-263 Traynor, C. 109-111 TRUELOVE, S.C. 349-354

Author Index v

Vallotton, M.B. 407–415 Van Brederode, J.F.M. 581–586 Van Dongen Torman, M.A. 581–586 Van Montfrans, G.A. 581–586 Veroni, M. 463–470 Vessby, B. 399–405 Vierhapper, H. 383–386

WALDHÄUSL, W. 383–386 WALLACE, E.C.H. 359–370 WALLER, D.G. 623–627 WALLMARK, B. 423–431

VINITSKI, S. 69-78

WALLMARK, B. 423–431 WARDLE, E.N. 247–252 WARREN, J.B. 475–479 WATSON, W.S. 117–118 WEGELIUS, O. 453–454 Wieling, W. 581–586 Wiles, C.M. 547–550 Williams, R. 113–116, 247– 252 Williamson, D.H. 349–354 Winterstein, G. 637–642 Wootton, R. 161–165 Wright, N.A. 643–648

Volume 64

SUBJECT INDEX

First and last page numbers of papers to which entries refer are given.

Page numbers marked with an asterisk refer to Editorial Reviews.

Absorption, intestinal 433-439, 527-535	p-Aminobenzoic acid, 24 h urine collections
Acetylsalicylic acid, 6-ketoprostaglandin F _{1α}	629–625
excretion 395–398	δ-Aminolaevulinic acid dehydratase 187–191
Acid activation, renin 481-486	δ -Aminolaevulinic acid synthase 187–191
Acid excretion, renal 565-572	Aminophylline, adductor pollicis fatigue 547-
Acid secretion, isolated oxyntic glands 423-431	550
Acylglycerol metabolism, adipose tissue 235-237	Ammonia excretion 101–108
Adductor pollicis, aminophylline 547-550	hepatic encephalopathy 247-252*
Adenosine diphosphate, placenta 239-241	Amniotic fluid, renin activation 481-486
Adenosine 3',5'-phosphate	Amyloidosis, secondary 453-454
parathyroid hormone 623-627	Amyotrophic lateral sclerosis, glucose tolerance
vascular, renal hypertension 355-358	601–609
Adenosine triphosphatase	Angiotensin II
duodenal mucosa 341–347	dietary sodium 295–301
erythrocyte sodium efflux 79–83	experimental hypertension 359-370
hepatic encephalopathy 247-252*	insulin 383–386
Adipose tissue	Antigens, tissue 113–116
brown 7-18, 19-23	a_1 -Antitrypsin
nicotinic acid 235–237	cigarette smoke 223–230
Adrenalectomy, baroreflexes 371-376	reactive systemic amyloidosis 453-454
Adrenaline, exercise 475–479, 593–599	Apolipoproteins 559-563*
α ₂ -Adrenergic receptors, platelets 265-272	Arrhythmias, digitalis intoxication 253-258*
Adrenocorticotropic hormone, dietary sodium 295–301	Arterial hypertrophy, renal hypertension 355-358
Airflow, intravenous ethanol 555-557	Arterio-coronary sinus differences 33-40
Airflow obstruction, chronic 487–495	Artery
Airway conductance, adrenaline 475–479	ADP degradation 239-241
Alanine	distensibility 455–461
myocardial exchange 33-40	Ascorbate oxidase, synovial fluid 551-553
protein metabolism 517–526	Ascorbate, rheumatoid disease 649-653
Albumin, sex hormone binding 307–314	Asialoglycoproteins, hepatic clearance 127-
Alcoholic cirrhosis, zinc metabolism 527–535	135*
Alcohol dehydrogenase, hepatic zinc 527-535	Aspirin see Acetylsalicylic acid
Aldosterone	Asthma, intravenous ethanol 555-557
diuretics 407–415	Athymic mice, human hepatocellular carcinoma
erythrocyte sodium-potassium pump 183- 186	643–648
insulin 383–386	Atrial pacing 33–40
Alkaline phosphatase, duodenal mucosa 341–	Atropine, static and dynamic handgrip 593-
347	599
Alkalosis	Autoimmunity 113–116
frusemide 565–572	Autonomic nervous system
potassium depletion 497–504	failure 587–591 hypoglycaemia 49–53
Almitrine, pulmonary haemodynamics 25–31	
Amino acids	Autonomic neuropathy, heart rate 581-585 Azo reductase, tissue activity 349-354
branched chain 321-331, 517-526	AZO reductase, ussue activity 343-334
protein metabolism 101–108, 231–233, 517–	·
526	Baroreceptors, arterial 455-461
	

Baroreflexes	Carbonic anhydrase, duodenal mucosa 341-
essential hypertension 259–263	347
post-adrenalectomy hypotension 371-376	Cardiac output
Bicarbonate, duodenal secretion 341-347	determination 289–293
[14C]Bicarbonate, respiratory 14CO ₂ 231-233	static and dynamic handgrip 593-599
Biliary cirrhosis 113–116	Carotid sinus radius, phenylephrine 455-
Bilirubin	461
biliary excretion 85-90	Catecholamines
hepatic conjugation 85-90	exercise 475-479
Bilirubin UDP-glucuronosyltransferase 85-90	orthostatic hypotension 587-591
Blood flow	Cell population kinetics, hepatocellular car-
cerebral 161-165	cinoma 643-648
coronary 33-40	Chest cage restriction, resistive load detection
meclofenamate 471–474	417–421
Blood platelets, α_2 -adrenergic receptors, 265–272	Cholesterol, plasma 91-99, 637-642
Blood pressure	Chronic obstructive pulmonary disease 213—
adrenaline 475–479	222
autonomic blockade 593–599	Cigarette smoke, α_1 -antitrypsin 223–230
autonomic failure 587–591	
insulin 383–386	Cilia, beating frequency 449-451
	Circadian rhythm, 18-hydroxycorticosterone 295–301
plasma renin 273–280	
post-adrenalectomy 371–376	Citrate, myocardial exchange 33–40 Cobalamin, pancreatic juice 193–205
Blood vessels	
ADP degradation 239–241	Coeliac disease, gliadins radioimmunoassay
collagen 355-358	655–659
coronary artery disease 33–40	Collagen, vascular, renal hypertension 355-358
distensibility 455–461	Converting enzyme inhibition 359–370
meclofenamate 471-474	Coproporphyrinogen oxidase 187–191
Blood volume	Coronary artery disease 33-40
regulation 281–287	Cor pulmonale 117–118
renal disease 141-152	Creatinine
Body fluids, distribution 153–160	24 h urine collection 629-635
Body temperature, nerve conduction 617–622	hypopituitary children 315-320
Body mineral content	Cyclic AMP see Adenosine 3',5'-phosphate
bed rest 537-540	Cysteamine, duodenal ulcers 341-347
physical exercise 541–546	Cytosol, duodenal mucosal enzymes 341–347
Brain, blood flow 161–165	
Braking effect, diuretics 565-572	Deoxyribonucleic acid, vascular, renal hyper-
Branched-chain amino acids	tension 355-358
metabolic fate 517-526	Diabetic autonomic neuropathy 581-585
protein kinetics 321–331	Diet-induced thermogenesis 7–18, 19–23
Brattleboro rats, short-term isolation 377–382	Diflunisal, diuretics 407-415
Bronchodilatation, intravenous ethanol 555-	Digitalis intoxication 253-258*
557	Diuretics
Brown adipose tissue 7–18, 19–23	diflunisal 407–415
Brush border, duodenal mucosal enzymes 341-	essential hypertension 259-263
347	frusemide 565-572
	renal prostaglandins 407-415
	'Down-regulation', α_2 -adrenergic receptors
Caeruloplasmin, synovial fluid 551-553	265–272
Captopril, renal hypertension 463–470	Duodenal ulcer, mucosal enzymes 341–347
Carbon dioxide	Dyspnoea, resistive load detection 417–421
almitrine 25–31	~ j = j = j = j = j = j = j = j = j = j
[14C]bicarbonate infusion 231–233	Elastase
cardiac output rebreathing technique 289–	α_1 -antitrypsin 223–230
293	inhibitory activity 453–454
ventilatory response 487–495	lung diseases 119–126*
Tommatory response To 1-433	rung discases 117-120

Electrolytes	Graft vs host disease 113-116
balance 377-382	Growth hormone, myofibrillar protein 315-320
leukaemic plasma 79–83	Growth retardation 161–165
Endopeptidases, lung diseases 119-126*	Gluten, dietary content 655-659
Endoplasmic reticulum, liver 303-306	Glycoproteins, hepatic clearance 127–135*
Endothelial cells, prostacyclin 387-394, 395-	
398	Haemodynamics
Energy balance, thermogenesis 7-18, 19-23	pulmonary 25–31
Energy metabolism, preterm infants 611–616	renal disease 141–152
Erythrocyte	Haemostasis
hyperthyroidism 441–447	acetylsalicylic acid 395–398
potassium 167–176, 177–182	
purine 333–340	polyunsaturated fatty acids 91–99
rubidium 183–186	Haem synthesis, sulphides 187–191
	Handgrip
sodium 79–83, 161–176, 177–182, 441–447	haemodynamic responses 593-599
Ethanol, asthma 555-557	heart rate 581-585
Exercise	Heart
catecholamines 475–479	cardiac output 289-293, 593-599
plasma adrenaline 475–479	digitalis intoxication 253-258*
purine transport and metabolism 333-340	failure 573–580
vertebral bone loss 541-546	ischaemic disease 273–280
Extracellular fluid volume 153–160	Heart failure, lactic acidosis 573-580
	Heart rate
False neurotransmitters 247–252*	adrenaline 475-479
Fatigue, muscle	autonomic neuropathy 581-585
aminophylline 547-550	muscle activity 581-585, 593-599
low-frequency 55-62	Hepatectomy, cardiovascular function 573-580
Fatty acids	Hepatic encephalopathy 247–252*
essential 91-99	Hepatic endoplasmic reticulum 303-306
myocardial exchange 33-40	Hepatocellular carcinoma, growth characteristics
Ferroxidase, synovial fluid 551-553	in athymic mice 643–645
Fick method, indirect 289–293	Hydrochlorothiazide, renal prostaglandins
Fluid balance, short-term isolation 377–382	407–415
a-Foetoprotein, tumour secretion 643-648	Hydrogen peroxide, hydroxyl radicals 649-653
Fractures, bone mineral 541-546	Hydroxyl radicals, rheumatoid disease 649–653
Frusemide	18-Hydroxycorticosterone, circadian rhythm
chloride 565–572	295–301
erythrocyte sodium transport 79–83	Hypercapnia, almitrine infusion 25–31
sodium 565–572	Hypercholesterolaemia, plasma exchange 637-
Cartain alanda asid assession 400 401	642
Gastric glands, acid secretion 423–431	Hyperlipidaemia, lipoprotein variants 559-563*
Gastroscopic biopsy, isolated oxyntic glands	Hypertension
423–431	a ₂ -adrenergic receptors 265-272
Globulin, sex hormone binding 307–314	baroreflexes 259-263
Glucose	experimental 359-370, 463-470
adrenaline 475-479	Goldblatt 359-370
myocardial exchange 33-40	plasma renin 273-280
tolerance test 601-609	renal 141-152, 355-358, 463-470
triglyceride metabolism 511-516	vasopressin 377–382
turnover, thyroid failure 41–47	Hyperthyroidism
Glutamate, myocardial exchange 33-40	erythrocyte sodium pumps 441-447
Glutamine, protein metabolism 517-526	glucose turnover 41–47
Glycine, colorectal tumours 101–108	Hypoglycaemia, autonomic reaction 49-53
Glycogen, abdominal surgery 109-111	Hypopituitarism, myofibrillar protein 315-320
Goldblatt hypertension, renin-angiotensin system	Hypotension
359–370	orthostatic 587–591
Gonadotrophin-releasing hormone 1-6*	post-adrenalectomy 371-376

Hypothyroidism	Liver disease
glucose turnover 41-47	alcoholic cirrhosis 527-535
nerve conduction 617-622	biliary cirrhosis 113-116
	cirrhosis, 3-methylhistidine 243-246
Immersion, sympathectomy responses 281–287	hepatic encephalopathy 247-252*
Immobilization, bone mineral content 537–540	hepatocellular carcinoma 643-648
Indocyanine green, hepatic extraction 207–212	Loaded breathing 417–421
Indomethacin, renal prostaglandins 407–415	Lung
Infant, preterm, protein turnover 611–616	haemodynamics 25-31
Injury, protein kinetics 321–331	hypoxaemia 213–222
Inspiratory muscle function 487–495	regional deposition of particles 69-78
Insulin	Lung disease
blood pressure 383-386	cardiac output determination 289-293
triglyceride metabolism 511-516	chronic airflow obstruction 487–495
Interstitial compliance 153–160	obstructive, hypoxaemia 213-222
Intestinal absorption, peptides 433-439	proteinase inhibitors 119-126*
Intrapleural pressure gradient 69-78	Lysosomes, duodenal mucosal enzymes 341-
Ionic strength, cilia beating frequency 449-451	347
Iron, caeruloplasmin 551-553	Luxuskonsumption 7–18, 19–23
Iron salts, hydroxyl radicals 649-653	
Isolation, cardiovascular and renal effects 377-382	Macaca fascicularis, blood volume homoeostasis 281–287
302	Mammary adenocarcinoma, experimental 303-306
Kaliuresis, frusemide 565-572	Meclofenamate, peripheral vasculature 471-
6-Ketoprostaglandin F _{1a} , acetylsalicylic acid 395–398	474
Kidney body fluid distribution 153-160	Mercaptans, hepatic encephalopathy 247–252* Metabolic acidosis 573–580
hypertension 141–152, 355–358, 463–470	Metabolic alkalosis, potassium depletion 497-
renal stone 399–405	504
renin 463–470	3-Methylhistidine, cirrhosis 243-246
10mm 405-470	Methylhistidine, colorectal tumours 101–108
•	3-Methylhistidine, excretion in preterm infants
Lactate, myocardial exchange 33-40	611–616
Lactic acidosis, cardiovascular system 573–580	N^{τ} -Methylhistidine, hypopituitary children
Leucine, protein metabolism 231–233, 517–526	315–320
Leucocytes 505 510	Metoprolol, static and dynamic handgrip 593-
potassium 505–510	599
purine concentrations 333–340	Milk, human 611–616
Lipid transport 559–563*	Mitochondria, duodenal mucosal enzymes 341-
Lipogenesis, nicotinic acid 235–237	347
Lipolysis, nicotinic acid 235–237	Muscle, skeletal
Lipoprotein	aminophylline 547–550
cholesterol 91–99	hypopituitarism 315–320
hypercholesterolaemia 637–642	low-frequency fatigue 55-62
molecular variants 559–563*	mass 315–320
plasma exchange 637–642	pain 55-62
Lithium therapy, parathyroid hormone 623–627	wasting, glucose tolerance 601-609
Liver	Muscle, smooth, arterial 455-461
acidaemia 573–580	Myeloid leukaemic blast cell, erythrocyte sodium
	efflux 79–83
circulation 207–212 glycogen in abdominal surgery 109–111	Myotonic dystrophy, glucose tolerance 601-
glycoprotein clearance 127–135*	609
hepatectomy 573–580	Nolonono mastadanesta de la constanta de la co
indocyanine green extraction 207–212	Naloxone, post-adrenalectomy hypotension 371-376
progesterone binding 303–306	
broportions and all 201-200	Natriuresis, frusemide 565-572

Neoplasm, protein turnover 101–108	Progesterone binding, liver 303–306
Nerve conduction 617–622	Prostacyclin
Nicotinic acid, acylglycerol metabolism 235-	acetylsalicylic acid 395-398
237	cultured endothelial cells 387-394
Nitrogen metabolism 101–108	Prostaglandins
Nocturnal hypoxaemia 213–222	acetylsalicylic acid 395-398
Noradrenaline	diuretics 407–415
exercise 475–479	platelets 63–68, 91–99
static and dynamic handgrip 593-599	renal blood flow 471–474
	Prostaglandin E ₂ , diuretics 407–415
Oedema, cor pulmonale 117–118	Prostaglandin $F_{2\alpha}$, diuretics 407–415
Oestradiol, binding to plasma proteins 307-314	Proteases, pancreatic juice 193-205
Orthostatic hypotension 587–591	Protein
Osteoporosis	biosynthesis 101–108
bed rest 537-540	kinetics 321–331
physical exercise 541-546	purine-rich diet 399–405
Ouabain, erythrocyte 79-83, 183-186	turnover 231–233, 611–616
6-Oxoprostaglandin E ₁ , platelet release 63-68	Proteinase inhibitors, lung diseases 119–126*
Oxygen saturation, obstructive pulmonary	Proteolysis, lung diseases 119–126*
disease 213–222	Pulmonary emphysema 119-126*
Oxyntic glands, acid secretion 423-431	Pupil size, hypoglycaemia 49-53
	Purine
Pain, muscle contractions 55-62	protein-rich diet 399–405
Pancreatic juice, human 193–205	transport and metabolism 333-340
Parathyroid hormone, lithium therapy 623-627	
Particles, inhalation and lung deposition 69–78	R3230 AC rat mammary adenocarcinoma
Peptides, small-intestinal absorption 433–439	303–306
pH, cilia beating frequency 449-451	Radioimmunoassay, α - and β -gliadins 655–659
Pharmacokinetic model 207–212	Receptors, hormone 1-6*
Phenformin, lactic acidosis 573-580	Rectum, tumours 101–108
Phenols, hepatic encephalopathy 247-252*	Renal hypertension
Phenylephrine, carotid sinus radius 455–461	arterial cyclic AMP 355-358
Pituitary gland 1-6*	renin-angiotensin system 463-470
Placenta artery, ADP degradation 239-241	Renal stone disease, purine-rich protein diet
Plasma exchange, hypercholesterolaemia 637–	399–405
642	Renin
Plasma membrane, duodenal mucosal enzymes	acid activation 481–486
341–347	diuretics 407–415
Plasma proteins, sex hormone binding 307–314	epidemiology 273–280
Platelets	experimental hypertension 359–370
fatty acids 91–99	inactive 481–486
6-oxoprostaglandin E ₁ -like substance 63-68	renal hypertension 463-470
plasma exchange in hypercholesterolaemia	trypsin-activatable 137-140
637–642	Renin-angiotensin system, renal hypertension
prostacyclin production 387–394	463–470
Posture heart rate 581–585	Respiratory compensation, metabolic alkalosis 497–504
lung particles 69–78	· · · · · · · · · · · · · · · · · · ·
vascular responses 661–662	Respiratory fuel selection 517–526 Rheumatoid arthritis 453–454, 551–553
Potassium	Rheumatoid disease, hydroxyl radicals 649–
	653
depletion 497–504	000 .
erythrocyte 167–176, 183–186 frusemide 565–572	Saline expansion 152 160
hypokalaemia 167–176, 177–182	Saline expansion 153–160 Salivation, hypoglycaemia 49–53
insulin 383–386	Sanvation, hypogrycaemia 49–55 Sex differences
leucocyte 505–510	bilirubin conjugation 85–90
total body 505–510	platelet prostaglandins 63–68
ioiai body 303-310	practice prostagranding 03-00

erythrocyte sodium 177-182

intestinal 433-439

Triamterine, renal prostaglandins 407-415 Sex hormone binding globulin 307–314 Triglyceride, insulin 511-516 Signal Detection Theory, inspiratory loads 417-421 Triglycerides, plasma 91–99 Skeletal muscle see Muscle, skeletal Trypsin-activatable renin 137–140 Small intestine, peptide absorption 433–439 Tyrosine kinetics 321-331 Smoking, plasma renin activity 273-280 Smooth muscle see Muscle, smooth Umbilical artery, ADP degeneration 239-241 Sodium Urate, renal stone disease 399-405 erythrocyte 79-83, 167-176, 177-182, 183-Urea, excretion 101-108 186, 441-447 Ureteric ligation, experimental hypertension excretion 463-470 463-470 frusemide 565-572 Urine 18-hydroxycorticosterone 295–301 24 h collection 629-635 Sodium-potassium pump, erythrocyte 183-186 purines 333-340 Spine, bone loss 537-540, 541-546 Urolithiasis, purine-rich protein diet 399-405 Sputum, α_1 -antitrypsin 223–230 Uroporphyrinogen decarboxylase 187–191 Standing, heart rate 581-585 Uroporphyrinogen synthase 187-191 Stomach, isolated oxyntic glands 423-431 Sulphapyridine, tissue and bacterial splitting 349-354 Vagus nerve, heart rate control 581-585 Sulphasalazine, tissue and bacterial splitting Vascular endothelium, prostacyclin 387-394, 349-354 395-398 Sulphides, haem synthesis 187-191 Vascular reactivity, essential hypertension 259-Superoxide, rheumatoid disease 649-653 263 Supersaturation, urolithiasis 399-405 Vasopressin Surgery, liver glycogen 109-111 chronic lithium therapy 623-627 Synovial fluid, caeruloplasmin 551-553 isolation-induced hypertension 377-382 Sweating, hypoglycaemia 49-53 renal prostaglandins 407-415 Sympathetic nervous system Ventilation carbon dioxide 487-495 cardiopulmonary afferent nerves 281–287 thermogenesis 7-18, 19-23 regional 69-78 Synovial fluid, rheumatoid disease 649-653 Vertebrae, bone mineral 537-540, 541-546 Viscosity, cilia beating frequency 449-451 Visual evoked responses, thyroid dysfunction Temperature, nerve conduction 617-622 617-622 Testosterone, binding to plasma proteins 307-Volume regulation, sympathetic nervous system 314 Thermogenesis, diet-induced 7-18, 19-23 281-287 Thyroid gland, failure 41-47 Thyrotoxicosis Wheat gliadin, radioimmunoassay 655–659 glucose turnover 41-47 Whole-body radioactivity, zinc metabolism nerve conduction 617-622 527-535 L-Thyroxine, nerve conduction 617-622 Tilt, heart rate 581-585 Xenograft, hepatocellular carcinoma 643-648 Transport

Zinc, alcoholic cirrhosis 527–535