

CONTENTS

Vol. 80, No. 1

January 1991

| | |
|--|-------|
| Guidance for authors | i-ix |
| EDITORIAL REVIEW | |
| Glucocorticoid-suppressible hyperaldosteronism: a paradigm of arrested adrenal zonation? <i>by</i> A. GANGULY | 1-7 |
| <hr/> | |
| Cyclosporin A and the humoral response to blood transfusion in the rat: definition of antigens which suppress alloantibody formation <i>by</i> D. J. PROPPER, M. C. JONES, K. N. STEWART, G. R. D. CATTO and D. A. POWER | 9-15 |
| Effect of plasma from patients with essential hypertension on vascular resistance in the isolated perfused rat kidney <i>by</i> J. BACHMANN, H. SCHLÜTER, W. STORKEBAUM, H. WITZEL, F. WESSELS and W. ZIDEK | 17-23 |
| Dose-response relationships for the effects of insulin on glucose and fat metabolism in injured patients and control subjects <i>by</i> A. A. HENDERSON, K. N. FRAYN, C. S. B. GALASKO and R. A. LITTLE | 25-32 |
| Changes in electrocardiographic patterns at different stages of Chagas' heart disease in rats <i>by</i> R. B. BESTETTI, L. Z. PINTO, E. G. SOARES, G. MUCCILLO and J. S. M. OLIVEIRA | 33-37 |
| Cardiovascular reflex responses in Bangladeshi and European subjects <i>by</i> M. A. RAHMAN, I. FARQUHAR and T. BENNETT | 39-45 |
| Effects of dichloroacetate and bicarbonate on haemodynamic parameters in healthy volunteers <i>by</i> B. LUDVIK, G. PEER, A. BERZLANOVICH, S. STIFTER and H. GRAF | 47-51 |
| Reduced plasma noradrenaline concentrations in simple-obese and diabetic obese patients <i>by</i> A. V. ASTRUP, N. J. CHRISTENSEN and L. BREUM | 53-58 |
| Mechanism of reserpine-induced acute gastric mucosal injury in the rat: role of endogenous 5-hydroxytryptamine <i>by</i> A. S. SALIM | 59-64 |
| Differentiation between the intensity of breathlessness and the distress it evokes in normal subjects during exercise <i>by</i> R. C. WILSON and P. W. JONES | 65-70 |
| Abnormal renal glucose handling in X-linked hypophosphataemic mice <i>by</i> R. C. MÜHLBAUER and H. FLEISCH | 71-76 |
| Limits of brain tolerance to daily increments in serum sodium in chronically hyponatraemic rats treated with hypertonic saline or urea: advantages of urea <i>by</i> A. SOUPART, A. STENUIT, O. PERIER and G. DECAUX | 77-84 |
| CORRESPONDENCE SECTION | |
| The ambiguity of changes in the rate constants of fluxes <i>by</i> I. M. GLYNN | 85-86 |

Vol. 80, No. 2

February 1991

| | |
|---|---------|
| Lymphocytotoxins in sera from highly sensitized multiparous dialysis patients: antibody class, relationship with the HLA and with paternal antigens <i>by</i> D. J. PROPPER, W. A. LEHENY, S. J. URBANIAK, G. R. D. CATTO and A. M. MACLEOD | 87-93 |
| Na ⁺ /H ⁺ antiport and buffering capacity in human polymorphonuclear and mononuclear leucocytes <i>by</i> V. FRIGHI, L. L. NG, A. LEWIS and H. DHAR | 95-99 |
| Platelet thromboxane A ₂ receptors in type I diabetes <i>by</i> P. A. MODESTI, R. ABBATE, G. F. GENSINI, A. COLELLA and G. G. NERI SERNERI | 101-105 |

| | |
|--|---------|
| Effect of theophylline and dipyridamole on the respiratory response to isocapnic hypoxia in normal human subjects <i>by</i> S. T. PARSONS, T. L. GRIFFITHS, J. M. L. CHRISTIE and S. T. HOLGATE | 107-112 |
| Serial changes in pulmonary haemodynamics during human pregnancy: a non-invasive study using Doppler echocardiography <i>by</i> S. C. ROBSON, S. HUNTER, R. J. BOYS and W. DUNLOP | 113-117 |
| Functional properties <i>in vitro</i> of systemic small arteries from rabbits fed a cholesterol-rich diet for 12 weeks <i>by</i> U. SIMONSEN, E. EHRNROOTH, L. U. GERDES, O. FAERGEMANN, J. BUCH, F. ANDREASEN and M. J. MULVANY | 119-129 |
| Differential effects of atrial natriuretic peptide and dopamine on urinary protein excretion in chronic glomerulonephritis <i>by</i> Y. HIRATA, M. ISHII, K. FUKUI, H. HAYAKAWA, E. SUZUKI, T. SUGIMOTO, K. KIMURA, H. MATSUOKA and T. SUGIMOTO | 131-136 |
| Abnormal erythrocyte choline transport in patients with chronic renal failure <i>by</i> F. C. FERVENZA, D. MEREDITH, J. C. ELLORY and B. M. HENDRY | 137-141 |
| Glomerular binding and contractile response to angiotensin II in rats with chronic experimental cirrhosis of the liver <i>by</i> L. M. VILLAMEDIANA, M. VELO, A. OLIVERA, L. HERNANDO, C. CARAMELO and J. M. LÓPEZ-NOVOA | 143-147 |
| Oral carbidopa has no effect on the renal response to angiotensin II in normal man <i>by</i> D. W. EADINGTON, C. P. SWAINSON and M. R. LEE | 149-154 |
| Indocyanine Green elimination in patients with liver disease and in normal subjects <i>by</i> E. BURNS, D. R. TRIGER, G. T. TUCKER and N. D. S. BAX | 155-160 |
| Tubular reabsorption of pepsinogen isozymogens in man studied by the inhibition of tubular protein reabsorption with dibasic amino acids <i>by</i> M. A. G. J. TEN DAM, A. ZWIERS, J. B. A. CRUSIUS, G. PALS, G. J. VAN KAMP, S. G. M. MEUWISSEN, A. J. M. DONKER and R. W. TEN KATE | 161-166 |
| Reassessment of the single intravenous injection method with inulin for measurement of the glomerular filtration rate in man <i>by</i> L. F. PRESCOTT, S. FREESTONE and J. A. N. MCAUSLANE | 167-176 |
| The contribution of the large intestine to blood acetate in man <i>by</i> W. SCHEPPACH, E. W. POMARE, M. ELIA and J. H. CUMMINGS | 177-182 |

Vol. 80, No. 3

March 1991

EDITORIAL REVIEW

| | |
|--|---------|
| Local control of human peripheral vascular tone: implications for drug therapy <i>by</i> N. BENJAMIN and P. VALLANCE | 183-190 |
| Orotidine accumulation in human erythrocytes during allopurinol therapy: association with high urinary oxypurinol-7-riboside concentrations in renal failure and in the Lesch-Nyhan syndrome <i>by</i> H. A. SIMMONDS, S. REITER, P. M. DAVIES and J. S. CAMERON | 191-197 |
| Fasting plasma glucose levels and endogenous androgens in non-diabetic post-menopausal women <i>by</i> K.-T. KHAW and E. BARRETT-CONNOR | 199-203 |
| Alterations of platelet membrane microviscosity in essential hypertension <i>by</i> K.-H. LE QUAN SANG, T. MONTENAY-GARESTIER and M.-A. DEVYNCK | 205-211 |
| Hepatic phosphofructokinase-1 activity and fructose 2,6-bisphosphate levels in patients with abdominal sepsis <i>by</i> J. ARNOLD, M. J. HAMER and M. IRVING | 213-217 |
| Renal and cardiovascular effects of exogenous insulin in healthy volunteers <i>by</i> R. O. B. GANS, L. v. D. TOORN, H. J. G. BILO, J. J. P. NAUTA, R. J. HEINE and A. J. M. DONKER | 219-225 |
| Cardiac and whole-body [³ H]noradrenaline kinetics during adrenaline infusion in man <i>by</i> A. J. McCANCE and J. C. FORFAR | 227-233 |

| | |
|--|---------|
| Secondary ischaemia in rabbit skin flaps: the roles played by thromboxane and free radicals <i>by</i> K. R. KNIGHT, M. F. ANGEL, D. A. LEPORE, P. A. ABBEY, L. I. ARNOLD, K. A. GRAY, C. G. MELLOW and B. MCC. O'BRIEN | 235-240 |
| Plasma exudation into airways induced by inhaled platelet-activating factor: effect of peptidase inhibition <i>by</i> J. O. LÖTVALL, W. ELWOOD, K. TOKUYAMA, P. J. BARNES and K. F. CHUNG | 241-247 |
| Influence of neuropathy on the microvascular response to local heating in the human diabetic foot <i>by</i> M. J. STEVENS, M. E. EDMONDS, S. L. E. DOUGLAS and P. J. WATKINS | 249-256 |
| Neuronal re-uptake of noradrenaline by sympathetic nerves in humans <i>by</i> G. EISENHOFER, M. D. ESLER, I. T. MEREDITH, C. FERRIER, G. LAMBERT and G. JENNINGS | 257-263 |
| Inhibition of neutral endopeptidase (EC 3.4.24.11) leads to an atrial natriuretic factor-mediated natriuretic, diuretic and antihypertensive response in rodents <i>by</i> N. B. SHEPPERSON, P. L. BARCLAY, J. A. BENNETT and G. M. R. SAMUELS | 265-269 |
| Diurnal blood pressure variation in quadriplegic chronic spinal cord injury patients <i>by</i> H. KRUM, W. J. LOUIS, D. J. BROWN, G. P. JACKMAN and L. G. HOWES | 271-276 |
| Supplement 24: Medical Research Society Communications | 1P-38P |

Vol. 80, No. 4

April 1991

EDITORIAL REVIEW

| | |
|--|---------|
| Stable isotopes in clinical research: safety reaffirmed <i>by</i> P. J. H. JONES and S. T. LEATHERDALE | 277-280 |
|--|---------|

RAPID COMMUNICATION

| | |
|--|---------|
| Achlorhydria-induced hypergastrinaemia: the role of bacteria <i>by</i> J. CALAM, R. A. GOODLAD, C. Y. LEE, B. RATCLIFFE, M. E. COATES, G. W. H. STAMP and N. A. WRIGHT | 281-284 |
| Transcription, storage and release of atrial natriuretic factor in the failing human heart <i>by</i> T. A. FISCHER, M. HAASS, R. DIETZ, R. C. WILLENBROCK, W. SAGGAU, R. E. LANG and W. KÜBLER | 285-291 |
| How important are suppression of aldosterone and stimulation of atrial natriuretic peptide secretion in the natriuretic response to an acute sodium load in man? <i>by</i> D. R. J. SINGER, D. G. SHIRLEY, N. D. MARKANDU, M. A. MILLER, M. G. BUCKLEY, A. L. SUGDEN, G. A. SAGNELLA and G. A. MACGREGOR | 293-299 |
| A study of thyroglobulin and peroxidase activity in the thyroid tissue of patients with non-endemic non-toxic nodular goitre <i>by</i> T. MORIES, J. M. MIRALLES, A. REGLERO, S. FELIPE, J. J. CORRALES and L. C. GARCIA | 301-307 |
| Long-term reproducibility of Borg scale estimates of breathlessness during exercise <i>by</i> R. C. WILSON and P. W. JONES | 309-312 |
| Triacylglycerol in the human erythrocyte: quantification and fatty acyl turnover in normal subjects and chronic alcoholics <i>by</i> B. LERIQUE, M. GASTALDI and J. BOYER | 313-318 |
| Total body calcium in patients with inflammatory bowel disease: a longitudinal study <i>by</i> S. J. S. RYDE, D. CLEMENTS, W. D. EVANS, R. MOTLEY, W. D. MORGAN, C. EVANS, J. RHODES and J. E. COMPSTON | 319-324 |
| Adrenoceptors on blood cells from patients with primary Raynaud's phenomenon <i>by</i> S. J. GRAAFSMA, H., WOLLERSHEIM, H. T. DROSTE, M. A. G. J. TEN DAM, L. J. H. VAN TITS, J. REYENGA, J. F. RODRIGUES DE MIRANDA and T. THIEN | 325-331 |
| Autonomic mechanisms underlying intraocular pressure changes during insulin-induced hypoglycaemia in normal human subjects: effects of pharmacological blockade <i>by</i> D. A. HEPBURN, B. M. FISHER, I. THOMSON, T. BARRIE and B. M. FRIER | 333-338 |
| Renal and extra-renal levels of renin mRNA in experimental hypertension <i>by</i> N. J. SAMANI, W. J. BRAMMAR and J. D. SWALES | 339-344 |

| | |
|--|---------|
| Measurement of whole-body protein turnover in insulin-dependent (type 1) diabetic patients during insulin withdrawal and infusion: comparison of [¹³ C]leucine and [² H ₃]phenylalanine methodologies <i>by</i> P. J. PACY, G. N. THOMPSON and D. HALLIDAY | 345-352 |
| Extent of pulmonary emphysema in man and its relation to the loss of elastic recoil <i>by</i> M. GUGGER, G. GOULD, M. F. SUDLOW, P. K. WRAITH and W. MACNEE | 353-358 |
| Basal fuel homeostasis in symptomatic human immunodeficiency virus infection <i>by</i> M. J. T. HOMMES, J. A. ROMIJN, E. ENDERT, J. K. M. EEF TINCK SCHATTENKERK and H. P. SAUERWEIN | 359-365 |
| Effects of contractile activity on muscle damage in the dystrophin-deficient <i>mdx</i> mouse <i>by</i> A. McARDLE, R. H. T. EDWARDS and M. J. JACKSON | 367-371 |
| Doxepin therapy for postprandial symptomatic hypoglycaemic patients: neurochemical, hormonal and metabolic disturbances <i>by</i> F. LECHIN, B. VAN DER DIJS, A. LECHIN, M. LECHIN, E. COLL-GARCÍA, H. JARA, A. CABRERA, V. JIMÉNEZ, F. GÓMEZ, D. TOVAR, I. RADA and L. AROCHA | 373-384 |
| Differential stimulation by oxygen-free-radical-altered immunoglobulin G of the production of superoxide and hydrogen peroxide by human polymorphonuclear leucocytes <i>by</i> H. A. KLEINVELD, W. SLUTTER, A. M. C. BOONMAN, A. J. G. SWAAK, C. E. HACK and J. F. KOSTER | 385-391 |
| Urinary excretion of nitrogenous and non-nitrogenous compounds in the chronic ethanol-fed rat <i>by</i> V. R. PREEDY, B. HAMMOND, R. A. ILES, S. E. C. DAVIES, J. D. GANDY, R. A. CHALMERS and T. J. PETERS | 393-400 |
| CORRESPONDENCE SECTION | |
| The kidney and essential hypertension <i>by</i> I. H. MILLS | 401-402 |
| ANNOUNCEMENT | |
| Enhancement of the pulmonary vasoconstriction reaction to alveolar hypoxia in systemic high blood pressure <i>by</i> M. D. GUAZZI, M. BERTI, E. DORIA, C. FIORENTINI, C. GALLI, M. PEPI and G. TAMBORINI (Clin. Sci. 1989; 76, 589-94) | 403 |

Vol. 80, No. 5

May 1991

EDITORIAL REVIEW

| | |
|---|---------|
| Role of proto-oncogenes in the control of myocardial cell growth and function <i>by</i> M. D. GAMMAGE and J. A. FRANKLYN | 405-411 |
| Role of the wrist cuff in forearm plethysmography <i>by</i> J. LENDERS, G.-J. JANSSEN, P. SMITS and T. THIEN | 413-417 |
| Hypothalamic regulatory peptides in obese and lean Zucker rats <i>by</i> G. WILLIAMS, H. M. CARDOSO, Y. C. LEE, J. M. BALL, M. A. GHATEI, M. J. STOCK and S. R. BLOOM | 419-426 |
| Influence of the plasma protein concentration on renal function <i>by</i> A. D. CUMMING and A. LINTON | 427-433 |
| Renal sensitivity to endothelium-derived-relaxing-factor-mediated vasodilatation in the spontaneously hypertensive rat <i>by</i> G. A. BURTON, J. HAYLOR and A. DE JONGE | 435-441 |
| Effects of treatment with nasal continuous positive airway pressure on atrial natriuretic peptide and arginine vasopressin release during sleep in patients with obstructive sleep apnoea <i>by</i> J. KRIEGER, M. FOLLENIUS, E. SFORZA, G. BRANDENBERGER and J. D. PETER | 443-449 |
| Clarification of the identity of the major M2 autoantigen in primary biliary cirrhosis <i>by</i> S. P. M. FUSSEY, S. M. WEST, J. G. LINDSAY, C. I. RAGAN, O. F. W. JAMES, M. F. BASSENDINE and S. J. YEAMAN | 451-455 |
| Skeletal muscle degradation and nitrogen wasting in rats with chronic metabolic acidosis <i>by</i> B. WILLIAMS, E. LAYWARD and J. WALLS | 457-462 |
| Effect of destruction of the vascular endothelium upon pressure/flow relations and endothelium-dependent vasodilatation in resistance beds of spontaneously hypertensive rats <i>by</i> M. D. RANDALL, G. R. THOMAS and C. R. HILEY | 463-469 |

| | |
|---|---------|
| Amino acid metabolism in human subcutaneous adipose tissue <i>in vivo</i> by K. N. FRAYN, K. KHAN, S. W. COPPACK and M. ELIA | 471-474 |
| Aldosterone, atrial natriuretic factor and sodium intake as determinants of the natriuretic response to head-out water immersion in healthy subjects by L. C. WARNER, G. A. MORALI, J. A. MILLER, A. G. LOGAN, K. L. SKORECKI and L. M. BLENDIS | 475-480 |
| Role of prostaglandins in the natriuresis of head-out water immersion in humans by T. J. RABELINK, H. A. KOOMANS and E. J. DORHOUT MEES | 481-488 |
| Plasma profiles and removal rates of inorganic sulphate, and their influence on serum ionized calcium, in patients on maintenance haemodialysis by M. MARANGELLA, M. PETRARULO, D. COSSEDDU, C. VITALE and F. LINARI | 489-495 |
| Enhanced urinary excretion of leukotriene E ₄ by patients with multiple trauma with or without adult respiratory distress syndrome by J. FAULER, D. TSIKAS, M. HOLCH, A. SEEKAMP, M. L. NERLICH, J. STURM and J. C. FRÖLICH | 497-504 |
| Ranitidine suppresses aluminium absorption in man by R. S. C. RODGER, G. S. MURALIKRISHNA, D. J. HALLS, J. B. HENDERSON, J. A. H. FORREST, A. I. MACDOUGALL and G. S. FELL | 505-508 |
| Evidence for altered Na ⁺ /H ⁺ antiport activity in cultured skeletal muscle cells and vascular smooth muscle cells from the spontaneously hypertensive rat by J. E. DAVIES, L. L. NG, M. AMEEN, P. D. SYME and J. K. ARONSON | 509-516 |
| Deglycosylation of α ₁ -proteinase inhibitor is impaired in the faeces of patients with active inflammatory bowel disease (Crohn's disease) by C. MIZON, J. EL YAMANI, J.-F. COLOMBEL, P. MAES, M. BALDUYCK, A. LAINE, A. CORTOT, A. TARTAR and J. MIZON | 517-523 |
| Effects of acute insulin-induced hypoglycaemia on haemostasis, fibrinolysis and haemorheology in insulin-dependent diabetic patients and control subjects by B. M. FISHER, J. D. QUIN, A. RUMLEY, S. E. LENNIE, M. SMALL, A. C. MACCUISH and G. D. O. LOWE | 525-531 |

Vol. 80, No. 6

June 1991

EDITORIAL REVIEW

| | |
|--|---------|
| Working hypotheses for the investigation of clinical hypertension by M. E. SAFAR | 533-538 |
| Intracellular free magnesium in human lymphocytes and the response to lectins by L. L. NG, J. E. DAVIES and M. C. GARRIDO | 539-547 |
| A cross-sectional study of platelet cyclic AMP in healthy and hypertensive pregnant women by E. H. HORN, J. COOPER, S. HEPTINSTALL and P. C. RUBIN | 549-558 |
| Glutathione depletion during experimental damage to rat skeletal muscle and its relevance to Duchenne muscular dystrophy by M. J. JACKSON, M. H. BROOKE, K. KAISER and R. H. T. EDWARDS | 559-564 |
| Hypotension induced by intravascular administration of nerve growth factor in the rat by Q. YAN, S. L. SETTLE and M. R. WILKINS | 565-569 |
| Energy expenditure and substrate metabolism measured by 24 h whole-body calorimetry in patients receiving cyclic and continuous total parenteral nutrition by E. PULLICINO, G. R. GOLDBERG and M. ELIA | 571-582 |
| Increased symptoms of hypoglycaemia in the standing position in insulin-dependent diabetes mellitus by I. B. HIRSCH, S. R. HELLER and P. E. CRYER | 583-586 |
| Measurement of tumour protein synthesis <i>in vivo</i> in human colorectal and breast cancer and its variability in separate biopsies from the same tumour by S. D. HEYS, K. G. M. PARK, M. A. McNURLAN, A. G. CALDER, V. BUCHAN, K. BLESSING, O. EREMIN and P. J. GARLICK | 587-593 |
| Noradrenaline release in skeletal muscle and in adipose tissue studied by microdialysis by B. GRÖNLUND, A. ASTRUP, P. BIE and N. J. CHRISTENSEN | 595-598 |

| | |
|---|---------|
| Na ⁺ ,K ⁺ -ATPase activity and hormones in single nephron segments from nephrotic rats <i>by</i> B. VOGT and H. FAVRE | 599-604 |
| Early loss of neurogenic inflammation in the human diabetic foot <i>by</i> D. WALMSLEY and P. G. WILES | 605-610 |
| Increased blood antioxidant systems of runners in response to training load <i>by</i> J. D. ROBERTSON, R. J. MAUGHAN, G. G. DUTHIE and P. C. MORRICE | 611-618 |
| Twenty-four hour intragastric acidity and plasma gastrin concentration profiles in female and male subjects <i>by</i> E. J. PREWETT, J. T. L. SMITH, C. U. NWOKOLO, A. M. SAWYERR and R. E. POUNDER | 619-624 |
| Effect of α -ketoglutarate infusions on organ balances of glutamine and glutamate in anaesthetized dogs in the catabolic state <i>by</i> E. ROTH, J. KARNER, A. ROTH-MERTEN, S. WINKLER, L. VALENTINI and K. SCHAUPP | 625-631 |
| Resting energy expenditure in chronic cardiac failure <i>by</i> M. RILEY, J. S. ELBORN, W. R. MCKANE, N. BELL, C. F. STANFORD and D. P. NICHOLLS | 633-639 |