467-478

Clinical SCIENCE

COMMENTS	
Does oxidized low-density lipoprotein occur in vivo? by M. J. Jackson (see accompanying paper pp. 489–495)	393
Cerebrovascular reactivity measurement using near-infrared spectroscopy by D. Grosset (see accompanying paper pp. 497–504)	395
PAPERS	
Importance of glutamine metabolism in murine macrophages and human monocytes to L-arginine biosynthesis and rates of nitrite or urea production	207 407
by C. Murphy and P. Newsholme	397–407
Renal arginine metabolism in fasted rats with subacute short bowel syndrome by C. H. C. Dejong, C. F. M. Welters, N. E. P. Deutz, E. Heineman and P. B. Soeters	409–418
Studies in vivo of ω -gliadins in gluten sensitivity (coeliac sprue disease) by A. Ensari, M. N. Marsh, K. J. Moriarty, C. M. Moore, R. J. Fido and A. S. Tatham	419–424
Isolation and characterization of the viscous, high-molecular-mass microbial carbohydrate fraction from faeces of healthy subjects and patients with Crohn's disease and the consequences for a therapeutic approach	
by J. G. H. Ruseler-van Embden, L. M. C. van Lieshout, D. J. Binnema and M. P. Hazenberg	425–433
Inhibition of renal II β -hydroxysteroid dehydrogenase <i>in vivo</i> by carbenoxolone in the rat and its relationship to sodium excretion	
by K. J. Sewell, D. G. Shirley, A. E. Michael, A. Thompson, D. P. Norgate and R. J. Unwin	435–443
A comparison of the development of renal hypertension in male and female rats by R. Okuniewski, E.A. Davis, B. Jarrott and R. E. Widdop	445–451
Arrhythmogenic effect of ventriculography in patients with left ventricular dilation and/or hypertrophy	
by A. S. Serafi, S. J. Evans and J. V. Jones	453–458
Administration of albumin to patients with sepsis syndrome: a possible beneficial role in plasma thiol repletion	
by G. J. Quinlan, M. P. Margarson, S. Mumby, T. W. Evans and J. M. C. Gutteridge	459–465
Porcine hepatic response to sepsis and its amplification by an adrenergic receptor a_1 agonist and a β_2 antagonist	

by D. Tighe, R. Moss and D. Bennett

Randomized clinical trial comparing the effectiveness of two dietary interventions for	
patients with hyperlipidaemia	
by C. Ní Mhurchú, B. M. Margetts and V. Speller	479–487
Identification of oxidized low-density lipoprotein in human serum by NMR spectroscopy	
by J. Jankowski, JR. Nofer, M. Tepel, B. Griewel, H. Schlüter, G. Assmann and W. Zidek	489–495
Evaluation of potential factors affecting the measurement of cerebrovascular reactivity by near-infrared spectroscopy	
by R. Totaro, G. Barattelli, V. Quaresima, A. Carolei and M. Ferrari	497–504
Contractile response of isolated human hepatic arteries to a -adrenoceptor agonist is not impaired in patients with cirrhosis	
by P.W. F. Hadoke, J. F. Dillon, T. G. John, S. W. Walker, P. C. Hayes and B. C. Williams	505–511
Inhaled corticosteroid therapy reduces the early morning peak in cortisol and aldosterone	
by A. M. Wilson, F. I. Sims, A. D. Struthers and R. I. I. inworth	513-517